

**WO0140276**

Publication Title:

NOVEL SERPENTINE TRANSMEMBRANE ANTIGENS EXPRESSED IN HUMAN CANCERS AND USES THEREOF

Abstract:

Described is a novel family of cell surface serpentine transmembrane antigens. Two of the proteins in this family are exclusively or predominantly expressed in the prostate, as well as in prostate cancer, and thus members of this family have been termed "STEAP" (<u>S</u>ix <u>T</u>ransmembrane <u>E</u>pithelial <u>A</u>ntigen of the <u>P</u>rostate). Four particular human STEAPs are described and characterized herein. The human STEAPs exhibit a high degree of structural conservation among them but show no significant structural homology to any known human proteins. The prototype member of the STEAP family, STEAP-1, appears to be a type IIIa membrane protein expressed predominantly in prostate cells in normal human tissues. Structurally, STEAP-1 is a 339 amino acid protein characterized by a molecular topology of six transmembrane domains and intracellular N- and C- termini, suggesting that it folds in a "serpentine" manner into three extracellular and two intracellular loops. STEAP-1 protein expression is maintained at high levels across various stages of prostate cancer. Moreover, STEAP-1 is highly over-expressed in certain other human cancers.

-----  
Data supplied from the esp@cenet database - <http://ep.espacenet.com>

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 June 2001 (07.06.2001)

PCT

(10) International Publication Number  
**WO 01/40276 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/00**

(21) International Application Number: PCT/US00/33040

(22) International Filing Date: 6 December 2000 (06.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/455,486 6 December 1999 (06.12.1999) US

(71) Applicant: **UROGENESYS, INC.** [US/US]; 1701 Colorado Avenue, Santa Monica, CA 90404 (US).

(72) Inventors: **AFAR, Daniel, E., H.**; 435 Visitacion Avenue, Brisbane, CA 94005 (US). **HUBERT, Rene, S.**; 1644 North Occidentia Boulevard, Los Angeles, CA 90026 (US). **RAITANO, Arthur, B.**; 10807 Cushdon Avenue, Los Angeles, CA 90064 (US). **SAFFRAN, Douglas, C.**; 11720 Bellagio Road, No. 204, Los Angeles, CA 90049 (US). **MITCHELL, Steve, Chappell**; 2417 1/2

4th Street, Santa Monica, CA 90405 (US). **FARIS, Mary**; 2538 Almaden Court, Los Angeles, CA 90077 (US). **JAKOBOVITS, Aya**; 3135 Hutton Drive, Beverly Hills, CA 90210 (US).


(74) Agent: **CANADY, Karen, S.**; Gates & Cooper LLP, Suite 1050, 6701 Center Drive West, Los Angeles, CA 90045 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

[Continued on next page]

(54) Title: NOVEL SERPENTINE TRANSMEMBRANE ANTIGENS EXPRESSED IN HUMAN CANCERS AND USES THEREOF



(57) Abstract: Described is a novel family of cell surface serpentine transmembrane antigens. Two of the proteins in this family are exclusively or predominantly expressed in the prostate, as well as in prostate cancer, and thus members of this family have been termed "STEAP" (Six Transmembrane Epithelial Antigen of the Prostate). Four particular human STEAPs are described and characterized herein. The human STEAPs exhibit a high degree of structural conservation among them but show no significant structural homology to any known human proteins. The prototype member of the STEAP family, STEAP-1, appears to be a type IIIa membrane protein expressed predominantly in prostate cells in normal human tissues. Structurally, STEAP-1 is a 339 amino acid protein characterized by a molecular topology of six transmembrane domains and intracellular N- and C- termini, suggesting that it folds in a "serpentine" manner into three extracellular and two intracellular loops. STEAP-1 protein expression is maintained at high levels across various stages of prostate cancer. Moreover, STEAP-1 is highly over-expressed in certain other human cancers.



WO 01/40276 A2



IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

— *Without international search report and to be republished upon receipt of that report.*

**NOVEL SERPENTINE TRANSMEMBRANE ANTIGENS**  
**EXPRESSED IN HUMAN CANCERS AND USES THEREOF**

This application claims the benefit of United States patent application number 09/455,486, filed December 6, 1999.

**FIELD OF THE INVENTION**

5           The invention described herein relates to a family of novel genes and their encoded proteins and tumor antigens, termed STEAPs, and to diagnostic and therapeutic methods and compositions useful in the management of various cancers, particularly including prostate cancer, colon cancer, bladder cancer, lung cancer, ovarian cancer and pancreatic cancer.

10

**BACKGROUND OF THE INVENTION**

Cancer is the second leading cause of human death next to coronary disease. Worldwide, millions of people die from cancer every year. In the United States alone, cancer causes the death of well over a half-million people annually, with some 1.4 million  
15 new cases diagnosed per year. While deaths from heart disease have been declining significantly, those resulting from cancer generally are on the rise. In the early part of the next century, cancer is predicted to become the leading cause of death.

Worldwide, several cancers stand out as the leading killers. In particular, carcinomas of the lung, prostate, breast, colon, pancreas, and ovary represent the primary  
20 causes of cancer death. These and virtually all other carcinomas share a common lethal feature. With very few exceptions, metastatic disease from a carcinoma is fatal. Moreover, even for those cancer patients who initially survive their primary cancers, common experience has shown that their lives are dramatically altered. Many cancer patients experience strong anxieties driven by the awareness of the potential for  
25 recurrence or treatment failure. Many cancer patients experience physical debilitations following treatment. Many cancer patients experience a recurrence.

Worldwide, prostate cancer is the fourth most prevalent cancer in men. In North America and Northern Europe, it is by far the most common male cancer and is the second leading cause of cancer death in men. In the United States alone, well over 40,000 men die annually of this disease - second only to lung cancer. Despite the  
5 magnitude of these figures, there is still no effective treatment for metastatic prostate cancer. Surgical prostatectomy, radiation therapy, hormone ablation therapy, and chemotherapy continue to be the main treatment modalities. Unfortunately, these treatments are ineffective for many and are often associated with undesirable consequences.

10 On the diagnostic front, the lack of a prostate tumor marker that can accurately detect early-stage, localized tumors remains a significant limitation in the management of this disease. Although the serum PSA assay has been a very useful tool, its specificity and general utility is widely regarded as lacking in several important respects.

Progress in identifying additional specific markers for prostate cancer has been  
15 improved by the generation of prostate cancer xenografts that can recapitulate different stages of the disease in mice. The LAPC (Los Angeles Prostate Cancer) xenografts are prostate cancer xenografts that have survived passage in severe combined immune deficient (SCID) mice and have exhibited the capacity to mimic disease progression, including the transition from androgen dependence to androgen independence and the  
20 development of metastatic lesions (U.S. Patent No. 6,107,540; Klein et al., 1997, Nat. Med.3:402). More recently identified prostate cancer markers include PCTA-1 (Su et al., 1996, Proc. Natl. Acad. Sci. USA 93: 7252), prostate stem cell antigen (PSCA) (Reiter et al., 1998, Proc. Natl. Acad. Sci. USA 95: 1735), and STEAP (Hubert et al., 1999, Proc. Natl. Acad. Sci. USA 96: 14523).

25 While previously identified markers such as PSA, PSM, PCTA and PSCA have facilitated efforts to diagnose and treat prostate cancer, there is need for the identification of additional markers and therapeutic targets for prostate and related cancers in order to further improve diagnosis and therapy.

### SUMMARY OF THE INVENTION

The present invention relates to a novel family of cell surface serpentine transmembrane antigens. Two of the proteins in this family are exclusively or predominantly expressed in the prostate, as well as in prostate cancer, and thus members of this family have been termed “STEAP” (Six Transmembrane Epithelial Antigen of the Prostate). Four particular human STEAPs are described and characterized herein. The human STEAPs exhibit a high degree of structural conservation among them but show no significant structural homology to any known human proteins.

The prototype member of the STEAP family, STEAP-1, appears to be a type IIIa membrane protein expressed predominantly in prostate cells in normal human tissues. Structurally, STEAP-1 is a 339 amino acid protein characterized by a molecular topology of six transmembrane domains and intracellular N- and C- termini, suggesting that it folds in a “serpentine” manner into three extracellular and two intracellular loops. STEAP-1 protein expression is maintained at high levels across various stages of prostate cancer. Moreover, STEAP-1 is highly over-expressed in certain other human cancers. In particular, cell surface expression of STEAP-1 has been definitively confirmed in a variety of prostate and prostate cancer cells, lung cancer, bladder cancer cells and colon cancer cells. These characteristics indicate that STEAP-1 is a specific cell-surface tumor antigen expressed at high levels in prostate, bladder, colon, and other cancers.

A second member of the family, STEAP-2, is a 454 amino acid protein with a predicted molecular topology similar to that of STEAP-1. STEAP-2, like STEAP-1, is prostate-specific in normal human tissues and is also expressed in prostate cancer. Alignment of the STEAP-2 and STEAP-1 ORFs shows 54.9% identity over a 237 amino acid residue overlap, and the locations of the six putative transmembrane domains in STEAP-2 coincide with the locations of the transmembrane domains in STEAP-1 (FIG. 11A-B).

STEAP-3 and STEAP-4 are also described herein. These are also structurally related, and show unique expression profiles. In particular, STEAP-3 and STEAP-4 appear to show a different tissue restriction patterns. An amino acid sequence alignment of all four STEAPs is shown in FIG. 11A-B.

The invention provides polynucleotides corresponding or complementary to all or part of the STEAP gene as described herein, mRNAs, and/or coding sequences, preferably in isolated form, including polynucleotides encoding STEAP proteins and fragments thereof, DNA, RNA, DNA/RNA hybrid, and related molecules, 5 polynucleotides or oligonucleotides complementary to the STEAP gene or mRNA sequences or parts thereof, and polynucleotides or oligonucleotides which hybridize to the STEAP gene, mRNAs, or to STEAP-encoding polynucleotides. Also provided are means for isolating cDNAs and the genes encoding STEAP. Recombinant DNA molecules containing STEAP polynucleotides, cells transformed or transduced with such 10 molecules, and host vector systems for the expression of STEAP gene products are also provided.

The invention further provides STEAP proteins and polypeptide fragments thereof. The invention further provides antibodies that bind to STEAP proteins and polypeptide fragments thereof, including polyclonal and monoclonal antibodies, murine 15 and other mammalian antibodies, chimeric antibodies, humanized and fully human antibodies, and antibodies labeled with a detectable marker, and antibodies conjugated to radionuclides/radioisotopes, toxins or other therapeutic compositions. The invention further provides methods for detecting the presence of STEAP polynucleotides and proteins in various biological samples, as well as methods for identifying cells that express a 20 STEAP. The invention further provides various therapeutic compositions and strategies for treating prostate and other cancers, including particularly, antibody, vaccine and small molecule therapy..

The invention further provides methods for detecting the presence of STEAP polynucleotides and proteins in various biological samples, as well as methods for 25 identifying cells that express STEAP. The invention further provides various therapeutic compositions and strategies, including particularly, antibody, vaccine and small molecule therapy, for treating cancers of the prostate. The extracellular nature of this protein presents a number of therapeutic approaches using molecules that target STEAP and its function, as well as molecules that target other proteins, factors and ligands that interact 30 with STEAP. These therapeutic approaches include antibody therapy with anti-STEAP antibodies, small molecule therapies, and vaccine therapies. In addition, given its up-regulated expression in prostate cancer, STEAP is useful as a diagnostic, staging and/or

prognostic marker for prostate cancer and, similarly, may be a marker for other cancers expressing this protein. In particular, STEAP-1 provides an excellent marker for identifying prostate cancer metastases.

## 5 **BRIEF DESCRIPTION OF THE FIGURES**

FIG. 1. STEAP-1 structure. 1A-B: Nucleotide and deduced amino acid sequences of STEAP-1 (8P1B4) clone 10 cDNA (SEQ ID NOS. 1 and 2, respectively). The start methionine is indicated in bold at amino acid residue position 1 and six putative transmembrane domains are indicated in bold and are underlined. 1C: Schematic representation of STEAP-1 transmembrane orientation; amino acid residues bordering the predicted extracellular domains are indicated and correspond to the numbering scheme of FIG. 1A-B. 1D: G/C rich 5' non-coding sequence of the STEAP-1 gene (SEQ ID NO: 3) as determined by overlapping sequences of clone 10 and clone 3.

FIG. 2. Predominant expression of STEAP-1 in prostate tissue. First strand cDNA was prepared from 16 normal tissues, the LAPC xenografts (4AD, 4AI and 9AD) and HeLa cells. Normalization was performed by PCR using primers to actin and GAPDH. Semi-quantitative PCR, using primers derived from STEAP-1 (8P1D4) cDNA (FIG. 1A-B), shows predominant expression of STEAP-1 in normal prostate and the LAPC xenografts. The following primers were used to amplify STEAP-1:

8P1D4.1 5' ACTTTGTTGATGACCAGGATTGGA 3' (SEQ ID NO: 4)

8P1D4.2 5' CAGAACTTCAGCACACACAGGAAC 3' (SEQ ID NO: 5).

FIG. 2A. Lanes are as follows: 1 brain; 2 prostate; 3 LAPC-4 AD; 4 LAPC-4 AI; 5 LAPC-9 AD; 6 HeLa; 7 murine cDNA; 8 negative control.

FIG. 2B. Lanes are as follows: 1 brain; 2 heart; 3 kidney; 4 liver; 5 lung; 6 pancreas; 7 placenta; 8 skeletal muscle.

FIG. 2C. Lanes are as follows: 1 colon; 2 ovary; 3 leukocytes; 4 prostate; 5 small intestine; 6 spleen; 7 testis; 8 thymus.



FIG. 3. Northern blot analyses of STEAP-1 expression in various normal human tissues and prostate cancer xenografts, showing predominant expression of STEAP-1 in prostate tissue.

5

FIG. 3A: Two multiple tissue northern blots (Clontech) were probed with a full length STEAP cDNA clone 10 (FIG. 1A-B). Size standards in kilobases (kb) are indicated on the side. Each lane contains 2 µg of mRNA that was normalized by using a β-actin probe. A1 brain; A2 amygdala; A3 caudate nucleus; A4 cerebellum; A5 cerebral cortex; A6 frontal lobe; A7 hippocampus; A8 medulla oblongata; B1 occipital lobe; B2 putamen; B3 substantia nigra; B4 temporal lobe; B5 thalamus; B6 sub-thalamic nucleus; B7 spinal cord; C1 heart; C2 aorta; C3 skeletal muscle; C4 colon; C5 bladder; C6 uterus; **C7 prostate**; C8 stomach; D1 testis; D2 ovary; D3 pancreas; D4 pituitary gland; D5 adrenal gland; D6 thyroid gland; D7 salivary gland; D8 mammary gland; E1 kidney; **E2 liver**; E3 small intestine; E4 spleen; E5 thymus ; E6 peripheral leukocytes; E7 lymph node; E8 bone marrow; F1 appendix; **F2 lung**; **F3 trachea**; F4 placenta; G1 fetal brain; G2 fetal heart; G3 fetal kidney; **G4 fetal liver**; G5 fetal spleen; G6 fetal thymus; G7 fetal lung.

FIG. 3B: Multiple tissue RNA dot blot (Clontech, Human Master Blot cat# 7770-1) probed with STEAP-1 cDNA clone 10 (FIG. 1A-B), showing approximately five-fold greater expression in prostate relative to other tissues with significant detectable expression.

FIG. 4A-B. Nucleotide sequence (SEQ ID NO: 6) of STEAP-1 GTH9 clone corresponding to the 4 kb message on northern blots (FIG. 3A). The sequence contains an intron of 2399 base pairs relative to the STEAP-1 clone 10 sequence of FIG. 1A-B; coding regions are nucleotides 96-857 and 3257- 3510 (indicated in bold). The start ATG is in bold and underlined, the STOP codon is in bold and underlined, and the intron-exon boundaries are underlined.

FIG. 5. Expression of STEAP-1 in prostate and multiple cancer cell lines and prostate cancer xenografts. Xenograft and cell line filters were prepared with 10 µg of total RNA per lane. The blots were analyzed using the STEAP-1 clone 10 as probe. All RNA samples were normalized by ethidium bromide staining and subsequent analysis with a β-actin probe. FIG. 5A: Expression in various cancer cell lines and xenografts and prostate. Lanes as follows: (1) PrEC cells, (2) normal prostate tissue, (3) LAPC-4 AD xenograft, (4) LAPC-4 AI xenograft, (5) LAPC-9 AD xenograft, (6) LAPC-9 AI xenograft, (7) LNCaP cells, (8) PC-3 cells, (9) DU145 cells, (10) PANC-1 cells, (11) BxPC-3 cells, (12) HPAC cells, (13) Capan-1 cells, (14) CACO-2 cells, (15) LOVO cells, (16) T84 cells, (17) COLO-205 cells, (18) KCL-22 cells (acute lymphocytic leukemia, ALL), (19) HT1197 cells, (20) SCABER cells, (21) UM-UC-3 cells, (22) TCCSUP cells, (23) J82 cells, (24) 5637 cells, (25) RD-ES cells (Ewing sarcoma, EWS), (26) CAMA-1 cells, (27) DU4475 cells, (28) MCF-7 cells, (29) MDA-MB-435s cells, (30) NTERA-2 cells, (31) NCCIT cells, (32) TERA-1 cells, (33) TERA-2 cells, (34) A431 cells, (35) HeLa cells, (36) OV-1063 cells, (37) PA-1 cells, (38) SW 626 cells, (39) CAO-3 cells. FIG. 5B: The expression of STEAP-1 in subcutaneously (sc) grown LAPC xenografts compared to the expression in LAPC-4 and LAPC-9 xenografts grown in the tibia (it) of mice.

FIG. 6A-B. Western blot analysis of STEAP-1 protein expression in tissues and multiple cell lines. Western blots of cell lysates prepared from prostate cancer xenografts and cell lines were probed with a polyclonal anti-STEAP-1 antibody preparation. The samples contain 20 µg of protein and were normalized with anti-Grb-2 probing of the western blots.

FIG. 7A-B. Cell surface biotinylation of STEAP-1. FIG. 7A: Cell surface biotinylation of 293T cells transfected with vector alone or with vector containing cDNA encoding 6His-tagged STEAP-1. Cell lysates were immunoprecipitated with specific antibodies, transferred to a membrane and probed with horseradish peroxidase-conjugated streptavidin. Lanes 1-4 and 6 correspond to immunoprecipitates from lysates prepared from STEAP-1 expressing 293T cells. Lanes 5 and 7 are immunoprecipitates

from vector transfected cells. The immunoprecipitations were performed using the following antibodies: (1) sheep non-immune, (2) anti-Large T antigen, (3) anti- CD71 (transferrin receptor), (4) anti-His, (5) anti-His, (6) anti-STEAP-1, (7) anti-STEAP-1. FIG. 7B: Prostate cancer (LNCaP, PC-3, DU145), bladder cancer (UM-UC-3, TCCSUP) and colon cancer (LOVO, COLO) cell lines were either biotinylated (+) or not (-) prior to lysis. Western blots of streptavidin-gel purified proteins were probed with anti-STEAP-1 antibodies. Molecular weight markers are indicated in kilodaltons (kD).

FIG. 8. Immunohistochemical analysis of STEAP-1 expression using anti-STEAP-1 polyclonal antibody. Tissues were fixed in 10% formalin and embedded in paraffin. Tissue sections were stained using anti-STEAP-1 polyclonal antibodies directed towards the N-terminal peptide. FIG. 8A: LNCaP cells probed in the presence of N-terminal STEAP-1 peptide 1, FIG. 8B: LNCaP plus non specific peptide 2, FIG. 8C: normal prostate tissue, FIG. 8D: grade 3 prostate carcinoma, FIG. 8E: grade 2, Gleason 7 prostate carcinoma, FIG. 8F: LAPC-9 AD xenograft, FIG. 8G: normal bladder, FIG. 8H: normal colon. All images are at 400x magnification.

FIG. 9A-D. Nucleotide (SEQ ID NO: 7) and deduced amino acid (SEQ ID NO: 8) sequences of STEAP-2 (98P4B6) clone GTD3 cDNA. The start methionine and Kozak sequence are indicated in bold, and the putative transmembrane domains are underlined in bold. The 5' UTR exhibits a high GC content of 72%.

FIG. 10A-E. Nucleotide (SEQ ID NO: 9) and deduced amino acid (SEQ ID NO: 10) sequences of STEAP-3. Kozak region is in bold.

25

FIG. 10F. Nucleotide sequences (SEQ ID NOS: 11-14, respectively) of dbEST database entries corresponding to additional STEAP family members obtained by searching with the protein sequence of STEAP-1.

FIG. 11. Primary structural comparisons of STEAP family proteins:

30

FIG. 11A-B. Amino acid sequence alignment of STEAPs 1-4 (SEQ ID NOS: 2, 8, 10 and 15, respectively) using PIMA program (PIMA 1.4 program at Internet address: <<http://dot.imgen.bcm.tmc.edu:9331/multi-align/multi-align.html>>); transmembrane domains identified by the SOSUI program (available at Internet address [www.tuat.ac.jp/~mitaku/adv\\_sosui/submit.html](http://www.tuat.ac.jp/~mitaku/adv_sosui/submit.html)) are in bold. PIMA maximal linkage clustering results shown; identical residues shown in bold.

FIG. 11C. Amino acid sequence alignment of STEAP-1 (8P1D4 clone 10; SEQ ID NO: 2) and STEAP-2 (98P4B6 clone GTD3; SEQ ID NO: 8) sequences. The alignment was performed using the SIM alignment program of the Baylor College of Medicine Search Launcher Web site. Transmembrane domains are indicated in boldface. The results show a 54.9% identity in a 237 residues overlap (Score: 717.0; Gap frequency: 0.0%).

FIG. 11D. Amino acid sequence alignment of STEAP-1 (SEQ ID NO: 2) and STEAP-3 (98P4B6 clone GTD3; SEQ ID NO: 10) sequences. Identical residues indicated with asterisks. SIM results: 40.9% identity in 264 residues overlap; Score: 625.0; Gap frequency: 0.0%.

FIG. 11E. Amino acid sequence alignment of STEAP-2 (SEQ ID NO: 8) and STEAP-3 (98P4B6 clone GTD3; SEQ ID NO: 10) sequences. Identical residues indicated with asterisks. SIM results: 47.8% identity in 416 residues overlap; Score: 1075.0; Gap frequency: 0.2%.

FIG. 12. Expression of STEAP-3 mRNA in normal tissues by northern blot (FIG. 12A-B) and RT-PCR (FIG. 12B). For RT-PCR analysis, first strand cDNA was prepared from 16 normal tissues. Normalization was performed by PCR using primers to actin and GAPDH. Semi-quantitative PCR, using primers to AI139607, shows predominant expression of AI139607 in placenta and prostate after 25 cycles of amplification. The following primers were used to amplify AI139607:

AI139607.1 5' TTAGGACAACCTTGATCACCAGCA 3' (SEQ ID NO: 16)  
AI139607.2 5' TGTCCAGTCCAACTGGGTTATTT 3' (SEQ ID NO: 17).

FIG. 12A. Lanes are as follows (from left to right): Panel 1: heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas; Panel 2: spleen, thymus, prostate, testis, ovary, small intestine, colon and white blood cells.

5

FIG. 12B. Lanes are as follows: 1 brain; 2 heart; 3 kidney; 4 liver; 5 lung; 6 pancreas; 7 placenta; 8 skeletal muscle.

FIG. 12C. Lanes are as follows: 1 colon; 2 ovary; 3 leukocytes; 4 prostate; 5 small intestine; 6 spleen; 7 testis; 8 thymus.

FIG. 13. Predominant expression of STEAP-4/R80991 in liver. First strand cDNA was prepared from 16 normal tissues. Normalization was performed by PCR using primers to actin and GAPDH. Semi-quantitative PCR, using primers to R80991, shows predominant expression of R80991 in liver after 25 cycles of amplification. The following primers were used to amplify R80991:

R80991.1      5' AGGGAGTTCAGCTTCGTTTCAGTC 3' (SEQ ID NO: 18)

R80991.2      5' GGTAGAACTTGTCGCGGCTCTCCT 3' (SEQ ID NO: 19).

FIG. 13A. Lanes are as follows: 1 brain; 2 heart; 3 kidney; 4 liver; 5 lung; 6 pancreas; 7 placenta; 8 skeletal muscle.

FIG. 13B. Lanes are as follows: 1 colon; 2 ovary; 3 leukocytes; 4 prostate; 5 small intestine; 6 spleen; 7 testis; 8 thymus.

25

FIG. 14. Predominant expression of STEAP-2 (98P4B6) in prostate tissue. First strand cDNA was prepared from 8 normal tissues, the LAPC xenografts (4AD, 4AI and 9AD) and HeLa cells. Normalization was performed by PCR using primers to actin and GAPDH. Semi-quantitative PCR, using primers to 98P4B6, shows predominant expression of 98P4B6 in normal prostate and the LAPC xenografts. The following primers were used to amplify STEAP II:

30

98P4B6.1      5' GACTGAGCTGGAAGTGAATTTGT 3' (SEQ ID NO: 20)

98P4B6.2      5' TTTGAGGAGACTTCATCTCACTGG 3' (SEQ ID NO: 21).

FIG. 14A. Lanes are as follows: 1 brain; 2 prostate; 3 LAPC-4 AD; 4 LAPC-4 AI; 5 LAPC-9 AD; 6 HeLa; 7 murine cDNA; 8 negative control.

5

FIG. 14B. Lanes are as follows: 1 colon; 2 ovary; 3 leukocytes; 4 prostate; 5 small intestine; 6 spleen; 7 testis; 8 thymus.

FIG. 15. Expression of the prostate-specific STEAP-2/98P4B6 gene in normal tissues and in prostate cancer xenografts determined by Northern blot analysis. Human normal tissue filters (A and B) were obtained from CLONTECH and contain 2 µg of mRNA per lane. Xenograft filter (C) was prepared with 10 µg of total RNA per lane. The blots were analyzed using the SSH derived 98P4B6 clone as probe. All RNA samples were normalized by ethidium bromide staining.

15

FIG. 15A. Lanes are as follows: 1 heart; 2 brain; 3 placenta; 4 lung; 5 liver; 6 skeletal muscle; 7 kidney; 8 pancreas.

FIG. 15B. Lanes are as follows: 1 spleen; 2 thymus; 3 prostate; 4 testis; 5 ovary; 6 small intestine; 7 colon; 8 leukocytes.

20

FIG. 15C. Lanes are as follows: 1 prostate; 2 LAPC-4 AD; 3 LAPC-4 AI; 4 LAPC-9 AD; LAPC-9AI.

FIG. 16. Expression of STEAP-2 in prostate and select cancer cell lines as determined by Northern blot analysis. Xenograft and cell line filters were prepared with 10 µg total RNA per lane. The blots were analyzed using an SSH derived 98P4B6 clone as probe. All RNA samples were normalized by ethidium bromide staining. Lanes are as follows: 1 prostate; 2 LAPC-4 AD; 3 LAPC-4 AI; 4 LAPC-9 AD; 5 LAPC-9 AI; 6 TsuPr1; 7 DU145; 8 LNCaP; 9 PC-3; 10 LAPC-4 CL; 11 PrEC; 12 HT1197; 13 SCaBER; 14 UM-UC-3; 15 TCCSUP; 16 J82; 17 5637; 18 RD-ES; 19 293T; 20 PANC-1;

30

21 BxPC-3; 22 HPAC; 23 Capan-1; 24 LS180; 25 SK-CO-1; 26 CaCo-2; 27 LoVo; 28 T84; 29 Colo-205; 30 BT-20; 31 CAMA-1; 32 DU4475; 33 MCF-7; 34 MDA-MB-435s; 35 NTERA-2; 36 NCCIT; 37 TERA-1; 38 TERA-2; 39 A431; 40 HeLa; 41 OV-1063; 42 PA-1; 43 SW626; 44 CAOV-3.

5

FIG. 17. Chromosomal localization of STEAP family members. The chromosomal localizations of the STEAP genes described herein were determined using the GeneBridge4 radiation hybrid panel (Research Genetics, Huntsville AL). The mapping for STEAP-2 and AI139607 was performed using the Stanford G3 radiation hybrid panel (Research Genetics, Huntsville AL).

10

FIG. 18. Schematic representation of Intron-Exon boundaries within the ORF of human STEAP-1 gene. A total of 3 introns (i) and 4 exons (e) were identified.

15

FIG. 19. Zooblot southern analysis of STEAP-1 gene in various species. Genomic DNA was prepared from several different organisms including human, monkey, dog, mouse, chicken and Drosophila. Ten micrograms of each DNA sample was digested with EcoRI, blotted onto nitrocellulose and probed with a STEAP-1 probe. Size standards are indicated on the side in kilobases (kb).

20

FIG. 20. Southern blot analysis of mouse BAC with a STEAP-1 probe. DNA was prepared from human cells to isolate genomic DNA and from a mouse BAC clone (12P11) that contains the mouse STEAP gene. Each DNA sample was digested with EcoRI, blotted onto nitrocellulose and probed. Eight micrograms of genomic DNA was compared to 250 ng of mouse BAC DNA. Lanes are as follows: (1) 1 kb ladder; (2) human female genomic; (3) 12P11 BAC mus; (4) human female genomic; (5) 12P11 BAC mus; (6) 3T3.

25

FIG. 21A. Immunohistochemical staining using a sheep polyclonal antibody directed against STEAP-1 and showing pericellular staining in a bladder cancer specimen.

30

FIG. 21B. Immunohistochemical staining using a sheep polyclonal antibody directed against STEAP-1 and showing pericellular staining in a second bladder cancer specimen.

5           FIG. 21C. Immunohistochemical staining using a sheep polyclonal antibody directed against STEAP-1 and showing pericellular staining in a lung cancer specimen.

FIG. 21D. Immunohistochemical staining using a sheep polyclonal antibody directed against STEAP-1 and showing pericellular staining in a second lung cancer  
10       specimen.

FIG. 22A. STEAP-2 expression in normal prostate shown by RNA in situ hybridization with an antisense probe.

15           FIG. 22B. STEAP-2 expression in normal prostate by RNA in situ hybridization using a sense probe as control.

FIG. 23A. STEAP-2 expression in prostate cancer shown by RNA in situ hybridization with an antisense probe.

20

FIG. 23B. STEAP-2 expression in prostate cancer control for RNA in situ hybridization using a sense probe.

FIG. 24. Expression of STEAP-2 in various cancer tissues as examined using  
25       RT-PCR. Lane 1 represents a sample from an LAPC4 AD xenograft; lane 2 is LAPC9  
AD xenograft; lane 3 is LAPC9 AD<sup>2</sup> xenograft (grown with human bone explant); lane 4  
is LAPC9 AD IT (grown intratibially); lane 5 is pooled tissue from colon cancer patients;  
lane 6 is pooled tissue from lung cancer patients; M represents a marker lane; lane 7 is  
patient normal prostate tissue; lane 8 is patient prostate cancer tissue; lane 9 is pooled  
30       tissue from kidney cancer patients; lane 10 is pooled tissue from bladder cancer patients;  
lane 11 is HeLa cells; and lane 12 is a water blank.



FIG. 25. RNA dot blot analysis of STEAP-2 expression in 76 normal tissues, showing prostate-specific expression. RNA tissue sources: A1 whole brain; A2 cerebellum, left; A3 substantia nigra; A4 heart; A5 esophagus; A6 colon, transverse; A7 kidney; A8 lung; A9 liver; A10 HL60, leukemia; A11 fetal brain; B1 cerebral cortex; B2 cerebellum, right; B3 accumbens nucleus; B4 aorta; B5 stomach; B6 colon, descending; B7 skeletal muscle; B8 placenta; B9 pancreas; B10 HeLa, S3; B11 fetal heart; C1 frontal lobe; C2 corpus callosum; C3 thalamus; C4 atrium, left; C5 duodenum; C6 rectum; C7 spleen; C8 bladder; C9 adrenal gland; C10 K562, leukemia; C11 fetal kidney; D1 parietal lobe; D2 amygdala; D3 pituitary gland; D4 atrium, right; D5 jejunum; D6 blank; D7 thymus; D8 uterus; D9 thyroid gland; D10 MOLT-4, leukemia; D11 fetal liver; E1 occipital lobe; E2 caudate nucleus; E3 spinal cord; E4 ventricle, left; E5 ileum; E6 blank; E7 leukocytes; **E8 prostate**; E9 salivary gland; E10 RAJI, lymphoma; E11 fetal spleen; F1 temporal lobe; F2 hippocampus; F3 blank; F4 ventricle, right; F5 ileocecum; F6 blank; F7 lymph node; F8 testis; F9 mammary gland; F10 DAUDI, lymphoma; F11 fetal thymus; G1 paracentral gyrus; G2 medulla oblongata; G3 blank; G4 interventricular septum; G5 appendix; G6 blank; G7 bone marrow; G8 ovary; G9 blank; G10 SW480, colon cancer; G11 fetal lung; H1 pons; H2 putamen; H3 blank; H4 apex of the heart; H5 colon, ascending; H6 blank; H7 trachea; H8 blank; H9 blank; H10 A549, lung cancer; H11 blank.

20

FIG. 26. Western blot with anti-phosphotyrosine (4G10 mAb) showing that expression of STEAP-2 in PC3 cells is sufficient to induce phosphorylation of various proteins on tyrosine residues, including p150, p120 and p75. An overlay using anti-Grb2 Ab shows that the gel was equally loaded.

25

FIG. 27A. Western blot using anti-phospho-p38 showing that expression of STEAP-1 or STEAP-2 in PC3 cells activates the p38 kinase, as compared to PC3-neo controls. TNF-NaSaI and NaSaI, known p38 activators, served as positive controls.

FIG. 27B. Western blot, as in FIG. 27A, using anti-p38 to demonstrate equal protein loading on the gels.

FIG. 28A. Western blot using anti-phospho-ERK to examine activation of the ERK pathway in 1% FBS. PC3 cells expressing Ras (positive control), STEAP-1, STEAP-2, or neo (negative control), were grown in 1% FBS. The results show that, while expression of the neo control gene has no effect on ERK phosphorylation, expression of STEAP-1 and STEAP-2 induces ERK phosphorylation. An anti-ERK antibody was used to confirm the presence of ERK in all lanes.

FIG. 28B. Western blot using anti-phospho-ERK to examine activation of the ERK pathway in 0.1% and 10% FBS. PC3 cells expressing STEAP-1 or neo (negative control), were grown in either 0.1% or 10% FBS. The results confirm that expression of STEAP-1 is sufficient to induce activation of the ERK signaling cascade.

FIG. 29. Signaling in PC3-STEAP-1 cells mediated by odorants.

FIG. 29A. Anti-phosphotyrosine western blot of PC3 cells, stably expressing neo, were grown overnight in 0.1% FBS to allow for receptor occupancy, and then treated with citralva, ethylvanillin or IBMP. Treatment with 10% FBS was used as a control.

FIG. 29B. Anti-phosphotyrosine western blot of PC3 cells, stably expressing STEAP-1, were treated as described for FIG. 29A. The results show that citralva and ethylvanillin specifically induce the phosphorylation of p136-140 in PC3-STEAP-1 cells. In addition, citralva induces the *de novo* phosphorylation of a protein at 160-200kDa.

FIG. 30. Activation of the ERK cascade by odorants. Anti-ERK western blot of PC3 cells, stably expressing either neo or STEAP-1, were grown overnight in 0.1% FBS. Cells were then treated with citralva for 5 min. Treatment with 10% FBS was used as a control. Whole cell lysates were analyzed using anti-phospho-ERK. The results show that citralva induces the phosphorylation of ERK, and therefore activation of the ERK pathway, in a STEAP-1 specific manner.

FIG. 31A. Anti-STEAP-1 immunohistochemistry analysis on the LAPC-9 orthotopic prostate cancer tumor (magnification 200X). Cells expressing STEAP-1 show perinuclear staining.

5           FIG. 31B. Anti-STEAP-1 immunohistochemistry analysis on the LAPC-9 lymph node metastasis (magnification 400X). Cells expressing STEAP-1 show perinuclear staining.

10           FIG. 31C-D. Anti-STEAP-1 immunohistochemistry analysis on the LAPC-9 lung metastasis (magnification 800X). Cells expressing STEAP-1 show perinuclear staining.

15           FIG. 31E-F. Anti- PSA immunohistochemistry analysis on a LAPC-9 prostate cancer lung micrometastasis (magnification 800X). Cells expressing PSA show perinuclear staining.

FIG. 32A. Immunohistochemical detection of STEAP-1 showing intense pericellular staining in lymph node metastasis of a human patient.

20           FIG. 32B. Immunohistochemical detection of STEAP-1 showing intense pericellular staining in bone metastasis of a human patient.

FIG. 33. Western blot showing that anti-STEAP-1 murine pAb recognizes STEAP-1 protein in engineered cell lines and endogenous STEAP-1 protein in LNCaP cells. Lysates of LNCaP cells and 293T cells transfected with either pcDNA 3.1 MYC/HIS tagged STEAP-1 or neo empty vector, and RAT1 cells engineered to express STEAP-1 or a neo control gene, were separated by SDS-PAGE and transferred to nitrocellulose. The blot was then subjected to anti-STEAP western analysis using a 1:1000 dilution of serum from mice immunized with a GST-STEAP-1 fusion protein.

30

## **DETAILED DESCRIPTION OF THE INVENTION**

The present invention relates to a novel family of cell surface serpentine transmembrane antigens. Two of the proteins in this family are exclusively or predominantly expressed in the prostate, as well as in prostate cancer, and thus members of this family have been termed “STEAP” (Six Transmembrane Epithelial Antigen of the Prostate). Four particular human STEAPs are described and characterized herein. The human STEAPs exhibit a high degree of structural conservation among them but show no significant structural homology to any known human proteins. The present invention relates to methods and compositions for the diagnosis and therapy of prostate and other cancers, which methods utilize isolated polynucleotides corresponding to human STEAP genes, proteins encoded by the STEAP genes and fragments thereof, and antibodies capable of specifically recognizing and binding to STEAP proteins.

Unless otherwise defined, all terms of art, notations and other scientific terminology used herein are intended to have the meanings commonly understood by those of skill in the art to which this invention pertains. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not necessarily be construed to represent a substantial difference over what is generally understood in the art. The techniques and procedures described or referenced herein are generally well understood and commonly employed using conventional methodology by those skilled in the art, such as, for example, the widely utilized molecular cloning methodologies described in Sambrook et al., Molecular Cloning: A Laboratory Manual 2nd. edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. As appropriate, procedures involving the use of commercially available kits and reagents are generally carried out in accordance with manufacturer defined protocols and/or parameters unless otherwise noted.

As used herein, the terms “advanced prostate cancer”, “locally advanced prostate cancer”, “advanced disease” and “locally advanced disease” mean prostate cancers that have extended through the prostate capsule, and are meant to include stage C disease under the American Urological Association (AUA) system, stage C1 - C2 disease under

the Whitmore-Jewett system, and stage T3 - T4 and N+ disease under the TNM (tumor, node, metastasis) system. In general, surgery is not recommended for patients with locally advanced disease, and these patients have substantially less favorable outcomes compared to patients having clinically localized (organ-confined) prostate cancer.

- 5 Locally advanced disease is clinically identified by palpable evidence of induration beyond the lateral border of the prostate, or asymmetry or induration above the prostate base. Locally advanced prostate cancer is presently diagnosed pathologically following radical prostatectomy if the tumor invades or penetrates the prostatic capsule, extends into the surgical margin, or invades the seminal vesicles.

- 10 As used herein, the terms “metastatic prostate cancer” and “metastatic disease” mean prostate cancers that have spread to regional lymph nodes or to distant sites, and are meant to include stage D disease under the AUA system and stage TxNxM+ under the TNM system. As is the case with locally advanced prostate cancer, surgery is generally not indicated for patients with metastatic disease, and hormonal (androgen  
15 ablation) therapy is the preferred treatment modality. Patients with metastatic prostate cancer eventually develop an androgen-refractory state within 12 to 18 months of treatment initiation, and approximately half of these patients die within 6 months thereafter. The most common site for prostate cancer metastasis is bone. Prostate cancer bone metastases are, on balance, characteristically osteoblastic rather than osteolytic (i.e.,  
20 resulting in net bone formation). Bone metastases are found most frequently in the spine, followed by the femur, pelvis, rib cage, skull and humerus. Other common sites for metastasis include lymph nodes, lung, liver and brain. Metastatic prostate cancer is typically diagnosed by open or laparoscopic pelvic lymphadenectomy, whole body radionuclide scans, skeletal radiography, and/or bone lesion biopsy.

- 25 As used herein, the term “polynucleotide” means a polymeric form of nucleotides of at least 10 bases or base pairs in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide, and is meant to include single and double stranded forms of DNA.

- 30 As used herein, the term “polypeptide” means a polymer of at least 10 amino acids. Throughout the specification, standard three letter or single letter designations for amino acids are used.

As used herein, the terms “hybridize”, “hybridizing”, “hybridizes” and the like, used in the context of polynucleotides, are meant to refer to conventional hybridization conditions, preferably such as hybridization in 50% formamide/6XSSC/0.1% SDS/100 µg/ml ssDNA, in which temperatures for hybridization are above 37°C and  
5 temperatures for washing in 0.1XSSC/0.1% SDS are above 55°C, and most preferably to stringent hybridization conditions.

“Stringency” of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher  
10 temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable  
15 sequence, the higher the relative temperature that can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., Current Protocols in Molecular Biology, Wiley Interscience Publishers, (1995).

“Stringent conditions” or “high stringency conditions”, as defined herein, may be  
20 identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium  
25 chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency  
30 wash consisting of 0.1 x SSC containing EDTA at 55°C.

“Moderately stringent conditions” may be identified as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

In the context of amino acid sequence comparisons, the term “identity” is used to express the percentage of amino acid residues at the same relative positions that are the same. Also in this context, the term “homology” is used to express the percentage of amino acid residues at the same relative positions that are either identical or are similar, using the conserved amino acid criteria of BLAST analysis, as is generally understood in the art. For example, % identity values may be generated by WU-BLAST-2 (Altschul et al., Methods in Enzymology, 266: 460-480 (1996): <http://blast.wustl.edu/blast/README.html>). Further details regarding amino acid substitutions, which are considered conservative under such criteria, are provided below.

Additional definitions are provided throughout the subsections that follow.

### **STEAP POLYNUCLEOTIDES**

One aspect of the invention provides polynucleotides corresponding or complementary to all or part of a STEAP gene, mRNA, and/or coding sequence, preferably in isolated form, including polynucleotides encoding a STEAP protein and fragments thereof, DNA, RNA, DNA/RNA hybrid, and related molecules, polynucleotides or oligonucleotides complementary to a STEAP gene or mRNA sequence or a part thereof, and polynucleotides or oligonucleotides that hybridize to a STEAP gene, mRNA, or to a STEAP encoding polynucleotide (collectively, “STEAP polynucleotides”). As used herein, STEAP genes and proteins are meant to include the STEAP-1, STEAP-2 and STEAP-3 genes and proteins, and the gene and protein

corresponding to GenBank Accession number R80991 (STEAP-4), and the genes and proteins corresponding to other STEAP proteins and structurally similar variants of the foregoing. Such other STEAP proteins and variants will generally have coding sequences that are highly homologous to the STEAP coding sequence, and preferably will share at least about 50% amino acid identity and at least about 60% amino acid homology (using BLAST criteria), more preferably sharing 70% or greater homology (using BLAST criteria).

The STEAP family member gene sequences described herein encode STEAP proteins sharing unique highly conserved amino acid sequence domains which distinguish them from other proteins. Proteins which include one or more of these unique highly conserved domains may be related to the STEAP family members or may represent new STEAP proteins. Referring to FIG. 11A-B, which is an amino acid sequence alignment of the full STEAP-1, STEAP-2, and STEAP-3 protein sequences as well as the partial STEAP-4 sequence, it is clear that the STEAPs are closely related at the structural level. Referring to FIG. 11C, which is an amino acid sequence alignment of the full STEAP-1 and STEAP-2 protein sequences, close structural conservation is apparent, particularly in the predicted transmembrane domains. The STEAP-1 and STEAP-2 sequences share 54.9% identity over a 237 amino acid overlap. Additional amino acid sequence alignments between the STEAPs are shown in FIGS. 11D and 11E. These alignments show that STEAP-1 and STEAP-3 are 40.9% identical over a 264 amino acid region, while STEAP-2 and STEAP-3 are 47.8% identical over a 416 amino acid region.

A STEAP polynucleotide may comprise a polynucleotide having the nucleotide sequence of human STEAP-1 as shown in FIG. 1A-B, the nucleotide sequence of human STEAP-2 as shown in FIG. 9A-D, the nucleotide sequence of human STEAP-3 as shown in FIG. 10A-E, or the nucleotide sequence of STEAP-4 as shown in FIG. 10F, or a sequence complementary thereto, or a polynucleotide fragment of any of the foregoing. Another embodiment comprises a polynucleotide which encodes the human STEAP-1, STEAP-2, STEAP-3 or STEAP-4 protein amino acid sequences, a sequence complementary thereto, or a polynucleotide fragment of any of the foregoing. Another embodiment comprises a polynucleotide which is capable of hybridizing under stringent hybridization conditions to the human STEAP-1 cDNA shown in FIG. 1A-B, the



human STEAP-2 cDNA shown in FIG. 9A-D, the human STEAP-3 cDNA shown in FIG. 10A-E, or the STEAP-4 as shown in FIG. 10F, or to a polynucleotide fragment thereof.

Typical embodiments of the invention disclosed herein include STEAP

5 polynucleotides encoding specific portions of a STEAP mRNA sequence such as those that encode the protein and fragments thereof. For example, representative embodiments of the invention disclosed herein include: polynucleotides encoding about amino acid 1 to about amino acid 10 of a STEAP protein shown in Figure 11A-B, polynucleotides encoding about amino acid 20 to about amino acid 30 of a STEAP

10 protein shown in Figure 11A-B, polynucleotides encoding about amino acid 30 to about amino acid 40 of a STEAP protein shown in Figure 11A-B, polynucleotides encoding about amino acid 40 to about amino acid 50 of a STEAP protein shown in Figure 11A-B, polynucleotides encoding about amino acid 50 to about amino acid 60 of a STEAP protein shown in Figure 11A-B, polynucleotides encoding about amino acid 60 to about

15 amino acid 70 of a STEAP protein shown in Figure 11A-B, polynucleotides encoding about amino acid 70 to about amino acid 80 of a STEAP protein shown in Figure 11A-B, polynucleotides encoding about amino acid 80 to about amino acid 90 of a STEAP protein shown in Figure 11A-B and polynucleotides encoding about amino acid 90 to about amino acid 100 of a STEAP protein shown in Figure 11A-B, etc. Following this

20 scheme, polynucleotides (of at least 10 amino acids) encoding further portions of the amino acid sequence of a STEAP protein are typical embodiments of the invention. Such portions of a STEAP protein include amino acids 100-339 of a STEAP-1 protein shown in Figure 11A-B, or amino acids 100-454 of a STEAP-2 protein shown in Figure 11A-B, or amino acids 100-459 of a STEAP-3 protein shown in Figure 11A-B, and

25 amino acids 100-133 of a STEAP-4 protein shown in Figure 11A-B. Polynucleotides encoding larger portions of the STEAP protein are also contemplated. For example polynucleotides encoding from about amino acid 1 (or 20 or 30 or 40 etc.) to about amino acid 20, (or 30, or 40 or 50 etc.) of a STEAP protein shown in Figure 11A-B may be generated by a variety of techniques well known in the art.

30 Additional illustrative embodiments of the invention disclosed herein include STEAP polynucleotide fragments encoding one or more of the biological motifs contained within the STEAP protein sequence. In one embodiment, typical

polynucleotide fragments of the invention can encode one or more of the regions of STEAP that exhibit homology to other STEAP family members, such as one or more of the transmembrane domains. In another embodiment of the invention, typical polynucleotide fragments can encode sequences that are unique to one or more STEAP  
5 alternative splicing variants. In yet another embodiment, typical polynucleotide fragments can encode an immunogenic portion of a STEAP protein. One example of an immunogenic portion of a STEAP protein is amino acid residues 14 through 28 of the STEAP-1 amino acid sequence as shown in FIG. 1A-B (WKMKPRRNLEEDDYL; SEQ ID NO: 22).

10 The polynucleotides of the preceding paragraphs have a number of different specific uses. As STEAPs are differentially expressed in prostate and other cancers, these polynucleotides may be used in methods assessing the status of STEAP gene products in normal versus cancerous tissues. Typically, polynucleotides encoding specific regions of a STEAP protein may be used to assess the presence of perturbations  
15 (such as deletions, insertions, point mutations etc.) in specific regions of the STEAP gene products. Exemplary assays include both RT-PCR assays as well as single-strand conformation polymorphism (SSCP) analysis (see e.g. Marrogi et al., J. Cutan. Pathol. 26(8): 369-378 (1999), both of which utilize polynucleotides encoding specific regions of a protein to examine these regions within the protein. Assays and methods for analyzing  
20 sequences to detect single nucleotide polymorphisms are also available (Irizarry, et al., 2000, Nature Genetics 26(2):223-236.

Other specifically contemplated embodiments of the invention disclosed herein are genomic DNA, cDNAs, ribozymes, and antisense molecules, including morpholino anti-sense molecules, as well as nucleic acid molecules based on an alternative backbone or  
25 including alternative bases, whether derived from natural sources or synthesized. For example, antisense molecules can be RNAs or other molecules, including peptide nucleic acids (PNAs) or non-nucleic acid molecules such as phosphorothioate derivatives, that specifically bind DNA or RNA in a base pair-dependent manner. A skilled artisan can readily obtain these classes of nucleic acid molecules using the STEAP polynucleotides and  
30 polynucleotide sequences disclosed herein.

Antisense technology entails the administration of exogenous oligonucleotides that bind to a target polynucleotide located within the cells. The term "antisense" refers

to the fact that such oligonucleotides are complementary to their intracellular targets, e.g., STEAP. See for example, Jack Cohen, OLIGODEOXYNUCLEOTIDES, Antisense Inhibitors of Gene Expression, CRC Press, 1989; and Synthesis 1:1-5 (1988). The STEAP antisense oligonucleotides of the present invention include derivatives such as S-oligonucleotides (phosphorothioate derivatives or S-oligos, see, Jack Cohen, supra), which exhibit enhanced cancer cell growth inhibitory action. S-oligos (nucleoside phosphorothioates) are isoelectronic analogs of an oligonucleotide (O-oligo) in which a nonbridging oxygen atom of the phosphate group is replaced by a sulfur atom. The S-oligos of the present invention may be prepared by treatment of the corresponding O-oligos with 3H-1,2-benzodithiol-3-one-1,1-dioxide, which is a sulfur transfer reagent. See Iyer, R. P. et al, J. Org. Chem. 55:4693-4698 (1990); and Iyer, R. P. et al., J. Am. Chem. Soc. 112:1253-1254 (1990), the disclosures of which are fully incorporated by reference herein. Additional STEAP antisense oligonucleotides of the present invention include morpholino antisense oligonucleotides known in the art (see e.g. Partridge et al., 1996, Antisense & Nucleic Acid Drug Development 6: 169-175).

The STEAP antisense oligonucleotides of the present invention typically may be RNA or DNA that is complementary to and stably hybridizes with the first 100 N-terminal codons or last 100 C-terminal codons, or overlapping with the ATG start site, of the STEAP genome or the corresponding mRNA. While absolute complementarity is not required, high degrees of complementarity are preferred. Use of an oligonucleotide complementary to this region allows for the selective hybridization to STEAP mRNA and not to mRNA specifying other regulatory subunits of protein kinase. Preferably, the STEAP antisense oligonucleotides of the present invention are a 15 to 30-mer fragment of the antisense DNA molecule having a sequence that hybridizes to STEAP mRNA. Optionally, STEAP antisense oligonucleotide is a 30-mer oligonucleotide that is complementary to a region in the first 10 N-terminal codons and last 10 C-terminal codons of STEAP. Alternatively, the antisense molecules are modified to employ ribozymes in the inhibition of STEAP expression. L. A. Couture & D. T. Stinchcomb; *Trends Genet* 12: 510-515 (1996).

Further specific embodiments of this aspect of the invention include primers and primer pairs, which allow the specific amplification of the polynucleotides of the invention or of any specific parts thereof, and probes that selectively or specifically

hybridize to nucleic acid molecules of the invention or to any part thereof. Probes may be labeled with a detectable marker, such as, for example, a radioisotope, fluorescent compound, bioluminescent compound, a chemiluminescent compound, metal chelator or enzyme. Such probes and primers can be used to detect the presence of a STEAP polynucleotide in a sample and as a means for detecting a cell expressing a STEAP protein. Examples of such probes include polypeptides comprising all or part of a human STEAP cDNA sequence shown in Fig. 1A-B (SEQ ID NO: 1), FIG. 9A-D (SEQ ID NO: 7) or Fig. 10A-E (SEQ ID NO: 9). Examples of primer pairs capable of specifically amplifying STEAP mRNAs are also described in the Examples that follow. As will be understood by the skilled artisan, a great many different primers and probes may be prepared based on the sequences provided in herein and used effectively to amplify and/or detect a STEAP mRNA.

As used herein, a polynucleotide is said to be "isolated" when it is substantially separated from contaminant polynucleotides that correspond or are complementary to genes other than the STEAP gene or that encode polypeptides other than STEAP gene product or fragments thereof. A skilled artisan can readily employ nucleic acid isolation procedures to obtain an isolated STEAP polynucleotide.

The STEAP polynucleotides of the invention are useful for a variety of purposes, including but not limited to their use as probes and primers for the amplification and/or detection of the STEAP gene(s), mRNA(s), or fragments thereof; as reagents for the diagnosis and/or prognosis of prostate cancer and other cancers; as tools for identifying molecules that inhibit calcium entry specifically into prostate cells; as coding sequences capable of directing the expression of STEAP polypeptides; as tools for modulating or inhibiting the expression of the STEAP gene(s) and/or translation of the STEAP transcript(s); and as therapeutic agents.

### **MOLECULAR AND BIOCHEMICAL FEATURES OF THE STEAPs**

The invention relates to a novel family of proteins, termed STEAPs. Four STEAPs are specifically described herein by way of structural, molecular and biochemical features. As is further described in the Examples which follow, the STEAPs have been characterized in a variety of ways. For example, analyses of nucleotide coding and amino acid sequences were conducted in order to identify conserved structural elements

within the STEAP family. Extensive RT-PCR and Northern blot analyses of STEAP mRNA expression were conducted in order to establish the range of normal and cancerous tissues expressing the various STEAP messages. Western blot, immunohistochemical and flow cytometric analyses of STEAP protein expression were conducted to determine protein expression profiles, cell surface localization and gross molecular topology of STEAP.

The prototype member of the STEAP family, STEAP-1, is a six-transmembrane cell surface protein of 339 amino acids with no identifiable homology to any known human protein. The cDNA nucleotide and deduced amino acid sequences of human STEAP-1 are shown in FIG. 1A-B. A gross topological schematic of the STEAP-1 protein integrated within the cell membrane is shown in FIG. 1B. STEAP-1 expression is predominantly prostate-specific in normal tissues. Specifically, extensive analysis of STEAP-1 mRNA and protein expression in normal human tissues shows that STEAP-1 protein is predominantly expressed in prostate and, to a far smaller degree, in bladder. STEAP-1 mRNA is also relatively prostate specific, with only very low level expression detected in a few other normal tissues. In cancer, STEAP-1 mRNA and protein is consistently expressed at high levels in prostate cancer (including androgen-dependent and androgen-independent tumors) and during all stages of the disease. STEAP-1 is also expressed in other cancers. Specifically, STEAP-1 mRNA is expressed at very high levels in bladder, colon, pancreatic, and ovarian cancer (as well as other cancers). In addition, cell surface expression of STEAP-1 protein has been established in prostate, bladder, lung and colon cancers. Therefore, STEAP-1 has all of the hallmark characteristics of an excellent diagnostic and therapeutic target for the treatment of certain cancers, including particularly prostate, colon and bladder carcinomas.

A second member of the family, STEAP-2, is a 454 amino acid protein encoded by a distinct gene and having a predicted molecular topology similar to that of STEAP-1. The cDNA nucleotide and deduced amino acid sequences of STEAP-2 are shown in FIG. 9A-D. Amino acid alignment of the STEAP-1 and STEAP-2 sequences show a high degree of structural conservation (54.9% identity over a 237 amino acid residue overlap, and the locations of the six putative transmembrane domains in STEAP-1 and STEAP-2 coincide (FIGS. 11A-B, 11C). Structural homology between these STEAP-1 and STEAP-2 is highest in the regions spanned by the first putative extracellular loop to

the fifth transmembrane domain. However, some significant structural differences between STEAP-1 and STEAP-2 are apparent. For example, STEAP-2 exhibits a 205 a.a. long intracellular N-terminus (compared to 69 a.a. in STEAP-1) and a short 4 a.a. intracellular C-terminus (compared to 26 a.a. in STEAP-1). In addition, both the  
5 STEAP-1 and STEAP-2 genes are located on chromosome 7, but on different arms. These differences could imply significant differences in function and/or interaction with intracellular signaling pathways.

STEAP-2 is expressed only in normal prostate among human tissues tested (FIGS. 14 and 15) and is also expressed in prostate cancer (FIG. 15), and thus shows  
10 some similarity in expression profile to STEAP-1. However, STEAP-2 exhibits a different mRNA expression profile relative to STEAP-1 in prostate cancer samples (compare FIGS. 3 and 15) and in other non-prostate cancers tested (compare FIGS. 5 and 16). These differences in the expression profiles of STEAP-1 and STEAP-2 suggest that they are differentially regulated.

15 STEAP-3 and STEAP-4 appear to be closely related to both STEAP-1 and STEAP-2 on a structural level, and both appear to be transmembrane proteins as well. STEAP-3 is more related to STEAP-2 (47.8% identity) than to STEAP-1 (40.9% identity). STEAP-3 and STEAP-4 show unique expression profiles. STEAP-3, for example, appears to have an expression pattern which is predominantly restricted to  
20 placenta and, to a smaller degree, expression is seen in prostate but not in other normal tissues tested. STEAP-4 seems to be expressed predominantly in liver by RT-PCR analysis. Neither STEAP-3 nor STEAP-4 appear to be expressed in prostate cancer xenografts which exhibit high level STEAP-1 and STEAP-2 expression.

The STEAP proteins exhibit characteristics of proteins involved in cell signaling  
25 pathways. Specifically, STEAP-1 and STEAP-2, when expressed in PC3 cells, activate phosphorylation of p38, a protein involved in the MAPK signaling cascade. In addition, STEAP-2 expression induces tyrosine phosphorylation, and STEAP-1 mediates activation of tyrosine kinase in odorant-treated cells. These findings support the use of STEAP-related molecules and cells modified to express STEAP in high throughput  
30 assays to identify molecules capable of altering cellular signaling pathways, leading to the identification of novel therapeutic agents. In one embodiment, the assay identifies molecules capable of inhibiting STEAP function, which molecules are thereby capable of

modulating the progression of cancer or other disease associated with dysregulated cell growth.

Three of the four STEAPs described herein map to human chromosome 7 (STEAP-1, -2 and 3). Interestingly, STEAP-1 maps within 7p22 (7p22.3), a large region of allelic gain reported for both primary and recurrent prostate cancers (Visakorpi et al., 1995 *Cancer Res.* 55: 342, Nupponen et al., 1998 *American J. Pathol.* 153: 141), suggesting that up-regulation of STEAP-1 in cancer might include genomic mechanisms. In addition, both STEAP-2 and STEAP-3 locate to chromosome 7q21, suggesting that these two genes arose by gene duplication.

Other cell surface molecules that contain six transmembrane domains include ion channels (Dolly and Parcej, 1996 *J Bioenerg Biomembr* 28:231) and water channels or aquaporins (Reizer et al., 1993 *Crit Rev Biochem Mol Biol* 28:235). Structural studies show that both types of molecules assemble into tetrameric complexes to form functional channels (Christie, 1995, *Clin Exp Pharmacol Physiol* 22:944, Walz et al., 1997 *Nature* 387:624, Cheng et al., 1997 *Nature* 387:627). Immunohistochemical staining of STEAP-1 in the prostate gland seems to be concentrated at the cell-cell boundaries, with less staining detected at the luminal side. This may suggest a role for STEAP-1 in tight-junctions, gap-junctions, cell communication, adhesion or as a transporter protein.

To test these possibilities, xenopus oocytes (or other cells) expressing STEAP may be analyzed using voltage-clamp and patch-clamp experiments to determine if STEAP functions as an ion-channel. Oocyte cell volume may also be measured to determine if STEAP exhibits water channel properties. If STEAPs function as channel or gap-junction proteins, they may serve as excellent targets for inhibition using, for example, antibodies, small molecules, and polynucleotides capable of inhibiting expression or function. The restricted expression pattern in normal tissue, and the high levels of expression in cancer tissue suggest that interfering with STEAP function may selectively kill cancer cells.

Since the STEAP gene family is predominantly expressed in epithelial tissue, it seems possible that the STEAP proteins function as ion channels, transport proteins or gap-junction proteins in epithelial cell function. Ion channels have been implicated in proliferation and invasiveness of prostate cancer cells (Lalani et al., 1997, *Cancer Metastasis Rev* 16:29). Both rat and human prostate cancer cells contain sub-population

of cells with higher and lower expression levels of sodium channels. Higher levels of sodium channel expression correlate with more aggressive invasiveness in vitro (Smith et al., 1998, FEBS Lett. 423:19). Similarly, it has been shown that a specific blockade of sodium channels inhibits the invasiveness of PC-3 cells in vitro (Laniado et al., 1997, Am. J. Pathol. 150:1213), while specific inhibition of potassium channels in LNCaP cells inhibited cell proliferation (Skryma et al., 1997, Prostate 33:112). These reports suggest a role for ion channels in prostate cancer and also demonstrate that small molecules that inhibit ion channel function may interfere with prostate cancer proliferation.

#### 10 **ISOLATION OF STEAP-ENCODING NUCLEIC ACID MOLECULES**

The STEAP cDNA sequences described herein enable the isolation of other polynucleotides encoding STEAP gene product(s), as well as the isolation of polynucleotides encoding STEAP gene product homologues, alternatively spliced isoforms, allelic variants, and mutant forms of the STEAP gene product. Various molecular cloning methods that can be employed to isolate full length cDNAs encoding a STEAP gene are well known (See, for example, Sambrook, J. et al. Molecular Cloning: A Laboratory Manual, 2d edition., Cold Spring Harbor Press, New York, 1989; Current Protocols in Molecular Biology. Ausubel et al., Eds., Wiley and Sons, 1995). For example, lambda phage cloning methodologies may be conveniently employed, using commercially available cloning systems (e.g., Lambda ZAP Express, Stratagene). Phage clones containing STEAP gene cDNAs may be identified by probing with labeled STEAP cDNA or a fragment thereof. For example, in one embodiment, a STEAP cDNA or a portion thereof can be synthesized and used as a probe to retrieve overlapping and full length cDNAs corresponding to a STEAP gene. The STEAP gene itself may be isolated by screening genomic DNA libraries, bacterial artificial chromosome libraries (BACs), yeast artificial chromosome libraries (YACs), and the like, with STEAP DNA probes or primers.

#### **RECOMBINANT DNA MOLECULES AND HOST-VECTOR SYSTEMS**

30 The invention also provides recombinant DNA or RNA molecules containing a STEAP polynucleotide, including but not limited to phages, plasmids, phagemids, cosmids, YACs, BACs, as well as various viral and non-viral vectors well known in the



art, and cells transformed or transfected with such recombinant DNA or RNA molecules. As used herein, a recombinant DNA or RNA molecule is a DNA or RNA molecule that has been subjected to molecular manipulation in vitro. Methods for generating such molecules are well known (see, for example, Sambrook et al, 1989, 5 supra).

The invention further provides a host-vector system comprising a recombinant DNA molecule containing a STEAP polynucleotide within a suitable prokaryotic or eukaryotic host cell. Examples of suitable eukaryotic host cells include a yeast cell, a plant cell, or an animal cell, such as a mammalian cell or an insect cell (e.g., a baculovirus- 10 infectible cell such as an Sf9 cell). Examples of suitable mammalian cells include various prostate cancer cell lines such as LNCaP, PC-3, DU145, LAPC-4, TsuPr1, other transfectable or transducible prostate cancer cell lines, as well as a number of mammalian cells routinely used for the expression of recombinant proteins (e.g., COS, CHO, 293, 293T cells). More particularly, a polynucleotide comprising the coding sequence of a 15 STEAP may be used to generate STEAP proteins or fragments thereof using any number of host vector systems routinely used and widely known in the art.

A wide range of host vector systems suitable for the expression of STEAP proteins or fragments thereof are available, see for example, Sambrook et al., 1989, supra; Current Protocols in Molecular Biology, 1995, supra). Preferred vectors for mammalian 20 expression include but are not limited to pcDNA 3.1 myc-His-tag (Invitrogen) and the retroviral vector pSR $\alpha$ tkneo (Muller et al., 1991, MCB 11:1785). Using these expression vectors, STEAP may be preferably expressed in several prostate cancer and non-prostate cell lines, including for example 293, 293T, rat-1, 3T3, PC-3, LNCaP and TsuPr1. The host vector systems of the invention are useful for the production of a STEAP protein or 25 fragment thereof. Such host-vector systems may be employed to study the functional properties of STEAP and STEAP mutations.

Proteins encoded by the STEAP genes, or by fragments thereof, will have a variety of uses, including but not limited to generating antibodies and in methods for identifying ligands and other agents and cellular constituents that bind to a STEAP gene 30 product. Antibodies raised against a STEAP protein or fragment thereof may be useful in diagnostic and prognostic assays, imaging methodologies (including, particularly, cancer

imaging), and therapeutic methods in the management of human cancers characterized by expression of a STEAP protein, including but not limited to cancer of the prostate. Various immunological assays useful for the detection of STEAP proteins are contemplated, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), immunocytochemical methods, and the like. Such antibodies may be labeled and used as immunological imaging reagents capable of detecting prostate cells (e.g., in radioscintigraphic imaging methods). STEAP proteins may also be particularly useful in generating cancer vaccines, as further described below.

### **STEAP PROTEINS**

Another aspect of the present invention provides various STEAP proteins and polypeptide fragments thereof. As used herein, a STEAP protein refers to a protein that has or includes the amino acid sequence of human STEAP-1 as provided in FIG. 1A-B, human STEAP-2 as provided in FIG. 9A-D, human STEAP-3 as provided in FIG. 10A-E, the amino acid sequence of other mammalian STEAP homologs (e.g., STEAP-4) and variants, as well as allelic variants and conservative substitution mutants of these proteins that have STEAP biological activity, to the extent that such variants and homologs can be isolated/generated and characterized without undue experimentation following the methods outlined below. Fusion proteins that combine parts of different STEAP proteins or fragments thereof, as well as fusion proteins of a STEAP protein and a heterologous polypeptide, are also included. Such STEAP proteins will be collectively referred to as the STEAP proteins, the proteins of the invention, or STEAP. As used herein, the term "STEAP polypeptide" refers to a polypeptide fragment or a STEAP protein of at least 10 amino acids, preferably at least 15 amino acids.

A specific embodiment of a STEAP protein comprises a polypeptide having the amino acid sequence of human STEAP-1 as shown in FIG. 1A-B. Another embodiment of a STEAP protein comprises a polypeptide containing the STEAP-2 amino acid sequence as shown in FIG. 9A-D. Another embodiment comprises a polypeptide containing the STEAP-3 amino acid sequence of shown in FIG. 10A-E. Yet another embodiment comprises a polypeptide containing the partial STEAP-4 amino acid sequence of shown in FIG. 11A-B.

In general, naturally occurring allelic variants of human STEAP will share a high degree of structural identity and homology (e.g., 90% or more identity). Typically, allelic variants of the STEAP proteins will contain conservative amino acid substitutions within the STEAP sequences described herein or will contain a substitution of an amino acid  
5 from a corresponding position in a STEAP homologue. One class of STEAP allelic variants will be proteins that share a high degree of homology with at least a small region of a particular STEAP amino acid sequence, but will further contain a radical departure from the sequence, such as a non-conservative substitution, truncation insertion or frame shift.

10 Conservative amino acid substitutions can frequently be made in a protein without altering either the conformation or the function of the protein. Such changes include substituting any of isoleucine (I), valine (V), and leucine (L) for any other of these hydrophobic amino acids; aspartic acid (D) for glutamic acid (E) and vice versa; glutamine (Q) for asparagine (N) and vice versa; and serine (S) for threonine (T) and vice  
15 versa. Other substitutions can also be considered conservative, depending on the environment of the particular amino acid and its role in the three-dimensional structure of the protein. For example, glycine (G) and alanine (A) can frequently be interchangeable, as can alanine (A) and valine (V). Methionine (M), which is relatively hydrophobic, can frequently be interchanged with leucine and isoleucine, and sometimes  
20 with valine. Lysine (K) and arginine (R) are frequently interchangeable in locations in which the significant feature of the amino acid residue is its charge and the differing pK's of these two amino acid residues are not significant. Still other changes can be considered "conservative" in particular environments.

STEAP proteins, including variants, comprise at least one epitope in common  
25 with a STEAP protein having an amino acid sequence shown in Fig. 11A-B, such that an antibody that specifically binds to a STEAP protein or variant will also specifically bind to the STEAP protein having an amino acid sequence shown in Fig. 11A-B. One class of STEAP protein variants shares 90% or more identity with an amino acid sequence of Fig. 11A-B. A more specific class of STEAP protein variants comprises an extracellular  
30 protein SCP motif as described above. Preferred STEAP protein variants are capable of exhibiting one or more of the defensin functions described herein, including, for

example, the ability to induce tumor death or to chemoattract and/or induce migration of cells.

STEAP proteins may be embodied in many forms, preferably in isolated form. As used herein, a protein is said to be “isolated” when physical, mechanical or chemical methods are employed to remove the STEAP protein from cellular constituents that are normally associated with the protein. A skilled artisan can readily employ standard purification methods to obtain an isolated STEAP protein. A purified STEAP protein molecule will be substantially free of other proteins or molecules that impair the binding of STEAP to antibody or other ligand. The nature and degree of isolation and purification will depend on the intended use. Embodiments of a STEAP protein include a purified STEAP protein and a functional, soluble STEAP protein. In one form, such functional, soluble STEAP proteins or fragments thereof retain the ability to bind antibody or other ligand.

The invention also provides STEAP polypeptides comprising biologically active fragments of the STEAP amino acid sequence, such as a polypeptide corresponding to part of the amino acid sequences for STEAP-1 as shown in FIG. 1A-B, STEAP-2 as shown in FIG. 9A-D, STEAP-3 as shown in FIG. 10A-E, or STEAP-4 as shown in FIG. 11A-B. Such polypeptides of the invention exhibit properties of a STEAP protein, such as the ability to elicit the generation of antibodies which specifically bind an epitope associated with a STEAP protein. Polypeptides comprising amino acid sequences which are unique to a particular STEAP protein (relative to other STEAP proteins) may be used to generate antibodies which will specifically react with that particular STEAP protein. For example, referring to the amino acid alignment of the STEAP structures shown in FIGS. 11A-E, the skilled artisan will readily appreciate that each molecule contains stretches of sequence unique to its structure. These unique stretches can be used to generate antibodies specific to a particular STEAP. Similarly, regions of conserved sequence may be used to generate antibodies that may bind to multiple STEAPs.

Embodiments of the invention disclosed herein include a wide variety of art accepted variants of STEAP proteins such as polypeptides having amino acid insertions, deletions and substitutions. STEAP variants can be made using methods known in the art such as site-directed mutagenesis, alanine scanning, and PCR mutagenesis. Site-

directed mutagenesis [Carter et al., *Nucl. Acids Res.*, 13:4331 (1986); Zoller et al., *Nucl. Acids Res.*, 10:6487 (1987)], cassette mutagenesis [Wells et al., *Gene*, 34:315 (1985)], restriction selection mutagenesis [Wells et al., *Philos. Trans. R. Soc. London Ser.A*, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the STEAP variant DNA. Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, *The Proteins*, (W.H. Freeman & Co., N.Y.); Chothia, J. *Mol. Biol.*, 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isosteric amino acid can be used.

As discussed above, embodiments of the claimed invention include polypeptides containing less than the full amino acid sequence of a STEAP protein shown in Figure 11A-B. For example, representative embodiments of the invention disclosed herein include polypeptides consisting of about amino acid 1 to about amino acid 10 of a STEAP protein shown in Figure 11A-B, polypeptides consisting of about amino acid 20 to about amino acid 30 of a STEAP protein shown in Figure 11A-B, polypeptides consisting of about amino acid 30 to about amino acid 40 of a STEAP protein shown in Figure 11A-B, polypeptides consisting of about amino acid 40 to about amino acid 50 of a STEAP protein shown in Figure 11A-B, polypeptides consisting of about amino acid 50 to about amino acid 60 of a STEAP protein shown in Figure 11A-B, polypeptides consisting of about amino acid 60 to about amino acid 70 of a STEAP protein shown in Figure 11A-B, polypeptides consisting of about amino acid 70 to about amino acid 80 of a STEAP protein shown in Figure 11A-B, polypeptides consisting of about amino acid 80 to about amino acid 90 of a STEAP protein shown in Figure 11A-B and polypeptides consisting of about amino acid 90 to about amino acid 100 of a STEAP protein shown in Figure 11A-B, etc. Following this scheme, polypeptides consisting of portions of the amino acid sequence of amino acids 100-339 of a STEAP-1 protein shown in Figure 11A-B, or amino acids 100-454 of a STEAP-2 protein shown in Figure 11A-B, or amino

acids 100-459 of a STEAP-3 protein shown in Figure 11A-B, or amino acids 100-133 of a STEAP-4 protein shown in Figure 11A-B, are typical embodiments of the invention. Polypeptides consisting of larger portions of the STEAP protein are also contemplated. For example polypeptides consisting of about amino acid 1 (or 20 or 30 or 40 etc.) to  
5 about amino acid 20, (or 30, or 40 or 50 etc.) of a STEAP protein shown in Figure 11A-B may be generated by a variety of techniques well known in the art.

Additional illustrative embodiments of the invention disclosed herein include STEAP polypeptides containing the amino acid residues of one or more of the biological motifs contained within a STEAP polypeptide sequence as shown in Fig. 11A-B.

10 STEAP polypeptides containing one or more of these motifs or other select regions of interest described herein will typically include an additional 5 to 25 or more amino acid residues of adjacent STEAP protein sequence on one or both sides of the selected motif(s). In one embodiment, typical polypeptides of the invention can contain one or more of the regions of STEAP that exhibit homology to one or more other STEAP  
15 proteins. In another embodiment, typical polypeptides of the invention can contain one or more immunogenic portions of a STEAP protein. One example of an immunogenic portion of a STEAP protein is amino acid residues 14 through 28 of the STEAP-1 amino acid sequence as shown in FIG. 1A-B (WKMKPRRNLEEDDYL; SEQ ID NO: 22). In another embodiment, typical polypeptides of the invention can contain one or more  
20 predicted HLA-A2 binding peptides such as amino acids 165-173 of STEAP-1, amino acids 86-94 of STEAP-1, amino acids 262-270 of STEAP-1, amino acids 302-310 of STEAP-1, amino acids 158-166 of STEAP-1, amino acids 227-235 of STEAP-2, amino acids 402-410 of STEAP-2, amino acids 307-315 of STEAP-2, amino acids 306-314 of STEAP-2, and amino acids 100-108 of STEAP-2. In another embodiment, typical  
25 polypeptides of the invention consist of all or part of a fragment of STEAP-2, such as amino acids 1-245, 2-204, 121-454, 153-165, 182-454, 183-387, 276-453, 345-358, or 419-454 of the STEAP-2 protein shown in FIG. 8. Related embodiments of these inventions include polypeptides containing combinations of the different motifs discussed above with preferable embodiments being those that contain no insertions, deletions or  
30 substitutions either within the motifs or the intervening sequences of these polypeptides.

STEAP polypeptides can be generated using standard peptide synthesis technology or using chemical cleavage methods well known in the art based on the

amino acid sequences of the human STEAP proteins disclosed herein. Alternatively, recombinant methods can be used to generate nucleic acid molecules that encode a polypeptide fragment of a STEAP protein. In this regard, the STEAP-encoding nucleic acid molecules described herein provide means for generating defined fragments of STEAP proteins. STEAP polypeptides are particularly useful in generating and characterizing domain specific antibodies (e.g., antibodies recognizing an extracellular or intracellular epitope of a STEAP protein), in identifying agents or cellular factors that bind to STEAP or a particular structural domain thereof, and in various therapeutic contexts, including but not limited to cancer vaccines. STEAP polypeptides containing particularly interesting structures can be predicted and/or identified using various analytical techniques well known in the art, including, for example, the methods of Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis, or on the basis of immunogenicity. Fragments containing such structures are particularly useful in generating subunit specific anti-STEAP antibodies or in identifying cellular factors that bind to STEAP.

In a specific embodiment described in the examples that follow, a secreted form of STEAP may be conveniently expressed in 293T cells transfected with a CMV-driven expression vector encoding STEAP with a C-terminal 6XHis and MYC tag (pcDNA3.1/mycHis, Invitrogen). The secreted HIS-tagged STEAP in the culture media may be purified using a nickel column and standard techniques. Alternatively, an AP-tag system may be used. Various constructs for expression of STEAP are described in the examples below.

Modifications of STEAP such as covalent modifications are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of an STEAP polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C- terminal residues of the STEAP. Another type of covalent modification of the STEAP polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence STEAP (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more

glycosylation sites that are not present in the native sequence STEAP. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Another type of covalent modification of STEAP comprises linking the STEAP  
5 polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The STEAP of the present invention may also be modified in a way to form a chimeric molecule comprising STEAP fused to another, heterologous polypeptide or  
10 amino acid sequence. In one embodiment, such a chimeric molecule comprises a fusion of the STEAP with a polyhistidine epitope tag, which provides an epitope to which immobilized nickel can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the STEAP. In an alternative embodiment, the chimeric molecule may comprise a fusion of the STEAP with an immunoglobulin or a particular  
15 region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of an STEAP polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the  
20 immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also US Patent No. 5,428,130 issued June 27, 1995.

### **STEAP ANTIBODIES**

25 Another aspect of the invention provides antibodies that bind to STEAP proteins and polypeptides. The most preferred antibodies will selectively bind to a STEAP protein and will not bind (or will bind weakly) to non-STEAP proteins and polypeptides. Anti-STEAP antibodies that are particularly contemplated include monoclonal and polyclonal antibodies as well as fragments containing the antigen-  
30 binding domain and/or one or more complementarity determining regions of these antibodies. As used herein, an antibody fragment is defined as at least a portion of the



variable region of the immunoglobulin molecule that binds to its target, i.e., the antigen binding region.

For some applications, it may be desirable to generate antibodies which specifically react with a particular STEAP protein and/or an epitope within a particular structural domain. For example, preferred antibodies useful for cancer therapy and diagnostic imaging purposes are those which react with an epitope in an extracellular region of the STEAP protein as expressed in cancer cells. Such antibodies may be generated by using the STEAP proteins described herein, or using peptides derived from predicted extracellular domains thereof, as an immunogen. In this regard, with reference to the STEAP-1 protein topological schematic shown in FIG 1B, regions in the extracellular loops between the indicated transmembrane domains may be selected as used to design appropriate immunogens for raising extracellular-specific antibodies.

STEAP antibodies of the invention may be particularly useful in prostate cancer therapeutic strategies, diagnostic and prognostic assays, and imaging methodologies. Similarly, such antibodies may be useful in the treatment, diagnosis, and/or prognosis of other cancers, to the extent STEAP is also expressed or overexpressed in other types of cancer. The invention provides various immunological assays useful for the detection and quantification of STEAP and mutant STEAP proteins and polypeptides. Such assays generally comprise one or more STEAP antibodies capable of recognizing and binding a STEAP or mutant STEAP protein, as appropriate, and may be performed within various immunological assay formats well known in the art, including but not limited to various types of radioimmunoassays, enzyme-linked immunosorbent assays (ELISA), enzyme-linked immunofluorescent assays (ELIFA), and the like. In addition, immunological imaging methods capable of detecting prostate cancer are also provided by the invention, including but limited to radioscintigraphic imaging methods using labeled STEAP antibodies. Such assays may be used clinically in the detection, monitoring, and prognosis of prostate cancer, particularly advanced prostate cancer.

STEAP antibodies may also be used in methods for purifying STEAP and mutant STEAP proteins and polypeptides and for isolating STEAP homologues and related molecules. For example, in one embodiment, the method of purifying a STEAP protein comprises incubating a STEAP antibody, which has been coupled to a solid matrix, with a lysate or other solution containing STEAP under conditions which permit

the STEAP antibody to bind to STEAP; washing the solid matrix to eliminate impurities; and eluting the STEAP from the coupled antibody. Other uses of the STEAP antibodies of the invention include generating anti-idiotypic antibodies that mimic the STEAP protein.

- 5           STEAP antibodies may also be used therapeutically by, for example, modulating or inhibiting the biological activity of a STEAP protein or targeting and destroying cancer cells expressing a STEAP protein. Antibody therapy of prostate and other cancers is more specifically described in a separate subsection below.

- 10           Various methods for the preparation of antibodies are well known in the art. For example, antibodies may be prepared by immunizing a suitable mammalian host using a STEAP protein, peptide, or fragment, in isolated or immunoconjugated form (Antibodies: A Laboratory Manual, CSH Press, Eds., Harlow, and Lane (1988); Harlow, Antibodies, Cold Spring Harbor Press, NY (1989)). Examples of protein immunogens include recombinant STEAP (expressed in a baculovirus system, mammalian system, 15 etc.), STEAP extracellular domain or extracellular loops of STEAP protein conjugated to one or more antibody constant regions, AP-tagged STEAP, etc. In addition, fusion proteins of STEAP may also be used, such as a fusion of STEAP with GST, maltose-binding protein (MBP), green fluorescent protein (GFP), HisMax-TOPO or MycHis (see Examples below).

- 20           In a particular embodiment, a GST fusion protein comprising all or most of an open reading frame amino acid sequence as shown in Fig. 11A-B may be produced and used as an immunogen to generate appropriate antibodies. Cells expressing or overexpressing STEAP may also be used for immunizations. Similarly, any cell engineered to express STEAP may be used. Such strategies may result in the production 25 of monoclonal antibodies with enhanced capacities for recognizing endogenous STEAP. Another useful immunogen comprises STEAP peptides linked to the plasma membrane of sheep red blood cells.

- The amino acid sequences of STEAP proteins as shown in Fig. 11A-B may be used to select specific regions of a STEAP protein for generating antibodies. For 30 example, hydrophobicity and hydrophilicity analyses of the STEAP amino acid sequence may be used to identify hydrophilic regions in the STEAP structure. Regions of the STEAP protein that show immunogenic structure, as well as other regions and domains,

can readily be identified using various other methods known in the art, such as Chou-Fasman, Garnier Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis. Peptides of STEAP predicted to bind HLA-A2 may be selected for the generation of antibodies or used to generate a CTL response. Such predicted HLA-A2 binding peptides include, but are not limited to, amino acids 165-173 of STEAP-1, amino acids 86-94 of STEAP-1, amino acids 262-270 of STEAP-1, amino acids 302-310 of STEAP-1, amino acids 158-166 of STEAP-1, amino acids 227-235 of STEAP-2, amino acids 402-410 of STEAP-2, amino acids 307-315 of STEAP-2, amino acids 306-314 of STEAP-2, and amino acids 100-108 of STEAP-2. As discussed in the examples below, immunogenicity has been demonstrated with STEAP, which was used to generate polyclonal and monoclonal antibodies using rabbits and mice, respectively. This B cell response (antibody production) is the result of an initial T cell response elicited by the immunogenic portions of STEAP.

Methods for preparing a protein or polypeptide for use as an immunogen and for preparing immunogenic conjugates of a protein with a carrier such as BSA, KLH, or other carrier proteins are well known in the art. In some circumstances, direct conjugation using, for example, carbodiimide reagents may be used; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, IL, may be effective. Administration of a STEAP immunogen is conducted generally by injection over a suitable period and with use of a suitable adjuvant, as is generally understood in the art. During the immunization schedule, titers of antibodies can be taken to determine adequacy of antibody formation.

STEAP monoclonal antibodies are preferred and may be produced by various means well known in the art. For example, immortalized cell lines which secrete a desired monoclonal antibody may be prepared using the standard hybridoma technology of Kohler and Milstein or modifications which immortalize producing B cells, a is generally known. The immortalized cell lines secreting the desired antibodies are screened by immunoassay in which the antigen is the STEAP protein or STEAP fragment. When the appropriate immortalized cell culture secreting the desired antibody is identified, the cells may be expanded and antibodies produced either from in vitro cultures or from ascites fluid.

The antibodies or fragments may also be produced, using current technology, by recombinant means. Regions that bind specifically to the desired regions of the STEAP protein can also be produced in the context of chimeric or CDR grafted antibodies of multiple species origin. Humanized or human STEAP antibodies may also be produced and are preferred for use in therapeutic contexts. Methods for humanizing murine and other non-human antibodies by substituting one or more of the non-human antibody CDRs for corresponding human antibody sequences are well known (see for example, Jones et al., 1986, *Nature* 321: 522-525; Riechmann et al., 1988, *Nature* 332: 323-327; Verhoeyen et al., 1988, *Science* 239:1534-1536). See also, Carter et al., 1993, *Proc. Nat'l Acad. Sci. USA* 89: 4285 and Sims et al., 1993, *J. Immunol.* 151: 2296. Methods for producing fully human monoclonal antibodies include phage display and transgenic animal technologies (for review, see Vaughan et al., 1998, *Nature Biotechnology* 16: 535-539).

Fully human STEAP monoclonal antibodies may be generated using cloning technologies employing large human Ig gene combinatorial libraries (i.e., phage display) (Griffiths and Hoogenboom, *Building an in vitro immune system: human antibodies from phage display libraries*. In: *Protein Engineering of Antibody Molecules for Prophylactic and Therapeutic Applications in Man*. Clark, M. (Ed.), Nottingham Academic, pp 45-64 (1993); Burton and Barbas, *Human Antibodies from combinatorial libraries*. *Id.*, pp 65-82). Fully human STEAP monoclonal antibodies may also be produced using transgenic mice engineered to contain human immunoglobulin gene loci as described in U.S. Patent No. 6,150,584 and in PCT Patent Application W098/24893, published December 3, 1997 (see also, Jakobovits, 1998, *Exp. Opin. Invest. Drugs* 7(4): 607-614). This method avoids the in vitro manipulation required with phage display technology and efficiently produces high affinity authentic human antibodies.

Reactivity of STEAP antibodies with a STEAP protein may be established by a number of well known means, including western blot, immunoprecipitation, ELISA, and FACS analyses using, as appropriate, STEAP proteins, peptides, STEAP expressing cells or extracts thereof.

A STEAP antibody or fragment thereof of the invention may be labeled with a detectable marker or conjugated to a second molecule, such as a cytotoxin or other therapeutic agent, and used for targeting the second molecule to a STEAP positive cell

(Vitetta, E.S. et al., 1993, Immunotoxin therapy, in DeVita, Jr., V.T. et al., eds., Cancer: Principles and Practice of Oncology, 4th ed., J.B. Lippincott Co., Philadelphia, 2624-2636). Examples of cytotoxic agents include, but are not limited to ricin, ricin A-chain, doxorubicin, maytansinoids, daunorubicin, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, Pseudomonas exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, alpha-sarcin, gelonin, mitogellin, retstrictocin, phenomycin, enomycin, curicin, crotin, calicheamicin, sapaonaria officinalis inhibitor, and glucocorticoid and other chemotherapeutic agents, as well as radioisotopes such as  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ . Suitable detectable markers include, but are not limited to, a radioisotope, a fluorescent compound, a bioluminescent compound, chemiluminescent compound, a metal chelator or an enzyme. Antibodies may also be conjugated to an anti-cancer pro-drug activating enzyme capable of converting the pro-drug to its active form. See, for example, US Patent No. 4,975,287.

Further, bi-specific antibodies specific for two or more STEAP epitopes may be generated using methods generally known in the art. Further, antibody effector functions may be modified to enhance the therapeutic effect of STEAP antibodies on cancer cells. For example, cysteine residues may be engineered into the Fc region, permitting the formation of interchain disulfide bonds and the generation of homodimers which may have enhanced capacities for internalization, ADCC and/or complement mediated cell killing (see, for example, Caron et al., 1992, J. Exp. Med. 176: 1191-1195; Shopes, 1992, J. Immunol. 148: 2918-2922). Homodimeric antibodies may also be generated by cross-linking techniques known in the art (e.g., Wolff et al., Cancer Res. 53: 2560-2565).

### **STEAP TRANSGENIC ANIMALS**

Nucleic acids that encode STEAP or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an

embryonic stage. A transgene is a DNA that is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding STEAP can be used to clone genomic DNA encoding STEAP in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain  
5 cells that express DNA encoding STEAP.

Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for STEAP transgene incorporation with tissue-specific enhancers. Transgenic animals that include a  
10 copy of a transgene encoding STEAP introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding STEAP. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with  
15 the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of STEAP can be used to construct a STEAP “knock out” animal that has a defective or altered gene encoding STEAP as a  
20 result of homologous recombination between the endogenous gene encoding STEAP and altered genomic DNA encoding STEAP introduced into an embryonic cell of the animal. For example, cDNA encoding STEAP can be used to clone genomic DNA encoding STEAP in accordance with established techniques. A portion of the genomic DNA encoding STEAP can be deleted or replaced with another gene, such as a gene  
25 encoding a selectable marker that can be used to monitor integration.

Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector (see e.g., Thomas and Capecchi, 1987, Cell 51:503 for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced  
30 DNA has homologously recombined with the endogenous DNA are selected (see e.g., Li et al., 1992, Cell 69:915). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras (see e.g., Bradley, in

*Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed., IRL, Oxford, 1987, pp. 113-152).

5 A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knockout animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the STEAP polypeptide.

10

### **MONITORING THE STATUS OF STEAP AND ITS PRODUCTS**

Assays that evaluate the status of a STEAP gene and STEAP gene products in an individual may provide information on the growth or oncogenic potential of a biological sample from this individual. For example, because STEAP mRNA is so highly expressed  
15 in prostate, and not in most normal tissue, and because its expression is associated with certain cancers, assays that evaluate the relative levels of STEAP mRNA transcripts or proteins in a biological sample may be used to diagnose a disease associated with STEAP dysregulation, such as cancer or benign prostatic hyperplasia (BPH), and may provide prognostic information useful in defining appropriate therapeutic options. Similarly, assays  
20 that evaluate the integrity STEAP nucleotide and amino acid sequences in a biological sample, may also be used in this context.

The finding that STEAP mRNA is expressed in prostate and other cancers, and not in most normal tissue, provides evidence that this gene is associated with dysregulated cell growth and therefore identifies this gene and its products as targets that the skilled  
25 artisan can use to evaluate biological samples from individuals suspected of having a disease associated with STEAP dysregulation. In another example, because the expression of STEAP is normally restricted to prostate, one can also evaluate biological samples taken from other tissues to detect STEAP expression as an indication of metastasis. For example, as shown in Figure 31A-F, antibodies directed to STEAP-1 provide a superior  
30 marker for detection of metastases, as compared to the conventional prostate cancer marker, PSA. Such a marker can be useful both in evaluation tissue biopsies and as part of

*in vivo* imaging strategies. In this context, the evaluation of the expression status of STEAP gene and its products can be used to gain information on the disease potential of a tissue sample. The terms "expression status" in this context is used to broadly refer to the variety of factors involved in the expression, function and regulation of a gene and its products  
5 such as the level of mRNA expression, the integrity of the expressed gene products (such as the nucleic and amino acid sequences) and transcriptional and translational modifications to these molecules.

The expression status of STEAP may provide information useful for predicting susceptibility to particular disease stages, progression, and/or tumor aggressiveness. The  
10 invention provides methods and assays for determining STEAP expression status and diagnosing cancers that express STEAP, such as cancers of the prostate. STEAP expression status in patient samples may be analyzed by a number of means well known in the art, including without limitation, immunohistochemical analysis, in situ hybridization, RT-PCR analysis on laser capture micro-dissected samples, western blot analysis of clinical  
15 samples and cell lines, and tissue array analysis. Typical protocols for evaluating the expression status of the STEAP gene and gene products can be found, for example in *Current Protocols In Molecular Biology*, Units 2 [Northern Blotting], 4 [Southern Blotting], 15 [Immunoblotting] and 18 [PCR Analysis], Frederick M. Ausubul et al. eds., 1995.

In one aspect, the invention provides methods for monitoring STEAP gene  
20 products by determining the status of STEAP gene products expressed by cells in a test tissue sample from an individual suspected of having a disease associated with dysregulated cell growth (such as hyperplasia or cancer) and then comparing the status so determined to the status of STEAP gene products in a corresponding normal sample, the presence of aberrant or altered status of STEAP gene products in the test sample relative  
25 to the normal sample providing an indication of the presence of dysregulated cell growth within the cells of the individual.

The invention additionally provides methods of examining a biological sample for evidence of dysregulated cellular growth. In one embodiment, the method comprises comparing the status of STEAP in the biological sample to the status of STEAP in a  
30 corresponding normal sample, wherein alterations in the status of STEAP in the biological sample are associated with dysregulated cellular growth. The status of STEAP



in the biological sample can be evaluated by, for example, examining levels of STEAP mRNA expression or levels of STEAP protein expression. In one embodiment, an alteration in the status of STEAP is identified by the presence of STEAP expressing cells in a biological sample from a tissue in which STEAP expressing cells are normally  
5 absent.

In another aspect, the invention provides assays useful in determining the presence of cancer in an individual, comprising detecting a significant increase in STEAP mRNA or protein expression in a test cell or tissue sample relative to expression levels in the corresponding normal cell or tissue. The presence of STEAP mRNA may, for  
10 example, be evaluated in tissue samples including but not limited to colon, lung, prostate, pancreas, bladder, breast, ovary, cervix, testis, head and neck, brain, stomach, bone, etc. The presence of significant STEAP expression in any of these tissues may be useful to indicate the emergence, presence and/or severity of these cancers or a metastasis of cancer originating in another tissue, since the corresponding normal tissues do not  
15 express STEAP mRNA or express it at lower levels.

In a related embodiment, STEAP expression status may be determined at the protein level rather than at the nucleic acid level. For example, such a method or assay would comprise determining the level of STEAP protein expressed by cells in a test tissue sample and comparing the level so determined to the level of STEAP expressed in a  
20 corresponding normal sample. In one embodiment, the presence of STEAP protein is evaluated, for example, using immunohistochemical methods. STEAP antibodies or binding partners capable of detecting STEAP protein expression may be used in a variety of assay formats well known in the art for this purpose. As shown in the accompanying examples, STEAP immunoreactivity is associated with prostate, bladder and lung cancer, as  
25 well as BPH and prostate cancer metastases.

In other related embodiments, one can evaluate the integrity STEAP nucleotide and amino acid sequences in a biological sample in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like. Such  
30 embodiments are useful because perturbations in the nucleotide and amino acid sequences are observed in a large number of proteins associated with a growth dysregulated phenotype (see e.g. Marrogi et al., J. Cutan. Pathol. 26(8): 369-378 (1999)). In this

context, a wide variety of assays for observing perturbations in nucleotide and amino acid sequences are well known in the art. For example, the size and structure of nucleic acid or amino acid sequences of STEAP gene products may be observed by the northern,  
Southern, western, PCR and DNA sequencing protocols discussed herein. In addition,  
5 other methods for observing perturbations in nucleotide and amino acid sequences such as single strand conformation polymorphism analysis are well known in the art (see e.g. U.S. Patent Nos. 5,382,510 and 5,952,170).

In another embodiment, one can examine the methylation status of the STEAP gene in a biological sample. Aberrant demethylation and/or hypermethylation of CpG  
10 islands in gene 5' regulatory regions frequently occurs in immortalized and transformed cells and can result in altered expression of various genes. For example, promoter hypermethylation of the pi-class glutathione S-transferase (a protein expressed in normal prostate but not expressed in >90% of prostate carcinomas) appears to permanently silence transcription of this gene and is the most frequently detected genomic alteration  
15 in prostate carcinomas (De Marzo et al., 1999, Am. J. Pathol. 155(6): 1985-1992). In addition, this alteration is present in at least 70% of cases of high-grade prostatic intraepithelial neoplasia (PIN) (Brooks et al., 1998, Cancer Epidemiol. Biomarkers Prev., 7:531-536).

In another example, expression of the LAGE-I tumor specific gene (which is not  
20 expressed in normal prostate but is expressed in 25-50% of prostate cancers) is induced by deoxy-azacytidine in lymphoblastoid cells, suggesting that tumoral expression is due to demethylation (Lethe et al., 1998, Int. J. Cancer 76(6): 903-908). In this context, a variety of assays for examining methylation status of a gene are well known in the art. For example, one can utilize in Southern hybridization approaches methylation-sensitive  
25 restriction enzymes which can not cleave sequences that contain methylated CpG sites in order to assess the overall methylation status of CpG islands.

In addition, MSP (methylation specific PCR) can rapidly profile the methylation status of all the CpG sites present in a CpG island of a given gene. This procedure involves initial modification of DNA by sodium bisulfite (which will convert all unmethylated  
30 cytosines to uracil) followed by amplification using primers specific for methylated versus unmethylated DNA. Protocols involving methylation interference can also be found for

example in *Current Protocols In Molecular Biology*, Units 12, Frederick M. Ausubel et al. eds., 1995.

In another related embodiment, the invention provides assays useful in determining the presence of cancer in an individual, comprising detecting a significant  
5 change in the STEAP alternative splice variants expressed in a test cell or tissue sample relative to expression levels in the corresponding normal cell or tissue. The monitoring of alternative splice variants of STEAP is useful because changes in the alternative splicing of proteins is suggested as one of the steps in a series of events that lead to the progression of cancers (see e.g. Carstens et al., *Oncogene* 15(25): 3059-3065 (1997)).

10 Gene amplification provides an additional method of assessing the status of STEAP. Gene amplification may be measured in a sample directly, for example, by conventional Southern blotting, northern blotting to quantitate the transcription of mRNA [Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the  
15 sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be  
20 detected.

In addition to the tissues discussed above, peripheral blood may be conveniently assayed for the presence of cancer cells, including but not limited to prostate cancers, using RT-PCR to detect STEAP expression. The presence of RT-PCR amplifiable STEAP mRNA provides an indication of the presence of the cancer. RT-PCR detection assays for  
25 tumor cells in peripheral blood are currently being evaluated for use in the diagnosis and management of a number of human solid tumors. In the prostate cancer field, these include RT-PCR assays for the detection of cells expressing PSA and PSM (Verkaik et al., 1997, *Urol. Res.* 25: 373-384; Ghossein et al., 1995, *J. Clin. Oncol.* 13: 1195-2000; Heston et al., 1995, *Clin. Chem.* 41: 1687-1688). RT-PCR assays are well known in the art.

30 A related aspect of the invention is directed to predicting susceptibility to developing cancer in an individual. In one embodiment, a method for predicting susceptibility to cancer comprises detecting STEAP mRNA or STEAP protein in a tissue

sample, its presence indicating susceptibility to cancer, wherein the degree of STEAP mRNA expression present is proportional to the degree of susceptibility. In a specific embodiment, the presence of STEAP in prostate tissue is examined, with the presence of STEAP in the sample providing an indication of prostate cancer susceptibility (or the  
5 emergence or existence of a prostate tumor). In a closely related embodiment, one can evaluate the integrity STEAP nucleotide and amino acid sequences in a biological sample in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like, with the presence of one or more perturbations in STEAP gene products in the sample providing an indication of cancer susceptibility (or the  
10 emergence or existence of a tumor).

Yet another related aspect of the invention is directed to methods for gauging tumor aggressiveness. In one embodiment, a method for gauging aggressiveness of a tumor comprises determining the level of STEAP mRNA or STEAP protein expressed by cells in a sample of the tumor, comparing the level so determined to the level of STEAP  
15 mRNA or STEAP protein expressed in a corresponding normal tissue taken from the same individual or a normal tissue reference sample, wherein the degree of STEAP mRNA or STEAP protein expression in the tumor sample relative to the normal sample indicates the degree of aggressiveness. In a specific embodiment, aggressiveness of prostate tumors is evaluated by determining the extent to which STEAP is expressed in the tumor cells, with  
20 higher expression levels indicating more aggressive tumors. In a closely related embodiment, one can evaluate the integrity STEAP nucleotide and amino acid sequences in a biological sample in order to identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like, with the presence of one or more perturbations indicating more aggressive tumors.

Yet another related aspect of the invention is directed to methods for observing the progression of a malignancy in an individual over time. In one embodiment, methods for observing the progression of a malignancy in an individual over time comprise determining the level of STEAP mRNA or STEAP protein expressed by cells in a sample of the tumor, comparing the level so determined to the level of STEAP mRNA or STEAP protein  
25 expressed in an equivalent tissue sample taken from the same individual at a different time, wherein the degree of STEAP mRNA or STEAP protein expression in the tumor sample over time provides information on the progression of the cancer. In a specific  
30

embodiment, the progression of a cancer is evaluated by determining the extent to which STEAP expression in the tumor cells alters over time, with higher expression levels indicating a progression of the cancer. In a closely related embodiment, one can evaluate the integrity STEAP nucleotide and amino acid sequences in a biological sample in order to  
5 identify perturbations in the structure of these molecules such as insertions, deletions, substitutions and the like, with the presence of one or more perturbations indicating a progression of the cancer.

The above diagnostic approaches may be combined with any one of a wide variety of prognostic and diagnostic protocols known in the art. For example, another  
10 embodiment of the invention disclosed herein is directed to methods for observing a coincidence between the expression of STEAP gene and STEAP gene products (or perturbations in STEAP gene and STEAP gene products) and a factor that is associated with malignancy as a means of diagnosing and prognosticating the status of a tissue sample. In this context, a wide variety of factors associated with malignancy may be utilized such as  
15 the expression of genes otherwise associated with malignancy (including PSA, PSCA and PSM expression) as well as gross cytological observations (see e.g. Bocking et al., 1984, Anal. Quant. Cytol. 6(2):74-88; Epstein, 1995, Hum. Pathol. 1995 Feb;26(2):223-9; Thorson et al., 1998, Mod. Pathol. 11(6):543-51; Baisden et al., 1999, Am. J. Surg. Pathol. 23(8):918-24). Methods for observing a coincidence between the expression of STEAP  
20 gene and STEAP gene products (or perturbations in STEAP gene and STEAP gene products) and an additional factor that is associated with malignancy are useful, for example, because the presence of a set or constellation of specific factors that coincide provides information crucial for diagnosing and prognosticating the status of a tissue sample.

25 In a typical embodiment, methods for observing a coincidence between the expression of STEAP gene and STEAP gene products (or perturbations in STEAP gene and STEAP gene products) and a factor that is associated with malignancy entails detecting the overexpression of STEAP mRNA or protein in a tissue sample, detecting the overexpression of PSA mRNA or protein in a tissue sample, and observing a coincidence  
30 of STEAP mRNA or protein and PSA mRNA or protein overexpression. In a specific embodiment, the expression of STEAP and PSA mRNA in prostate tissue is examined. In a preferred embodiment, the coincidence of STEAP and PSA mRNA overexpression in

the sample provides an indication of prostate cancer, prostate cancer susceptibility or the emergence or existence of a prostate tumor.

Methods for detecting and quantifying the expression of STEAP mRNA or protein are described herein and use standard nucleic acid and protein detection and quantification technologies well known in the art. Standard methods for the detection and quantification of STEAP mRNA include in situ hybridization using labeled STEAP riboprobes, northern blot and related techniques using STEAP polynucleotide probes, RT-PCR analysis using primers specific for STEAP, and other amplification type detection methods, such as, for example, branched DNA, SISBA, TMA and the like. In a specific embodiment, semi-quantitative RT-PCR may be used to detect and quantify STEAP mRNA expression as described in the Examples that follow. Any number of primers capable of amplifying STEAP may be used for this purpose, including but not limited to the various primer sets specifically described herein. Standard methods for the detection and quantification of protein may be used for this purpose. In a specific embodiment, polyclonal or monoclonal antibodies specifically reactive with the wild-type STEAP protein may be used in an immunohistochemical assay of biopsied tissue. Antibodies directed against STEAP protein can also be used to detect STEAP in a patient specimen (e.g., blood, urine, semen or other sample) using conventional techniques such as fluorescence-activated cell sorting (FACS) and/or ELISA.

#### **IDENTIFYING MOLECULES THAT INTERACT WITH STEAP**

The STEAP protein sequences disclosed herein allow the skilled artisan to identify proteins, small molecules and other agents that interact with STEAP and pathways activated by STEAP via any one of a variety of art accepted protocols. For example one can utilize one of the variety of so-called interaction trap systems (also referred to as the “two-hybrid assay”). In such systems, molecules that interact reconstitute a transcription factor and direct expression of a reporter gene, the expression of which is then assayed. Typical systems identify protein-protein interactions in vivo through reconstitution of a eukaryotic transcriptional activator and are disclosed for example in U.S. Patent Nos. 5,955,280, 5,925,523, 5,846,722 and 6,004,746.

Alternatively one can identify molecules that interact with STEAP protein sequences by screening peptide libraries. In such methods, peptides that bind to selected receptor molecules such as STEAP are identified by screening libraries that encode a random or controlled collection of amino acids. Peptides encoded by the libraries are expressed as fusion proteins of bacteriophage coat proteins, and bacteriophage particles are then screened against the receptors of interest. Peptides having a wide variety of uses, such as therapeutic or diagnostic reagents, may thus be identified without any prior information on the structure of the expected ligand or receptor molecule. Typical peptide libraries and screening methods that can be used to identify molecules that interact with STEAP protein sequences are disclosed for example in U.S. Patent Nos. 5,723,286 and 5,733,731.

Alternatively, cell lines expressing STEAP can be used to identify protein-protein interactions mediated by STEAP. This possibility can be examined using immunoprecipitation techniques as shown by others (Hamilton BJ, et al. Biochem. Biophys. Res. Commun. 1999, 261:646-51). Typically STEAP protein can be immunoprecipitated from STEAP expressing prostate cancer cell lines using anti-STEAP antibodies. Alternatively, antibodies against His-tag can be used in a cell line engineered to express STEAP (vectors mentioned above). The immunoprecipitated complex can be examined for protein association by procedures such as western blotting, <sup>35</sup>S-methionine labeling of proteins, protein microsequencing, silver staining and two dimensional gel electrophoresis.

Small molecules that interact with STEAP can be identified through related embodiments of such screening assays. For example, small molecules can be identified that interfere with STEAP function, including molecules that interfere with STEAP's ability to bind to cells and/or to modulate tumor formation, progression, migration and/or apoptosis. Typical methods are discussed for example in U.S. Patent No. 5,928,868 and include methods for forming hybrid ligands in which at least one ligand is a small molecule. In an illustrative embodiment, the hybrid ligand is introduced into cells that in turn contain a first and a second expression vector. Each expression vector includes DNA for expressing a hybrid protein that encodes a target protein linked to a coding sequence for a transcriptional module. The cells further contains a reporter gene, the expression of which is conditioned on the proximity of the first and second hybrid

proteins to each other, an event that occurs only if the hybrid ligand binds to target sites on both hybrid proteins. Those cells that express the reporter gene are selected and the unknown small molecule or the unknown hybrid protein is identified.

A typical embodiment of this invention consists of a method of screening for a molecule that interacts with a STEAP amino acid sequence shown in Fig. 11A-B, comprising the steps of contacting a population of molecules with the STEAP amino acid sequence, allowing the population of molecules and the STEAP amino acid sequence to interact under conditions that facilitate an interaction, determining the presence of a molecule that interacts with the STEAP amino acid sequence and then separating molecules that do not interact with the STEAP amino acid sequence from molecules that do interact with the STEAP amino acid sequence. In a specific embodiment, the method further includes purifying a molecule that interacts with the STEAP amino acid sequence. In a preferred embodiment, the STEAP amino acid sequence is contacted with a library of peptides. Additional assays for identifying molecules that modulate STEAP function are described in the Examples that follow.

The invention additionally provides a method of screening for a molecule that modulates the activity of STEAP or a STEAP-related pathway. The method comprises contacting a molecule with a cell that expresses a STEAP protein and determining the activity of STEAP or a STEAP-related pathway. The determining can be via use of one of the phosphorylation assays described in the examples that follow, such as western blotting with an antibody directed to a phosphorylated signaling molecule. Alterations in the phosphorylation of the signaling molecule indicate a candidate molecule that modulates the activity of STEAP or a STEAP-related pathway.

## **THERAPEUTIC METHODS AND COMPOSITIONS**

The identification of STEAP as a prostate cancer protein opens a number of therapeutic approaches to the treatment of prostate and other STEAP-associated cancers. As discussed above, STEAP is a transmembrane protein, and its interaction with other cells and molecules likely plays a role in the regulation of the prostate environment and the initiation, development and/or progression of cancer. STEAP can be targeted for therapy via approaches aimed at inhibiting activity of the STEAP protein,



inhibiting the binding or association of STEAP protein with other cells and molecules, inhibiting transcription or translation of STEAP, and/or via the use of cancer vaccines based on STEAP. The therapeutic strategy can thus be designed to inhibit a function of the molecule or to target the STEAP molecule itself.

5           The expression profile of STEAP is reminiscent of the MAGEs, PSA and PMSA, which are tissue-specific genes that are up-regulated in melanomas and other cancers (Van den Eynde and Boon, Int J Clin Lab Res. 27:81-86, 1997). Due to their tissue-specific expression and high expression levels in cancer, these molecules are currently being investigated as targets for cancer vaccines (Durrant, Anticancer Drugs 8:727-733, 10   1997; Reynolds et al., Int J Cancer 72:972-976, 1997). The expression pattern of STEAP provides evidence that it is likewise an ideal target for a cancer vaccine approach to prostate cancer, as its expression is not detected in most normal tissues.

          Accordingly, therapeutic approaches targeting particular motifs of STEAP, or aimed at inhibiting the activity of the STEAP protein, are expected to be useful for 15   patients suffering from prostate cancer and other cancers expressing STEAP. The therapeutic approaches aimed at inhibiting the activity of the STEAP protein generally fall into two classes. One class comprises various methods for inhibiting the binding or association of the STEAP protein with its binding partner or with other proteins. Another class comprises a variety of methods for inhibiting the transcription of the 20   STEAP gene or translation of STEAP mRNA.

#### **STEAP as a Target for Antibody-Based Therapy**

          The cell surface nature and expression profiles of the STEAPs in cancers including prostate cancer indicate that they are promising targets for antibody therapy of prostate and 25   other cancers expressing STEAPs. The experimental results described in the Examples herein provide compelling evidence that STEAP-1 and STEAP-2 are strongly expressed uniformly over the surface of glandular epithelial cells within prostate and prostate cancer cells. In particular, immunohistochemical analysis results show that the surface of human prostate epithelial cells (normal and cancer) appear to be uniformly coated with STEAP-1. 30   Biochemical analysis confirms the cell surface localization of STEAP-1 initially suggested

by its putative 6-transmembrane primary structural elements and by the pericellular staining plainly evident by immunohistochemical staining.

STEAP-1 and STEAP-2 are uniformly expressed at high levels over the surface of prostate glandular epithelia, an ideal situation for immunotherapeutic intervention strategies that target extracellular STEAP epitopes. Systemic administration of STEAP-immunoreactive compositions would be expected to result in extensive contact of the composition with prostate epithelial cells via binding to STEAP extracellular epitopes. Moreover, given the near absence of STEAP-1 protein expression in normal human tissues, there is ample reason to expect exquisite sensitivity without toxic, non-specific and/or non-target effects caused by the binding of the immunotherapeutic composition to STEAP-1 on non-target organs and tissues.

In addition to the high level expression of STEAP-1 in prostate and prostate cancer cells, STEAP-1 appears to be substantially over-expressed in a variety of other human cancers, including bladder, lung, colon, pancreatic and ovarian cancers. In particular, high level STEAP-1 mRNA expression is detected in all tested prostate cancer tissues and cell lines, and in most of the pancreatic, colon, and bladder cancer cell lines tested. High level expression of STEAP-1 is also observed in some ovarian cancer cell lines. Lower level expression is observed in some breast, testicular, and cervical cancer cell lines. Very high level expression is also detected in a Ewing sarcoma cell line. Applicants have shown that cell surface STEAP-1 protein is expressed in bladder, lung and colon cancers, while there is no detectable cell surface (or intracellular) STEAP-1 protein in normal colon and low expression in normal bladder. Antibodies specifically reactive with extracellular domains of STEAP-1 may be useful to treat these cancers systemically, either as toxin or therapeutic agent conjugates or as naked antibodies capable of inhibiting cell proliferation or function.

STEAP-2 protein is also expressed in prostate cancer, and in other cancers as well, including colon and lung cancers. STEAP-2 mRNA analysis by RT-PCR and northern blot show that expression is restricted to prostate in normal tissues, is also expressed in some prostate, pancreatic, colon, testicular, ovarian and other cancers. Therefore, antibodies reactive with STEAP-2 may be useful in the treatment of prostate and other cancers. Similarly, the expression of STEAP-3 and STEAP-4 (as well as other

STEAPs) may be associated with some cancers. Thus antibodies reactive with these STEAP family member proteins may also be useful therapeutically.

STEAP antibodies may be introduced into a patient such that the antibody binds to STEAP on the cancer cells and mediates the destruction of the cells and the tumor, inhibits  
5 the growth of the cells or the tumor, and/or eliminates STEAP function in the primary tumor, in circulating micrometastases, and/or in established metastases. The degree of tumor vascularization may provide guidance on which delivery approach is recommended. Similarly, the grade and/or stage of disease would be expected to provide useful information in this regard. For example, a higher grade, more advanced  
10 tumor may be more likely to seed metastases, suggesting systemic administration in order to treat or prevent the emergence of metastases. Mechanisms by which such antibodies exert a therapeutic effect may include complement-mediated cytotoxicity, antibody-dependent cellular cytotoxicity, modulating the physiologic function of STEAP, inhibiting ligand binding or signal transduction pathways, modulating tumor cell differentiation, altering  
15 tumor angiogenesis factor profiles, and/or by inducing apoptosis. STEAP antibodies conjugated to toxic or therapeutic agents may also be used therapeutically to deliver the toxic or therapeutic agent directly to STEAP-bearing tumor cells.

Cancer immunotherapy using anti-STEAP antibodies may follow the teachings generated from various approaches that have been successfully employed in the  
20 treatment of other types of cancer, including but not limited to colon cancer (Arlen et al., 1998, Crit. Rev. Immunol. 18:133-138), multiple myeloma (Ozaki et al., 1997, Blood 90:3179-3186; Tsunenari et al., 1997, Blood 90:2437-2444), gastric cancer (Kasprzyk et al., 1992, Cancer Res. 52:2771-2776), B-cell lymphoma (Funakoshi et al., 1996, J. Immunother. Emphasis Tumor Immunol. 19:93-101), leukemia (Zhong et al., 1996, Leuk. Res. 20:581-589), colorectal cancer (Moun et al., 1994, Cancer Res. 54:6160-6166);  
25 Velders et al., 1995, Cancer Res. 55:4398-4403), and breast cancer (Shepard et al., 1991, J. Clin. Immunol. 11:117-127). Some therapeutic approaches involve conjugation of naked antibody to a toxin, such as the conjugation of <sup>131</sup>I to anti-CD20 antibodies (e.g., Bexxar, Coulter Pharmaceutical), while others involve co-administration of antibodies and other  
30 therapeutic agents, such as Herceptin<sup>TM</sup> (trastuzumab) with paclitaxel (Genentech, Inc.).

For treatment of prostate cancer, for example, STEAP antibodies can be administered in conjunction with radiation, chemotherapy or hormone ablation.

Although STEAP antibody therapy may be useful for all stages of cancer, antibody therapy may be particularly appropriate in advanced or metastatic cancers.

5 Treatment with the antibody therapy of the invention may be indicated for patients who have received previously one or more chemotherapy, while combining the antibody therapy of the invention with a chemotherapeutic or radiation regimen may be preferred for patients who have not received chemotherapeutic treatment. Additionally, antibody therapy may enable the use of reduced dosages of concomitant chemotherapy,  
10 particularly for patients who do not tolerate the toxicity of the chemotherapeutic agent very well.

It may be desirable for some cancer patients to be evaluated for the presence and level of STEAP expression, preferably using immunohistochemical assessments of tumor tissue, quantitative STEAP imaging, or other techniques capable of reliably indicating the  
15 presence and degree of STEAP expression. Immunohistochemical analysis of tumor biopsies or surgical specimens may be preferred for this purpose. Methods for immunohistochemical analysis of tumor tissues are well known in the art.

Anti-STEAP monoclonal antibodies useful in treating prostate and other cancers include those that are capable of initiating a potent immune response against the tumor  
20 and those that are capable of direct cytotoxicity. In this regard, anti-STEAP monoclonal antibodies (mAbs) may elicit tumor cell lysis by either complement-mediated or antibody-dependent cell cytotoxicity (ADCC) mechanisms, both of which require an intact Fc portion of the immunoglobulin molecule for interaction with effector cell Fc receptor sites or complement proteins. In addition, anti-STEAP mAbs that exert a direct  
25 biological effect on tumor growth are useful in the practice of the invention. Potential mechanisms by which such directly cytotoxic mAbs may act include inhibition of cell growth, modulation of cellular differentiation, modulation of tumor angiogenesis factor profiles, and the induction of apoptosis. The mechanism by which a particular anti-STEAP mAb exerts an anti-tumor effect may be evaluated using any number of in vitro  
30 assays designed to determine ADCC, ADMMC, complement-mediated cell lysis, and so forth, as is generally known in the art.

The anti-tumor activity of a particular anti-STEAP mAb, or combination of anti-STEAP mAbs, may be evaluated in vivo using a suitable animal model. For example, xenogenic prostate cancer models wherein human prostate cancer explants or passaged xenograft tissues are introduced into immune compromised animals, such as nude or SCID mice, are appropriate in relation to prostate cancer and have been described (U.S. Patent No. 6,107,540; Klein et al., 1997, Nature Medicine 3: 402-408). For Example, PCT Patent Application WO98/16628, Sawyers et al., published April 23, 1998, describes various xenograft models of human prostate cancer capable of recapitulating the development of primary tumors, micrometastasis, and the formation of osteoblastic metastases characteristic of late stage disease. Efficacy may be predicted using assays which measure inhibition of tumor formation, tumor regression or metastasis, and the like.

The use of murine or other non-human monoclonal antibodies, or human/mouse chimeric mAbs may induce moderate to strong immune responses in some patients. In some cases, this will result in clearance of the antibody from circulation and reduced efficacy. In the most severe cases, such an immune response may lead to the extensive formation of immune complexes that, potentially, can cause renal failure. Accordingly, preferred monoclonal antibodies used in the practice of the therapeutic methods of the invention are those that are either fully human or humanized and that bind specifically to the target STEAP antigen with high affinity but exhibit low or no antigenicity in the patient.

Therapeutic methods of the invention contemplate the administration of single anti-STEAP mAbs as well as combinations, or cocktails, of different mAbs. Such mAb cocktails may have certain advantages inasmuch as they contain mAbs that target different epitopes, exploit different effector mechanisms or combine directly cytotoxic mAbs with mAbs that rely on immune effector functionality. Such mAbs in combination may exhibit synergistic therapeutic effects. In addition, the administration of anti-STEAP mAbs may be combined with other therapeutic agents, including but not limited to various chemotherapeutic agents, androgen-blockers, and immune modulators (e.g., IL-2, GM-CSF). The anti-STEAP mAbs may be administered in their "naked" or unconjugated form, or may have therapeutic agents conjugated to them.

The anti-STEAP monoclonal antibodies used in the practice of the method of the invention may be formulated into pharmaceutical compositions comprising a carrier suitable for the desired delivery method. Suitable carriers include any material which when combined with the anti-STEAP mAbs retains the anti-tumor function of the antibody and is non-reactive with the subject's immune systems. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like.

The anti-STEAP antibody formulations may be administered via any route capable of delivering the antibodies to the tumor site. Potentially effective routes of administration include, but are not limited to, intravenous, intraperitoneal, intramuscular, intratumor, intradermal, and the like. The preferred route of administration is by intravenous injection. A preferred formulation for intravenous injection comprises the anti-STEAP mAbs in a solution of preserved bacteriostatic water, sterile unpreserved water, and/or diluted in polyvinylchloride or polyethylene bags containing 0.9% sterile Sodium Chloride for Injection, USP. The anti-STEAP mAb preparation may be lyophilized and stored as a sterile powder, preferably under vacuum, and then reconstituted in bacteriostatic water containing, for example, benzyl alcohol preservative, or in sterile water prior to injection.

Treatment will generally involve the repeated administration of the anti-STEAP antibody preparation via an acceptable route of administration such as intravenous injection (IV), typically at a dose in the range of about 0.1 to about 10 mg/kg body weight. Doses in the range of 10-500 mg mAb per week may be effective and well tolerated. Based on clinical experience with the Herceptin mAb in the treatment of metastatic breast cancer, an initial loading dose of approximately 4 mg/kg patient body weight IV followed by weekly doses of about 2 mg/kg IV of the anti-STEAP mAb preparation may represent an acceptable dosing regimen. Preferably, the initial loading dose is administered as a 90 minute or longer infusion. The periodic maintenance dose may be administered as a 30 minute or longer infusion, provided the initial dose was well tolerated. However, as one of skill in the art will understand, various factors will influence the ideal dose regimen in a particular case. Such factors may include, for example, the binding affinity and half life of the mAb or mAbs used, the degree of STEAP overexpression in the patient, the extent of circulating shed STEAP antigen, the

desired steady-state antibody concentration level, frequency of treatment, and the influence of chemotherapeutic agents used in combination with the treatment method of the invention.

5 Optimally, patients should be evaluated for the level of circulating shed STEAP antigen in serum in order to assist in the determination of the most effective dosing regimen and related factors. Such evaluations may also be used for monitoring purposes throughout therapy, and may be useful to gauge therapeutic success in combination with evaluating other parameters (such as serum PSA levels in prostate cancer therapy).

#### 10 **Inhibition of STEAP Protein Function**

The invention includes various methods and compositions for inhibiting the binding of STEAP to its binding partner or ligand, or its association with other protein(s) as well as methods for inhibiting STEAP function.

#### 15 *Inhibition of STEAP With Recombinant Proteins*

In one approach, recombinant molecules that are capable of binding to STEAP thereby preventing STEAP from accessing/binding to its binding partner(s) or associating with other protein(s) are used to inhibit STEAP function. Such recombinant molecules may, for example, contain the reactive part(s) of a STEAP specific antibody  
20 molecule. In a particular embodiment, the STEAP binding domain of a STEAP binding partner may be engineered into a dimeric fusion protein comprising two STEAP ligand binding domains linked to the Fc portion of a human IgG, such as human IgG1. Such IgG portion may contain, for example, the C<sub>H</sub>2 and C<sub>H</sub>3 domains and the hinge region, but not the C<sub>H</sub>1 domain. Such dimeric fusion proteins may be administered in soluble form to  
25 patients suffering from a cancer associated with the expression of STEAP, including but not limited to prostate cancer, where the dimeric fusion protein specifically binds to STEAP thereby blocking STEAP interaction with a binding partner and/or modulating STEAP function. Such dimeric fusion proteins may be further combined into multimeric proteins using known antibody linking technologies.

30

*Inhibition of STEAP With Intracellular Antibodies*

In another approach, recombinant vectors encoding single chain antibodies that specifically bind to STEAP may be introduced into STEAP expressing cells via gene transfer technologies, wherein the encoded single chain anti-STEAP antibody is

5 expressed intracellularly, binds to STEAP protein, and thereby inhibits its function. Methods for engineering such intracellular single chain antibodies are well known. Such intracellular antibodies, also known as “intrabodies”, may be specifically targeted to a particular compartment within the cell, providing control over where the inhibitory activity of the treatment will be focused. This technology has been successfully applied  
10 in the art (for review, see Richardson and Marasco, 1995, TIBTECH vol. 13). Intrabodies have been shown to virtually eliminate the expression of otherwise abundant cell surface receptors. See, for example, Richardson et al., 1995, Proc. Natl. Acad. Sci. USA 92: 3137-3141; Beerli et al., 1994, J. Biol. Chem. 269: 23931-23936; Deshane et al., 1994, Gene Ther. 1: 332-337.

15 Single chain antibodies comprise the variable domains of the heavy and light chain joined by a flexible linker polypeptide, and are expressed as a single polypeptide. Optionally, single chain antibodies may be expressed as a single chain variable region fragment joined to the light chain constant region. Well known intracellular trafficking signals may be engineered into recombinant polynucleotide vectors encoding such single  
20 chain antibodies in order to precisely target the expressed intrabody to the desired intracellular compartment. For example, intrabodies targeted to the endoplasmic reticulum (ER) may be engineered to incorporate a leader peptide and, optionally, a C-terminal ER retention signal, such as the KDEL amino acid motif. Intrabodies intended to exert activity in the nucleus may be engineered to include a nuclear localization signal.  
25 Lipid moieties may be joined to intrabodies in order to tether the intrabody to the cytosolic side of the plasma membrane. Intrabodies may also be targeted to exert function in the cytosol. For example, cytosolic intrabodies may be used to sequester factors within the cytosol, thereby preventing them from being transported to their natural cellular destination.

30 In one embodiment, STEAP intrabodies are designed to bind specifically to a particular STEAP domain. For example, cytosolic intrabodies that specifically bind to



the STEAP protein may be used to prevent STEAP related molecules from gaining access to the nucleus, thereby preventing it from exerting any biological activity within the nucleus.

5 In order to direct the expression of such intrabodies specifically to particular tumor cells, the transcription of the intrabody may be placed under the regulatory control of an appropriate tumor-specific promoter and/or enhancer. In order to target intrabody expression specifically to prostate, for example, the PSA promoter and/or promoter/enhancer may be utilized (See, for example, U.S. Patent No. 5,919,652).

#### 10 **Inhibition of STEAP Transcription or Translation**

Within another class of therapeutic approaches, the invention provides various methods and compositions for inhibiting the transcription of the STEAP gene. Similarly, the invention also provides methods and compositions for inhibiting the translation of STEAP mRNA into protein.

15 In one approach, a method of inhibiting the transcription of the STEAP gene comprises contacting the STEAP gene with a STEAP antisense polynucleotide. In another approach, a method of inhibiting STEAP mRNA translation comprises contacting the STEAP mRNA with an antisense polynucleotide. In another approach, a STEAP specific ribozyme may be used to cleave the STEAP message, thereby inhibiting  
20 translation. Such antisense and ribozyme based methods may also be directed to the regulatory regions of the STEAP gene, such as the STEAP promoter and/or enhancer elements. Similarly, proteins capable of inhibiting a STEAP gene transcription factor may be used to inhibit STEAP mRNA transcription. The various polynucleotides and compositions useful in the aforementioned methods have been described above. The  
25 use of antisense and ribozyme molecules to inhibit transcription and translation is well known in the art.

Other factors that inhibit the transcription of STEAP through interfering with STEAP transcriptional activation may also be useful for the treatment of cancers expressing STEAP. Similarly, factors that are capable of interfering with STEAP  
30 processing may be useful for the treatment of cancers expressing STEAP. Cancer treatment methods utilizing such factors are also within the scope of the invention.

### **General Considerations for Therapeutic Strategies**

Gene transfer and gene therapy technologies may be used for delivering therapeutic polynucleotide molecules to tumor cells synthesizing STEAP (i.e., antisense, ribozyme, polynucleotides encoding intrabodies and other STEAP inhibitory molecules). A number of gene therapy approaches are known in the art. Recombinant vectors encoding STEAP antisense polynucleotides, ribozymes, factors capable of interfering with STEAP transcription, and so forth, may be delivered to target tumor cells using such gene therapy approaches.

The above therapeutic approaches may be combined with any one of a wide variety of chemotherapy or radiation therapy regimens. These therapeutic approaches may also enable the use of reduced dosages of chemotherapy and/or less frequent administration, particularly in patients that do not tolerate the toxicity of the chemotherapeutic agent well.

The anti-tumor activity of a particular composition (e.g., antisense, ribozyme, intrabody), or a combination of such compositions, may be evaluated using various in vitro and in vivo assay systems. In vitro assays for evaluating therapeutic potential include cell growth assays, soft agar assays and other assays indicative of tumor promoting activity, binding assays capable of determining the extent to which a therapeutic composition will inhibit the binding of STEAP to a binding partner, etc.

In vivo, the effect of a STEAP therapeutic composition may be evaluated in a suitable animal model. For example, xenogeneic prostate cancer models wherein human prostate cancer explants or passaged xenograft tissues are introduced into immune compromised animals, such as nude or SCID mice, are appropriate in relation to prostate cancer and have been described (Klein et al., 1997, Nature Medicine 3: 402-408). For example, PCT Patent Application WO98/16628, Sawyers et al., published April 23, 1998, describes various xenograft models of human prostate cancer capable of recapitulating the development of primary tumors, micrometastasis, and the formation of osteoblastic metastases characteristic of late stage disease. Efficacy may be predicted using assays that measure inhibition of tumor formation, tumor regression or metastasis, and the like. See, also, the Examples below.

In vivo assays that qualify the promotion of apoptosis may also be useful in evaluating potential therapeutic compositions. In one embodiment, xenografts from bearing mice treated with the therapeutic composition may be examined for the presence of apoptotic foci and compared to untreated control xenograft-bearing mice. The extent to which apoptotic foci are found in the tumors of the treated mice provides an indication of the therapeutic efficacy of the composition.

The therapeutic compositions used in the practice of the foregoing methods may be formulated into pharmaceutical compositions, including vaccine compositions, comprising a carrier suitable for the desired delivery method. Suitable carriers include any material that when combined with the therapeutic composition retains the anti-tumor function of the therapeutic composition and is non-reactive with the patient's immune system. Examples include, but are not limited to, any of a number of standard pharmaceutical carriers such as sterile phosphate buffered saline solutions, bacteriostatic water, and the like (see, generally, Remington's Pharmaceutical Sciences 16<sup>th</sup> Edition, A. Osal., Ed., 1980).

Therapeutic formulations may be solubilized and administered via any route capable of delivering the therapeutic composition to the tumor site. Potentially effective routes of administration include, but are not limited to, intravenous, parenteral, intraperitoneal, intramuscular, intratumor, intradermal, intraorgan, orthotopic, and the like. A preferred formulation for intravenous injection comprises the therapeutic composition in a solution of preserved bacteriostatic water, sterile unpreserved water, and/or diluted in polyvinylchloride or polyethylene bags containing 0.9% sterile Sodium Chloride for Injection, USP. Therapeutic protein preparations may be lyophilized and stored as sterile powders, preferably under vacuum, and then reconstituted in bacteriostatic water containing, for example, benzyl alcohol preservative, or in sterile water prior to injection.

Dosages and administration protocols for the treatment of cancers using the foregoing methods will vary with the method and the target cancer and will generally depend on a number of other factors appreciated in the art.

30

## CANCER VACCINES

The invention further provides cancer vaccines comprising a STEAP protein or fragment thereof, as well as DNA based vaccines. In view of the prostate- and tumor-restricted expression of STEAP, STEAP cancer vaccines are expected to be effective at specifically preventing and/or treating STEAP expressing cancers without creating non-specific effects on non-target tissues. The use of a tumor antigen in a vaccine for generating humoral and cell-mediated immunity for use in anti-cancer therapy is well known in the art and has been employed in prostate cancer using human PSMA and rodent PAP immunogens (Hodge et al., 1995, Int. J. Cancer 63: 231-237; Fong et al., 1997, J. Immunol. 159: 3113-3117). Such methods can be readily practiced by employing a STEAP protein, or fragment thereof, or a STEAP-encoding nucleic acid molecule and recombinant vectors capable of expressing and appropriately presenting the STEAP immunogen.

For example, viral gene delivery systems may be used to deliver a STEAP-encoding nucleic acid molecule. Various viral gene delivery systems that can be used in the practice of this aspect of the invention include, but are not limited to, vaccinia, fowlpox, canarypox, adenovirus, influenza, poliovirus, adeno-associated virus, lentivirus, and sindbus virus (Restifo, 1996, Curr. Opin. Immunol. 8: 658-663). Non-viral delivery systems may also be employed by using naked DNA encoding a STEAP protein or fragment thereof introduced into the patient (e.g., intramuscularly) to induce an anti-tumor response. In one embodiment, the full-length human STEAP cDNA may be employed.

In one embodiment, a STEAP cancer vaccine is based on the identification of immunogenic peptides within a STEAP amino acid sequence shown in Fig. 11A-B. As discussed further in the examples below, STEAPs have been shown to induce T and B cell responses. STEAP-1 and STEAP-2 polypeptides have been used to generate an immune response in mice and rabbits for the production of monoclonal and polyclonal antibodies. Thus, specific portions of STEAP, and polynucleotides encoding these portions, may be selected for the production of a cancer vaccine. One example of such a portion of a STEAP protein is amino acid residues 14 through 28 of the STEAP-1 amino acid sequence as shown in FIG. 1A-B (WKMKPRRNLEEDDYLY; SEQ ID NO: 22).

In another embodiment, STEAP nucleic acid molecules encoding specific cytotoxic T lymphocyte (CTL) epitopes may be employed. CTL epitopes can be determined using

specific algorithms (e.g., Epimer, Brown University) to identify peptides within a STEAP protein that are capable of optimally binding to specified HLA alleles. One suitable algorithm is the HLA Peptide Motif Search algorithm available at the Bioinformatics and Molecular Analysis Section (BIMAS) web site (<http://bimas.dcrt.nih.gov/>). This algorithm is based on binding of specific peptide sequences in the groove of HLA Class I molecules and specifically HLA-A2 (Falk et al., 1991, Nature 351:290-6; Hunt et al., 1992, Science 255:1261-3; Parker et al., 1992, J. Immunol. 149:3580-7; Parker et al., 1994, J. Immunol. 152:163-75). The HLA Peptide Motif Search algorithm allows location and ranking of 8-mer, 9-mer, and 10-mer peptides from a complete protein sequence for predicted binding to HLA-A2 as well as other Class I molecules. Most HLA-A2 binding peptides are 9-mers, favorably containing a leucine at position 2 and a valine or leucine at position 9 (Parker et al., 1992, J. Immunol. 149:3580-7). Actual binding of peptides to HLA-A2 can be evaluated by stabilization of HLA-A2 expression on the antigen processing defective cell line T2 (Xue et al., 1997, Prostate 30:73-8; Peshwa et al., 1998, Prostate 36:129-38). Immunogenicity of specific peptides can be evaluated *in vitro* by stimulation of CD8+ CTL in the presence of dendritic cells (Xue et al.; Peshwa et al., *supra*).

Specific STEAP peptides predicted to bind HLA-A2 and preferred for use in cancer vaccines include peptides corresponding to amino acids 165-173 of STEAP-1, amino acids 86-94 of STEAP-1, amino acids 262-270 of STEAP-1, amino acids 302-310 of STEAP-1, amino acids 158-166 of STEAP-1, amino acids 227-235 of STEAP-2, amino acids 402-410 of STEAP-2, amino acids 307-315 of STEAP-2, amino acids 306-314 of STEAP-2, and amino acids 100-108 of STEAP-2.

Various *ex vivo* strategies may also be employed. One approach involves the use of dendritic cells to present STEAP antigen to a patient's immune system. Dendritic cells express MHC class I and II, B7 co-stimulator, and IL-12, and are thus highly specialized antigen presenting cells. In prostate cancer, autologous dendritic cells pulsed with peptides of the prostate-specific membrane antigen (PSMA) are being used in a Phase I clinical trial to stimulate prostate cancer patients' immune systems (Tjoa et al., 1996, Prostate 28: 65-69; Murphy et al., 1996, Prostate 29: 371-380). Dendritic cells can be used to present STEAP peptides to T cells in the context of MHC class I and II molecules. In one embodiment, autologous dendritic cells are pulsed with STEAP peptides capable of binding to MHC molecules. In another embodiment, dendritic cells

are pulsed with the complete STEAP protein. Yet another embodiment involves engineering the overexpression of the STEAP gene in dendritic cells using various implementing vectors known in the art, such as adenovirus (Arthur et al., 1997, Cancer Gene Ther. 4: 17-25), retrovirus (Henderson et al., 1996, Cancer Res. 56: 3763-3770),  
5 lentivirus, adeno-associated virus, DNA transfection (Ribas et al., 1997, Cancer Res. 57: 2865-2869), and tumor-derived RNA transfection (Ashley et al., 1997, J. Exp. Med. 186: 1177-1182). Cells expressing STEAP may also be engineered to express immune modulators, such as GM-CSF, and used as immunizing agents.

Anti-idiotypic anti-STEAP antibodies can also be used in anti-cancer therapy as a  
10 vaccine for inducing an immune response to cells expressing a STEAP protein. Specifically, the generation of anti-idiotypic antibodies is well known in the art and can readily be adapted to generate anti-idiotypic anti-STEAP antibodies that mimic an epitope on a STEAP protein (see, for example, Wagner et al., 1997, Hybridoma 16: 33-40; Foon et al., 1995, J Clin Invest 96: 334-342; Herlyn et al., 1996, Cancer Immunol Immunother 43:  
15 65-76). Such an anti-idiotypic antibody can be used in cancer vaccine strategies.

Genetic immunization methods may be employed to generate prophylactic or therapeutic humoral and cellular immune responses directed against cancer cells expressing STEAP. Constructs comprising DNA encoding a STEAP protein/immunogen and appropriate regulatory sequences may be injected directly into muscle or skin of an  
20 individual, such that the cells of the muscle or skin take up the construct and express the encoded STEAP protein/immunogen. Expression of the STEAP protein immunogen results in the generation of prophylactic or therapeutic humoral and cellular immunity against prostate and other STEAP-expressing cancers. Various prophylactic and therapeutic genetic immunization techniques known in the art may be used (for review, see  
25 information and references published at Internet address [www.genweb.com](http://www.genweb.com)).

### **DIAGNOSTIC COMPOSITIONS AND KITS**

For use in the diagnostic and therapeutic applications described or suggested above, kits are also provided by the invention. Such kits may comprise a carrier means  
30 being compartmentalized to receive in close confinement one or more container means such as vials, tubes, and the like, each of the container means comprising one of the

separate elements to be used in the method. For example, one of the container means may comprise a probe that is or can be detectably labeled. Such probe may be an antibody or polynucleotide specific for a STEAP protein or a STEAP gene or message, respectively. Where the kit utilizes nucleic acid hybridization to detect the target nucleic acid, the kit may also have containers containing nucleotide(s) for amplification of the target nucleic acid sequence and/or a container comprising a reporter-means, such as a biotin-binding protein, such as avidin or streptavidin, bound to a reporter molecule, such as an enzymatic, florescent, or radioisotope label.

The kit of the invention will typically comprise the container described above and one or more other containers comprising materials desirable from a commercial and user standpoint, including buffers, diluents, filters, needles, syringes, and package inserts with instructions for use. A label may be present on the on the container to indicate that the composition is used for a specific therapy or non-therapeutic application, and may also indicate directions for either *in vivo* or *in vitro* use, such as those described above.

Accordingly, the invention also provides diagnostic compositions comprising STEAP-related molecules. Such molecules include the various STEAP polynucleotides, primers, probes, proteins, fragments, antibodies described herein. The molecules included in the diagnostic composition may optionally be labeled with a detectable marker. STEAP diagnostic compositions may further comprise appropriate buffers, diluents, and other ingredients as desired.

## **EXAMPLES**

Various aspects of the invention are further described and illustrated by way of the several examples that follow, none of which are intended to limit the scope of the invention.

### **Example 1: SSH-Generated Isolation of cDNA Fragment of the STEAP Gene**

#### *Materials and Methods*

#### **LAPC Xenografts:**

LAPC xenografts were obtained from Dr. Charles Sawyers (UCLA) and generated as described (Klein et al, 1997, Nature Med. 3: 402-408; Craft et al., 1999,

Cancer Res. 59: 5030-5036). Androgen dependent and independent LAPC-4 xenografts (LAPC-4 AD and AI, respectively) and LAPC-9 xenografts (LAPC-9 AD and AI, respectively) were grown in intact male SCID mice or in castrated males, respectively, and were passaged as small tissue chunks in recipient males. LAPC-4 AI xenografts were  
5 derived from LAPC-4 AD tumors and LAPC-9 AI xenografts were derived from LAPC-9 AD tumors. To generate the AI xenografts, male mice bearing LAPC AD tumors were castrated and maintained for 2-3 months. After the LAPC tumors re-grew, the tumors were harvested and passaged in castrated males or in female SCID mice.

LAPC-4 AD xenografts were grown intratibially as follows. LAPC-4 AD  
10 xenograft tumor tissue grown subcutaneously was minced into 1-2 mm<sup>3</sup> sections while the tissue was bathed in 1X Iscoves medium, minced tissue was then centrifuged at 1.3K rpm for 4 minutes, the supernatant was resuspended in 10 ml ice cold 1X Iscoves medium and centrifuged at 1.3K rpm for 4 minutes. The pellet was then resuspended in 1X Iscoves with 1% pronase E and incubated for 20 minutes at room temperature with  
15 mild rocking agitation followed by incubation on ice for 2-4 minutes. Filtrate was centrifuged at 1.3K rpm for 4 minutes, and the pronase was removed from the aspirated pellet by resuspending in 10 ml Iscoves and re-centrifuging. Clumps of cells were then plated in PrEGM medium and grown overnight. The cells were then harvested, filtered, washed 2X RPMI, and counted. Approximately 50,000 cells were mixed with an equal  
20 volume of ice-cold Matrigel on ice, and surgically injected into the proximal tibial metaphyses of SCID mice via a 27 gauge needle. After 10-12 weeks, LAPC-4 tumors growing in bone marrow were recovered.

#### Cell Lines and Tissues:

25 Human cell lines (e.g., HeLa) were obtained from the ATCC and were maintained in DMEM with 5% fetal calf serum. Human tissues for RNA and protein analyses were obtained from the Human Tissue Resource Center (HTRC) at the UCLA (Los Angeles, CA) and from QualTek, Inc. (Santa Barbara, CA).

#### 30 RNA Isolation:

Tumor tissue and cell lines were homogenized in Trizol reagent (Life Technologies, Gibco BRL) using 10 ml/ g tissue or 10 ml/ 10<sup>8</sup> cells to isolate total RNA.



Poly A RNA was purified from total RNA using Qiagen's Oligotex mRNA Mini and Midi kits. Total and mRNA were quantified by spectrophotometric analysis (O.D. 260/280 nm) and analyzed by gel electrophoresis.

5 Oligonucleotides:

The following HPLC purified oligonucleotides were used.

DPNCDN (cDNA synthesis primer):

5'TTTT**GATCAAGC**TT<sub>30</sub>3' (SEQ ID NO: 23)

10 Adaptor 1:

5'CTAATACGACTCACTATAGGGCTCGAGCGGCCGCCCCGGGCAG3'  
3'GGCCCGTC**CCTAG**5' (SEQ ID NO: 24,25)

Adaptor 2:

15 5'GTAATACGACTCACTATAGGGCAGCGTGGTCGCGGCCGAG3'  
3'CGGCTC**CCTAG**5' (SEQ ID NO: 26,27)

PCR primer 1:

5'CTAATACGACTCACTATAGGGC3' (SEQ ID NO: 28)

20 Nested primer (NP)1:

5'TCGAGCGGCCGCCCCGGGCAG**GA**3' (SEQ ID NO: 29)

Nested primer (NP)2:

5'AGCGTGGTCGCGGCCGAG**GA**3' (SEQ ID NO: 30)

25

Suppression Subtractive Hybridization:

Suppression Subtractive Hybridization (SSH) was used to identify cDNAs corresponding to genes which may be up-regulated in androgen dependent prostate cancer compared to benign prostatic hyperplasia (BPH).

30 Double stranded cDNAs corresponding to the LAPC-4 AD xenograft (tester) and the BPH tissue (driver) were synthesized from 2 µg of poly(A)+ RNA isolated from

xenograft and BPH tissue, as described above, using CLONTECH's PCR-Select cDNA Subtraction Kit and 1 ng of oligonucleotide RSACDN as primer. First- and second-strand synthesis were carried out as described in the Kit's user manual protocol (CLONTECH Protocol No. PT1117-1, Catalog No. K1804-1). The resulting cDNA  
5 was digested with Rsa I for 3 hrs. at 37°C. Digested cDNA was extracted with phenol/chloroform (1:1) and ethanol precipitated.

Driver cDNA (BPH) was generated by combining in a 4 to 1 ratio Rsa I digested BPH cDNA with digested cDNA from mouse liver, in order to ensure that murine genes were subtracted from the tester cDNA (LAPC-4 AD).

10 Tester cDNA (LAPC-4 AD) was generated by diluting 1 µl of Rsa I digested LAPC-4 AD cDNA (400 ng) in 5 µl of water. The diluted cDNA (2 µl, 160 ng) was then ligated to 2 µl of adaptor 1 and adaptor 2 (10 µM), in separate ligation reactions, in a total volume of 10 µl at 16°C overnight, using 400 u of T4 DNA ligase (CLONTECH). Ligation was terminated with 1 µl of 0.2 M EDTA and heating at 72°C for 5 min.

15 The first hybridization was performed by adding 1.5 µl (600 ng) of driver cDNA to each of two tubes containing 1.5 µl (20 ng) adaptor 1- and adaptor 2- ligated tester cDNA. In a final volume of 4 µl, the samples were overlaid with mineral oil, denatured in an MJ Research thermal cycler at 98°C for 1.5 minutes, and then were allowed to hybridize for 8 hrs at 68°C. The two hybridizations were then mixed together with an  
20 additional 1 µl of fresh denatured driver cDNA and were allowed to hybridize overnight at 68°C. The second hybridization was then diluted in 200 µl of 20 mM Hepes, pH 8.3, 50 mM NaCl, 0.2 mM EDTA, heated at 70°C for 7 min. and stored at -20°C.

#### PCR Amplification, Cloning and Sequencing of Gene Fragments Generated from SSH:

25 To amplify gene fragments resulting from SSH reactions, two PCR amplifications were performed. In the primary PCR reaction 1 µl of the diluted final hybridization mix was added to 1 µl of PCR primer 1 (10 µM), 0.5 µl dNTP mix (10 µM), 2.5 µl 10 x reaction buffer (CLONTECH) and 0.5 µl 50 x Advantage cDNA polymerase Mix (CLONTECH) in a final volume of 25 µl. PCR 1 was conducted using the following  
30 conditions: 75°C for 5 min., 94°C for 25 sec., then 27 cycles of 94°C for 10 sec, 66°C for

30 sec, 72°C for 1.5 min. Five separate primary PCR reactions were performed for each experiment. The products were pooled and diluted 1:10 with water. For the secondary PCR reaction, 1 µl from the pooled and diluted primary PCR reaction was added to the same reaction mix as used for PCR 1, except that primers NP1 and NP2 (10 µM) were used instead of PCR primer 1. PCR 2 was performed using 10-12 cycles of 94°C for 10 sec, 68°C for 30 sec, 72°C for 1.5 minutes. The PCR products were analyzed using 2% agarose gel electrophoresis.

The PCR products were inserted into pCR2.1 using the T/A vector cloning kit (Invitrogen). Transformed E. coli were subjected to blue/white and ampicillin selection. White colonies were picked and arrayed into 96 well plates and were grown in liquid culture overnight. To identify inserts, PCR amplification was performed on 1 ml of bacterial culture using the conditions of PCR1 and NP1 and NP2 as primers. PCR products were analyzed using 2% agarose gel electrophoresis.

Bacterial clones were stored in 20% glycerol in a 96 well format. Plasmid DNA was prepared, sequenced, and subjected to nucleic acid homology searches of the GenBank, dbEST, and NCI-CGAP databases.

#### RT-PCR Expression Analysis:

First strand cDNAs were generated from 1 µg of mRNA with oligo (dT)<sub>12-18</sub> priming using the Gibco-BRL Superscript Preamplification system. The manufacturers protocol was used and included an incubation for 50 min at 42°C with reverse transcriptase followed by RNase H treatment at 37°C for 20 min. After completing the reaction, the volume was increased to 200 µl with water prior to normalization. First strand cDNAs from 16 different normal human tissues were obtained from Clontech.

Normalization of the first strand cDNAs from multiple tissues was performed by using the primers 5'atatcgccgcgctcgtcgtcgacaa3' (SEQ ID NO: 31) and 5'agccacacgcagctcattgtagaagg 3' (SEQ ID NO: 32) to amplify β-actin. First strand cDNA (5 µl) was amplified in a total volume of 50 µl containing 0.4 µM primers, 0.2 µM each dNTPs, 1XPCR buffer (Clontech, 10 mM Tris-HCL, 1.5 mM MgCl<sub>2</sub>, 50 mM KCl, pH8.3) and 1X Klentaq DNA polymerase (Clontech). Five µl of the PCR reaction was removed at 18, 20, and 22 cycles and used for agarose gel electrophoresis. PCR was

performed using an MJ Research thermal cycler under the following conditions: initial denaturation was at 94°C for 15 sec, followed by a 18, 20, and 22 cycles of 94°C for 15, 65°C for 2 min, 72°C for 5 sec. A final extension at 72°C was carried out for 2 min. After agarose gel electrophoresis, the band intensities of the 283 bp  $\beta$ -actin bands from multiple tissues were compared by visual inspection. Dilution factors for the first strand cDNAs were calculated to result in equal  $\beta$ -actin band intensities in all tissues after 22 cycles of PCR. Three rounds of normalization were required to achieve equal band intensities in all tissues after 22 cycles of PCR.

To determine expression levels of the 8P1D4 gene, 5  $\mu$ l of normalized first strand cDNA was analyzed by PCR using 25, 30, and 35 cycles of amplification using the following primer pairs:

5' ACT TTG TTG ATG ACC AGG ATT GGA 3' (SEQ ID NO: 4)

5' CAG AAC TTC AGC ACA CAC AGG AAC 3' (SEQ ID NO: 5)

Semi quantitative expression analysis was achieved by comparing the PCR products at cycle numbers that give light band intensities.

### *Results*

Several SSH experiments were conducted as described in the Materials and Methods, supra, and led to the isolation of numerous candidate gene fragment clones. All candidate clones were sequenced and subjected to homology analysis against all sequences in the major public gene and EST databases in order to provide information on the identity of the corresponding gene and to help guide the decision to analyze a particular gene for differential expression. In general, gene fragments which had no homology to any known sequence in any of the searched databases, and thus considered to represent novel genes, as well as gene fragments showing homology to previously sequenced expressed sequence tags (ESTs), were subjected to differential expression analysis by RT-PCR and/or Northern analysis.

One of the cDNA clones, designated 8P1D4, was 436 bp in length and showed homology to an EST sequence in the NCI-CGAP tumor gene database. The full length

cDNA encoding the 8P1D4 gene was subsequently isolated using this cDNA and re-named STEAP-1. The 8P1D4 cDNA nucleotide sequence corresponds to nucleotide residues 150 through 585 in the STEAP-1 cDNA sequence as shown in FIG. 1A-B. Another clone, designated 28P3E1, 561 bp in length showed homology to a number of EST sequences in the NCI-CGAP tumor gene database or in other databases. Part of the 28P3E1 sequence (356 bp) is identical to an EST derived from human fetal tissue. After the full length STEAP-1 cDNA was obtained and sequenced, it became apparent that this clone also corresponds to STEAP-1 (more specifically, to residues 622 through the 3' end of the STEAP-1 nucleotide sequence as shown in FIG. 1A-B).

Differential expression analysis by RT-PCR using primers derived from the 8P1D4 cDNA clone showed that the 8P1D4 (STEAP-1) gene is expressed at approximately equal levels in normal prostate and the LAPC-4 and LAPC-9 xenografts (FIG. 2, panel A). Further RT-PCR expression analysis of first strand cDNAs from 16 normal tissues showed greatest levels of 8P1D4 expression in prostate. Substantially lower level expression in several other normal tissues (i.e., colon, ovary, small intestine, spleen and testis) was detectable only at 30 cycles of amplification in brain, pancreas, colon and small intestine (FIG. 2, panels B and C).

### **Example 2: Isolation of Full Length STEAP-1 Encoding cDNA**

The 436 bp 8P1D4 gene fragment (Example 1) was used to isolate additional cDNAs encoding the 8P1D4/STEAP-1 gene. Briefly, a normal human prostate cDNA library (Clontech) was screened with a labeled probe generated from the 436 bp 8P1D4 cDNA. One of the positive clones, clone 10, is 1195 bp in length and encodes a 339 amino acid protein having nucleotide and encoded amino acid sequences bearing no significant homology to any known human genes or proteins (homology to a rat Kidney Injury Protein recently described in International Application WO98/53071). The encoded protein contains at least 6 predicted transmembrane motifs implying a cell surface orientation (see FIG. 1A-B, predicted transmembrane motifs underlined). These structural features led to the designation "STEAP", for "Six Transmembrane Epithelial Antigen of the Prostate".

Subsequent identification of additional STEAP proteins led to the re-designation of the 8P1D4 gene product as "STEAP-1". The STEAP-1 cDNA and encoded amino acid sequences are shown in FIG. 1A-B and correspond to SEQ ID NOS: 1 and 2, respectively. STEAP-1 cDNA clone 10 was deposited with the American Type Culture  
5 Collection ("ATCC") (10801 University Blvd., Manassas, VA 20110-2209 USA) as plasmid 8P1D4 clone 10.1 on August 26, 1998 as ATCC Accession Number 98849. The STEAP-1 cDNA clone can be excised therefrom using EcoRI/XbaI double digest (EcoRI at the 5'end, XbaI at the 3'end).

### 10 **Example 3: STEAP-1 Gene and Protein Expression Analysis**

In order to begin to characterize the biological characteristics of STEAP-1, an extensive evaluation of STEAP-1 mRNA and STEAP-1 protein expression across a variety of human tissue specimens was undertaken. This evaluation included Northern blot, Western blot and immunohistochemical analysis of STEAP-1 expression in a large  
15 number of normal human tissues, human prostate cancer xenografts and cell lines, and various other human cancer cell lines.

#### **Example 3A: Northern Blot Analysis of STEAP-1 mRNA Expression in Normal Human Tissues**

20 Initial analysis of STEAP-1 mRNA expression in normal human tissues was conducted by northern blotting two multiple tissue blots obtained from Clontech (Palo Alto, California), comprising a total of 16 different normal human tissues, using labeled STEAP-1 clone 10 as a probe. RNA samples were quantitatively normalized with a  $\beta$ -actin probe. The results are shown in FIG. 3A. The highest expression level was  
25 detected in normal prostate, with an approximately 5-10 fold lower level of expression detected in colon and liver. These northern blots showed two transcripts of approximately 1.4 kb and 4.0 kb, the former of which corresponds to the full length STEAP-1 clone 10 cDNA, which encodes the entire STEAP-1 open reading frame. The larger transcript was separately cloned as a 3627 bp cDNA from a normal prostate  
30 library, the sequence of which contains a 2399 bp intron (FIG. 4).

This initial analysis was extended by using the STEAP-1 clone 10 probe to analyze an RNA dot blot matrix of 37 normal human tissues (Clontech, Palo Alto, CA; Human Master Blot™). The results are shown in FIG. 3B and show strong STEAP-1 expression only in prostate. Very low level STEAP-1 RNA expression was detected in liver, lung, trachea and fetal liver tissue, at perhaps a 5-fold lower level compared to prostate. No expression was detected in any of the remaining tissues. Based on these analyses, significant STEAP-1 expression appears to be prostate specific in normal tissues.

**Example 3B: Northern Blot Analysis of STEAP-1 mRNA Expression in Prostate Cancer Xenografts and Cell Lines**

To analyze STEAP-1 expression in human cancer tissues and cell lines, RNAs derived from human prostate cancer xenografts and an extensive panel of prostate and non-prostate cancer cell lines were analyzed by Northern blot using STEAP-1 cDNA clone 10 as probe. All RNA samples were quantitatively normalized by ethidium bromide staining and subsequent analysis with a labeled  $\beta$ -actin probe.

The results, presented in FIG. 5, show high level STEAP-1 expression in all the LAPC xenografts and all of the prostate cancer cell lines. Expression in the LAPC-9 xenografts was higher compared to the LAPC-4 xenografts, with no significant difference observed between androgen-dependent and androgen-independent sublines (FIG. 5A). Expression in the LAPC-4 xenografts was comparable to expression in normal prostate. Lower levels of expression were detected in PrEC cells (Clonetics), which represent the basal cell compartment of the prostate. Analysis of prostate cancer cell lines showed highest expression levels in LNCaP, an androgen dependent prostate carcinoma cell line. Significant expression was also detected in the androgen-independent cell lines PC-3 and DU145. High levels of STEAP expression were also detected in LAPC-4 and LAPC-9 tumors that were grown within the tibia of mice as a model of prostate cancer bone metastasis (FIG. 5B).

Significantly, very strong STEAP-1 expression was also detected in many of the non-prostate human cancer cell lines analyzed (FIG. 5A). Particularly high level expression was observed in RD-ES cells, an Ewing sarcoma (EWS) derived cell line.

Additionally, very high level expression was also detected in several of the colon cancer cell lines (e.g., CaCo-2, LoVo, T84 and Colo-205), bladder carcinoma cell lines (e.g., SCABER, UM-UC-3, TCCSUP and 5637), ovarian cancer cell lines (e.g., OV-1063 and SW 626) and pancreatic cancer cell lines (e.g., HPAC, Capan-1, PANC-1 and BxPC-3).  
5 These results, combined with the absence of strong expression in the corresponding normal tissues (FIG. 3), indicate that STEAP-1 may be generally up-regulated in these types (as well as other types) of human cancers.

10 **Example 3C: Western Blot Analysis of STEAP-1 Protein Expression in Prostate and Other Cancers**

A 15 mer peptide corresponding to amino acid residues 14 through 28 of the STEAP-1 amino acid sequence as shown in FIG. 1A-B (WKMKPRRNLEEDDYL; SEQ ID NO: 22) was synthesized and used to immunize sheep for the generation of sheep polyclonal antibodies towards the amino-terminus of the protein (anti-STEAP-1)  
15 as follows. The peptide was conjugated to KLH (keyhole limpet hemocyanin). The sheep was initially immunized with 400 µg of peptide in complete Freund's adjuvant. The animal was subsequently boosted every two weeks with 200 µg of peptide in incomplete Freund's adjuvant. Anti-STEAP antibody was affinity-purified from sheep serum using STEAP peptide coupled to affigel 10 (Bio Rad). Purified antibody is stored in phosphate-  
20 buffered saline with 0.1% sodium azide.

To test antibody specificity, the cDNA of STEAP-1 was cloned into a retroviral expression vector (pSRαtkneo, Muller et al., 1991, MCB 11:1785). NIH 3T3 cells were infected with retroviruses encoding STEAP-1 and were selected in G418 for 2 weeks. Western blot analysis of protein extracts of infected and uninfected NIH 3T3 cells  
25 showed expression of a protein with an apparent molecular weight of 36 kD only in the infected cells (FIG. 6, lanes marked "3T3 STEAP" AND "3T3").

The anti-STEAP-1 polyclonal antibody was used to probe western blots of cell lysates prepared from a variety of prostate cancer xenograft tissues, prostate cancer cell lines and other non-prostate cancer cell lines. Protein samples (20µg each) were  
30 quantitatively normalized by probing the blots with an anti-Grb-2 antibody.



The results are shown in FIG. 6. STEAP-1 protein was detected in all of the LAPC prostate cancer xenografts, all of the prostate cancer cell lines, a primary prostate cancer specimen and its matched normal prostate control. Highest STEAP-1 protein expression was detected in the LAPC-9 xenograft and in LNCaP cells, in agreement with the northern blot analysis described immediately above. High level expression was also observed in the bladder carcinoma cell line UM-UC-3. Expression in other cancer cell lines was also detectable (FIG. 6).

### 10 **Example 3D: Immunohistochemical Analysis of STEAP-1 Protein Expression in Prostate Tumor Biopsy and Surgical Specimens**

To determine the extent of STEAP-1 protein expression in clinical materials, tissue sections were prepared from a variety of prostate cancer biopsies and surgical samples for immunohistochemical analysis. Tissues were fixed in 10% formalin, embedded in paraffin, and sectioned according to standard protocol. Formalin-fixed, paraffin-embedded sections of LNCaP cells were used as a positive control. Sections were stained with a sheep anti-STEAP-1 polyclonal antibody directed against a STEAP-1 N-terminal epitope (as described immediately above). LNCaP sections were stained in the presence of an excess amount of the STEAP-1 N-terminal peptide immunogen used to generate the polyclonal antibody (peptide 1) or a non-specific peptide derived from a distinct region of the STEAP-1 protein (peptide 2; YQQVQQNKEDAWIEH; SEQ ID NO: 33).

The results are shown in FIG. 8. LNCaP cells showed uniformly strong pericellular staining in all cells (FIG. 8B). Excess STEAP N-terminal peptide (peptide 1) was able to competitively inhibit antibody staining (FIG. 8A), while peptide 2 had no effect (FIG. 8B). Similarly, uniformly strong pericellular staining was seen in the LAPC-9 (FIG. 8F) and LAPC-4 prostate cancer xenografts. These results are clear and suggest that the staining is STEAP specific. Moreover, these results show localization of STEAP to the plasma membrane, corroborating the biochemical findings presented in Example 4 below.

30 The results obtained with the various clinical specimens are shown in FIG. 8C (normal prostate tissue), FIG. 8D (grade 3 prostatic carcinoma), and FIG. 8E (grade 4 prostatic carcinoma), and are also included in the summarized results shown in Table 1.

Light to strong staining was observed in the glandular epithelia of all prostate cancer samples tested as well as in all samples derived from normal prostate or benign disease (e.g., BPH). The signal appears to be strongest at the cell membrane of the epithelial cells, especially at the cell-cell junctions (FIG. 8C-E) and is also inhibited with excess STEAP N-terminal peptide 1. Some basal cell staining is also seen in normal prostate (FIG. 8C), which is more apparent when examining atrophic glands. STEAP-1 seems to be expressed at all stages of prostate cancer since lower grades (FIG. 8D), higher grades (FIG. 8E) and metastatic prostate cancer (represented by LAPC-9; FIG. 8F; See also human patient metastases shown in FIG. 32A-B) all exhibit strong staining.

Immunohistochemical staining of a large panel of normal non-prostate tissues showed no detectable STEAP-1 expression in most of these normal tissues (Table 1). Normal bladder exhibited low levels of cell surface staining in the transitional epithelium (FIG. 8G). Pancreas, stomach, uterus, fallopian tubes and pituitary showed low levels of cytoplasmic staining (Table 1). It is unclear whether the observed cytoplasmic staining is specific or is due to non-specific binding of the antibody, since northern blotting showed little to no STEAP-1 expression in pancreas (FIG. 3). These results indicate that cell surface expression of STEAP-1 in normal tissues appears to be restricted to prostate and bladder.

**Table 1: Immunohistochemical Staining of Human Tissues With Anti-STEAP-1 Sheep Polyclonal Antibody**

Staining Intensity	Tissues*
<b>None</b>	Cerebellum, cerebral cortex, spinal cord, heart, skeletal muscle (2), artery, thymus, spleen (4), bone marrow, lymph node (3), lung (8), liver (4), ileum, kidney (2), testis (2), ovary, placenta, breast, adrenal gland (2), thyroid gland (2), skin (2)
<b>Light Cytoplasmic</b>	Ureter, bladder cancer (2/5), colon (4/7), colon cancer (4/7), fallopian tubes (2), pituitary gland, pancreas, stomach (1/2), uterus (1/2)

<b>Light Membrane</b>	Prostate cancer bone marrow metastasis (1/6), prostate cancer lymph node metastasis (1/5), lung cancer (1/6)
<b>Moderate to Strong Membrane</b>	Prostate (7/7), BPH (5/5), prostate cancer (20/20)**, prostate cancer lymph node metastasis (4/5), prostate cancer bone marrow metastasis (5/6), bladder (2/5), bladder cancer (3/5), lung cancer (3/6)

\*In cases where more than one sample was analyzed per tissue, the number in parenthesis indicates how many samples showed positive staining/total analyzed.

\*\*Prostate cancer grades varied from 3 to 5 and Gleason scores of 7 to 10.

#### 5     **Example 3E: Immunohistochemical Analysis of STEAP-1 Protein Expression in Bladder and Lung Cancer**

To determine the extent of STEAP-1 protein expression in other cancers, tissue from bladder and lung cancer clinical specimens were subjected to immunohistochemical analysis using the polyclonal antibody and methods as described in Example 3D. The results, shown in Figures 21A-D, reveal a pericellular staining pattern similar to that observed with prostate cancer tissue.

#### **Example 4: Biochemical Characterization of STEAP-1 Protein**

To initially characterize the STEAP-1 protein, cDNA clone 10 was cloned into the pcDNA 3.1 Myc-His plasmid (Invitrogen), which encodes a 6His tag at the carboxyl-terminus, transfected into 293T cells, and analyzed by flow cytometry using anti-His monoclonal antibody (His-probe, Santa Cruz) as well as the anti-STEAP-1 polyclonal antibody described above. Staining of cells was performed on intact cells as well as permeabilized cells. The results indicated that only permeabilized cells stained with both antibodies, suggesting that both termini of the STEAP-1 protein are localized intracellularly. It is therefore possible that one or more of the STEAP-1 protein termini are associated with intracellular organelles rather than the plasma membrane.

To determine whether STEAP-1 protein is expressed at the cell surface, intact STEAP-1-transfected 293T cells were labeled with a biotinylation reagent that does not

enter live cells. STEAP-1 was then immunoprecipitated from cell extracts using the anti-His and anti-STEAP antibodies. SV40 large T antigen, an intracellular protein that is expressed at high levels in 293T cells, and the endogenous cell surface transferrin receptor were immunoprecipitated as negative and positive controls, respectively. After immunoprecipitation, the proteins were transferred to a membrane and visualized with horseradish peroxidase-conjugated streptavidin. The results of this analysis are shown in FIG. 7. Only the transferrin receptor (positive control) and STEAP-1 were labeled with biotin, while the SV40 large T antigen (negative control) was not detectably labeled (FIG. 7A). Since only cell surface proteins are labeled with this technique, it is clear from these results that STEAP-1 is a cell surface protein. Combined with the results obtained from the flow cytometric analysis, it is clear that STEAP-1 is a cell surface protein with intracellular amino- and carboxyl- termini.

Furthermore, the above results together with the STEAP-1 secondary structural predictions, shows that STEAP-1 is a type IIIa membrane protein with a molecular topology of six potential transmembrane domains, 3 extracellular loops, 2 intracellular loops and two intracellular termini. A schematic representation of STEAP-1 protein topology relative to the cell membrane is shown in FIG. 1B.

In addition, prostate, bladder and colon cancer cells were directly analyzed for cell surface expression of STEAP-1 by biotinylation studies. Briefly, biotinylated cell surface proteins were affinity purified with streptavidin-gel and probed with the anti-STEAP-1 polyclonal antibody described above. Western blotting of the streptavidin purified proteins clearly show cell surface biotinylation of endogenous STEAP-1 in all prostate (LNCaP, PC-3, DU145), bladder (UM-UC-3, TCCSUP) and colon cancer (LoVo, Colo) cells tested, as well as in NIH 3T3 cells infected with a STEAP-1 encoding retrovirus, but not in non-expressing NIH 3T3 cells used as a negative control (FIG. 7B). In a further negative control, STEAP-1 protein was not detected in streptavidin precipitates from non-biotinylated STEAP expressing cells (FIG. 7B).

#### **Example 5: Constructs for Recombinant Expression of STEAP-1** *pGEX Constructs*

To express STEAP-1 in bacterial cells, portions of STEAP-1 were fused to the Glutathione S-transferase (GST) gene by cloning into pGEX-6P-1 (Amersham Pharmacia

Biotech, NJ). All constructs were made to generate recombinant STEAP-1 protein sequences with GST fused at the N-terminus and a six histidine epitope at the C-terminus. The six histidine epitope tag was generated by adding the histidine codons to the cloning primer at the 3' end of the ORF. A PreScission™ recognition site permits cleavage of the GST tag from STEAP-1. The ampicillin resistance gene and pBR322 origin permits selection and maintenance of the plasmid in *E. coli*. The following fragments of STEAP-1 were cloned into pGEX-6P-1:

Amino acids 148 to 251; amino acids 144 to 339; amino acids 39 to 253; amino acids 70 to 136; amino acids 254 to 313.

And into pGEX-4T: Amino acids 2 to 247; amino acids 135-318; and amino acids 144-339.

Exemplary additional constructs that can be made in pGEX-6P-1 spanning the following regions of the STEAP-1 protein include:

Amino acids 1 to 339; amino acids 1 to 144.

#### *pcDNA3.1/MycHis Construct*

To express STEAP-1 in mammalian cells, the 1,017 bp STEAP-1 ORF (with translational start Kozak consensus) was cloned into pcDNA3.1/MycHis\_Version B (Invitrogen, Carlsbad, CA). Protein expression is driven from the cytomegalovirus (CMV) promoter. The recombinant protein has the myc and six histidines fused to the C-terminus. The pcDNA3.1/MycHis vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Neomycin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*.

#### *pSRa Constructs*

To generate mammalian cell lines expressing STEAP-1 constitutively, the 1,017 bp ORF (with translational start Kozak consensus) was cloned into pSRa constructs. Amphotropic and ecotropic retroviruses are generated by transfection of pSRa constructs into the 293T-10A1 packaging line or co-transfection of pSRa and a helper

plasmid (φ-) in 293 cells, respectively. The retrovirus can be used to infect a variety of mammalian cell lines, resulting in the integration of the cloned gene, STEAP-1, into the host cell lines. Protein expression is driven from a long terminal repeat (LTR). The Neomycin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*. Additional pSRa constructs were made that fused the FLAG tag to the C and N-terminus to allow detection using anti-FLAG antibodies. The FLAG sequence 5' gat tac aag gat gac gac gat aag 3' (SEQ ID NO: 34) were added to cloning primer at the 3' end of the ORF. Another pSRa construct was made that contained amino acids 182 to 454. This truncated form of the STEAP-1 protein will help determine the function of the long extracellular N-Terminal region.

An Additional pSRa construct is being made to produce a myc/6 HIS fusion protein of the full length STEAP-1 protein.

#### 15 **Example 6: Identification and Structural Analysis of Other Human STEAPs**

STEAP-1 has no homology to any known human genes. In an attempt to identify additional genes that are homologous to STEAP-1, the protein sequence of STEAP-1 was used as an electronic probe to identify family members in the public EST (expression sequence tag) database (dbEST). Using the "tblastn" function in NCBI (National Center for Biotechnology Information), the dbEST database was queried with the STEAP-1 protein sequence. This analysis revealed additional putative STEAP-1 homologues or STEAP family members, as further described below.

In addition, SSH cloning experiments also identified a STEAP-1 related cDNA fragment, clone 98P4B6. This clone was isolated from SSH cloning using normal prostate cDNA as tester and LAPC-4 AD cDNA as driver. A larger partial sequence of the 98P4B6 clone was subsequently isolated from a normal prostate library; this clone encodes an ORF of 173 amino acids with close homology to the primary structure of STEAP-1, and thus was designated STEAP-2. A full length STEAP-2 cDNA of 2454 bp was isolated from a prostate library. The STEAP-2 nucleotide and encoded ORF amino acid sequences are shown in FIG. 9A-D. An amino acid alignment of the STEAP-1 and partial STEAP-2 primary structures is shown in FIGS. 11A-B and 11C. STEAP-1 and -2 share 61% identity over their 171 amino acid residue overlap (FIG. 11C). The STEAP-

2 cDNA has been deposited with the American Type Culture Collection ("ATCC"; 10801 University Blvd., Manassas, VA 20110-2209 USA) as plasmid 98P4B6-GTD3 on July 2, 1999 as ATCC Accession Number PTA-311.

The STEAP-2 cDNA (98P4B6-GTD3) contains a 355 bp 5'UTR (untranslated  
5 region) that is 72% GC rich, suggesting that it contains translational regulatory elements. The cDNA encodes an open reading frame (ORF) of 454 amino acids (a.a.) with six potential transmembrane domains. This is in contrast to STEAP-1, which is 339 a.a. in length. Alignment with STEAP-1 demonstrates 54.9% identity over a 237 amino acid overlap. Interestingly, the locations of the six putative transmembrane domains in  
10 STEAP-2 coincide with the locations of the transmembrane domains in STEAP-1 (see alignment). The homology of STEAP-2 with STEAP-1 is highest in the regions spanned by the first putative extracellular loop to the fifth transmembrane domain. This analysis and the sequence of STEAP-2 suggest some significant differences between STEAP-1 and STEAP-2: STEAP-2 exhibits a 205 a.a. long intracellular N-terminus (compared to  
15 69 a.a. in STEAP-1) and a short 4 a.a. intracellular C-terminus (compared to 26 a.a. in STEAP-1). These differences could imply significant differences in function and/or interaction with intracellular signaling pathways. To identify a unique mouse EST corresponding to STEAP-2, the unique N-terminus of STEAP-2 was used to query the dbEST database. One mouse EST was isolated (AI747886, mouse kidney) that may be  
20 used in the identification of mouse STEAP-2 and in expression analysis of STEAP-2 in mouse.

Two ESTs encoding ORFs bearing close homology to the STEAP-1 and STEAP-2 sequences were also identified by electronic probing with the STEAP-1 protein sequence. These ESTs (AI139607 and R80991) were provisionally designated  
25 STEAP-3 and STEAP-4. A full length cDNA encoding STEAP-3 was subsequently cloned, and its nucleotide and deduced amino acid sequences are shown in FIG. 10A-E. The STEAP-3 cDNA has been deposited with the American Type Culture Collection ("ATCC"; 10801 University Blvd., Manassas, VA 20110-2209 USA) on December 8, 1999 as plasmid pSTEAP-3 EBB4 as ATCC Accession Number PTA-1033. The  
30 nucleotide sequences of the ESTs corresponding to the STEAPs are reproduced in FIG. 10F.

An amino acid alignment of the structures of STEAP-1, STEAP-2, STEAP-3 and the partial sequence of the putative STEAP-4 is shown in FIG. 11A-B. This alignment shows a close structural similarity between all four STEAP family proteins, particularly in the predicted transmembrane domains. As indicated above, STEAP-1 and STEAP-2 demonstrate 54.9% identity over a 237 amino acid overlap. STEAP-1 and STEAP-3 are 40.9% identical over a 264 amino acid region, while STEAP-2 and STEAP-3 are 47.8% identical over a 416 amino acid region.

#### **Example 7: Constructs for Recombinant Expression of STEAP-2**

##### *pGEX Constructs*

To express STEAP-2 in bacterial cells, portions of STEAP-2 were fused to the Glutathione S-transferase (GST) gene by cloning into pGEX-6P-1 (Amersham Pharmacia Biotech, NJ). All constructs were made to generate recombinant STEAP-2 protein sequences with GST fused at the N-terminus and a six histidine epitope at the C-terminus. The six histidine epitope tag was generated by adding the histidine codons to the cloning primer at the 3' end of the ORF. A PreScission™ recognition site permits cleavage of the GST tag from STEAP-2. The ampicillin resistance gene and pBR322 origin permits selection and maintenance of the plasmid in E. coli. The following fragments of STEAP-2 were cloned into pGEX-6P-1:

Amino acids 287 to 390; amino acids 285 to 454; amino acids 193 to 454.

Additional exemplary constructs that can be made in pGEX-6P-1 spanning the following regions of the STEAP-2 protein include:

Amino acids 1 to 193; amino acids 1 to 454.

And in pGEX-4T: Amino acids 2-204; 183-387; and 276-453.

##### *pCRII Construct*

To generate sense and anti-sense riboprobes for RNA in situ investigations, a pCRII construct was generated using bp 367 to 877 of the GTD3 STEAP-2 cDNA. The pCRII vector has Sp6 and T7 promoters flanking the insert to drive the production of STEAP-2 RNA riboprobes which were used in RNA in situ hybridization experiments.



*pcDNA4/HisMax-TOPO Construct*

To express STEAP-2 in mammalian cells, the 1,362 bp STEAP-2 ORF was cloned into pcDNA4/HisMax-TOPO Version A (cat# K864-20, Invitrogen, Carlsbad, CA). Protein expression is driven from the cytomegalovirus (CMV) promoter and the SP163 translational enhancer. The recombinant protein has Xpress™ and six histidine epitopes fused to the N-terminus. In addition to this construct containing only the 1,362 bp ORF an additional construct was made with the C-terminus of the recombinant protein containing a 28 amino acid fusion resulting from vector sequences prior to the termination codon. The pcDNA4/HisMax-TOPO vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The Zeocin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in E. coli.

*pcDNA3.1/MycHis Construct*

To express STEAP-2 in mammalian cells, the 1,362 bp STEAP-2 ORF (with start Kozak consensus) was cloned into pcDNA3.1/MycHis\_Version A (Invitrogen, Carlsbad, CA). Protein expression is driven from the cytomegalovirus (CMV) promoter. The recombinant protein has the myc and six histidines fused to the C-terminus. The pcDNA3.1/MycHis vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The neomycin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in E. coli. Additional pcDNA3.1/MycHis constructs were generated using STEAP-2 amino acids 6 to 454, 121-454 and 182-454.

30

*pBlueBacHIS2a*

To express STEAP-1 in SF9 insect cells, the STEAP-1 ORF was cloned into pBlueBacHIS2A (Invitrogen, California). Protein expression is driven under the polyhedrin promoter. N-terminus poly HIS and Xpress tags allow for detection and purification of the recombinant STEAP-1 protein. A C-terminus enterokinase recognition site allows for cleavage of these tags. The ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*. Following cotransfection with AcMNPV DNA, a homologous recombination event occurs between these sequences resulting in a recombinant virus carrying the gene of interest and the polyhedrin. A variety of restriction enzyme sites for simplified subcloning are present. In addition, a reporter gene ( $\beta$ -galactosidase) conveniently identifies recombinant plaques, to eliminate tedious plaque screening.

*pcDNA3.1CT-GFP-TOPO Construct*

To express STEAP-2 in mammalian cells and to allow detection of the recombinant protein using fluorescence, the 1,362 bp ORF (with start Kozak consensus) was cloned into pcDNA3.1CT-GFP-TOPO (Invitrogen, CA). Protein expression is driven from the cytomegalovirus (CMV) promoter. The recombinant protein has the Green Fluorescent Protein (GFP) fused to the C-terminus facilitating non-invasive, *in vivo* detection and cell biology studies. The pcDNA3.1/MycHis vector also contains the bovine growth hormone (BGH) polyadenylation signal and transcription termination sequence to enhance mRNA stability along with the SV40 origin for episomal replication and simple vector rescue in cell lines expressing the large T antigen. The neomycin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in *E. coli*.

An additional construct with a N-terminal GFP fusion is being made in pcDNA3.1NT-GFP-TOPO spanning the entire length of the STEAP-2 protein.

*pSRa Constructs*

To generate mammalian cell lines expressing STEAP-2 constitutively, the 1362 bp ORF was cloned into pSRa constructs. Amphotropic and ecotropic retroviruses are

generated by transfection of pSRa constructs into the 293T-10A1 packaging line or co-transfection of pSRa and a helper plasmid ( $\phi$ -) in 293 cells, respectively. The retrovirus can be used to infect a variety of mammalian cell lines, resulting in the integration of the cloned gene, STEAP-2, into the host cell-lines. Protein expression is driven from a long terminal repeat (LTR). The neomycin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene and ColE1 origin permits selection and maintenance of the plasmid in E. coli. Additional pSRa constructs were made that fused the FLAG tag to the C and N-terminus to allow detection using anti-FLAG antibodies. The FLAG sequence 5' gat tac aag gat gac gac gat aag 3' (SEQ ID NO: 34) was added to cloning primer at the 3' end of the ORF. Another pSRa construct was made that contained amino acids 182 to 454. This truncated form of the STEAP-2 protein will help determine the function of the long extracellular N-Terminal region.

In addition, a pSRa construct can be made to produce a myc/6 HIS fusion protein of the full length STEAP protein.

#### **Example 8: Expression Analysis of STEAP-2 and Other Human STEAP Family Members**

##### **Example 8A: Tissue Specific Expression of STEAP Family Members in Normal Human Tissues**

RT-PCR analysis of STEAP-2 shows expression in all the LAPC prostate cancer xenografts and in normal prostate (FIG. 14, panel A). Analysis of 8 normal human tissues shows prostate-specific expression after 25 cycles of amplification (FIG. 14, panel B). Lower level expression in other tissues was detected only after 30 cycles of amplification. Northern blotting for STEAP-2 shows a pattern of 2 transcripts (approximately 3 and 8 kb in size) expressed only in prostate (and at significantly lower levels in the LAPC xenografts), with no detectable expression in any of the 15 other normal human tissues analyzed (FIG. 15, panel C). Thus, STEAP-2 expression in normal human tissues appears to be highly prostate-specific.

Expression analysis of STEAP family members in normal tissues was performed by Northern blot and/or RT-PCR. All STEAP family members appeared to exhibit

tissue restricted expression patterns. STEAP-3/AI139607 expression is shown in FIG. 12A (Northern) and FIG. 12B (RT-PCR). STEAP-4/R80991 expression is shown in FIG. 13.

5     **Example 8B: Expression of STEAP-2 in Various Cancer Cell Lines**

The RT-PCR results above suggested that the different STEAP family members exhibit different tissue expression patterns. Interestingly, STEAP-2, which appears very prostate-specific, seems to be expressed at lower levels in the LAPC xenografts. This is in contrast to STEAP-1, which is highly expressed in both normal and malignant  
10     prostate tissue.

To better characterize this suggested difference in the STEAP-2 prostate cancer expression profile (relative to STEAP-1), Northern blotting was performed on RNA derived from the LAPC xenografts, as well as several prostate and other cancer cell lines, using a STEAP-2 specific probe (labeled cDNA clone 98P4B6). The results are shown  
15     in FIG. 16 and can be summarized as follows. STEAP-2 is highly expressed in normal prostate and in some of the prostate cancer xenografts and cell lines. More particularly, very strong expression was observed in the LAPC-9 AD xenograft and the LNCaP cells. Significantly attenuated or no expression was observed in the other prostate cancer xenografts and cell lines. Very strong expression was also evident in the Ewing Sarcoma  
20     cell line RD-ES. Unlike STEAP-1, which is highly expressed in cancer cell lines derived from bladder, colon, pancreatic and ovarian tumors, STEAP-2 showed low to non-detectable expression in these same cell lines (compare with FIG. 5). Interestingly, STEAP-2 was also non-detectable in PrEC cells, which are representative of the normal basal cell compartment of the prostate. These results suggests that expression of  
25     STEAP-1 and STEAP-2 are differentially regulated. While STEAP-1 may be a gene that is generally up-regulated in cancer, STEAP-2 may be a gene that is more restricted to normal prostate and prostate cancer.

30     **Example 8C: Analysis of *in Situ* Expression of STEAP-2 RNA in Normal Prostate and Prostate Cancer**

The expression of STEAP-2 RNA was evaluated using *in situ* hybridization with a labeled antisense probe and using a labeled sense probe as control. All four of four

normal prostate tissue specimens examined were positive for STEAP (Figures 22A-B). Likewise, five of five prostate cancer tissue specimens were positive using STEAP RNA *in situ* hybridization (Figures 23A-B). A PIN specimen and a sample of LNCaP cells that were examined were positive as well.

5

**Example 8D: Analysis of STEAP-2 Expression in Cancer Tissues**

The expression of STEAP-2 in various cancer tissues was examined using RT-PCR. The results are shown in Figure 24, in which lane 1 represents a sample from an LAPC4 AD xenograft; lane 2 is LAPC9 AD xenograft; lane 3 is LAPC9 AD<sup>2</sup> xenograft (grown with human bone explant); lane 4 is LAPC9 AD IT (grown intratibially); lane 5 is pooled tissue from colon cancer patients; lane 6 is pooled tissue from lung cancer patients; M represents a marker lane; lane 7 is patient normal prostate tissue; lane 8 is patient prostate cancer tissue; lane 9 is pooled tissue from kidney cancer patients; lane 10 is pooled tissue from bladder cancer patients; lane 11 is HeLa cells; and lane 12 is a water blank. The highest expression is found in the three LAPC9 AD xenografts. High expression is observed in the LAPC4 AD xenograft as well, and in normal prostate and lung cancer. Significant expression was detected also in colon and bladder cancer. Lower expression was detected in prostate and kidney cancer patient samples.

20 **Example 8E: Analysis of STEAP-2 Expression in Normal Tissues**

The expression of STEAP-2 in 76 normal human tissues was examined using a dot blot analysis of RNA samples. As shown in Figure 25, STEAP-2 is expressed at very high levels only in normal prostate.

25 **Example 9: Chromosomal Localization of STEAP Genes**

The chromosomal localization of STEAP-1 was determined using the GeneBridge 4 Human/Hamster radiation hybrid (RH) panel (Walter et al., 1994, Nat. Genetics 7:22) (Research Genetics, Huntsville AL), while STEAP-2 and the STEAP homologues were mapped using the Stanford G3 radiation hybrid panel (Stewart et al., 30 1997, Genome Res. 7:422).

The following PCR primers were used for STEAP-1:

8P1D4.1 5' ACTTTGTTGATGACCAGGATTGGA 3' (SEQ ID NO: 4)

8P1D4.2 5' CAGAACTTCAGCACACACAGGAAC 3' (SEQ ID NO: 5)

5 The resulting STEAP-1 mapping vector for the 93 radiation hybrid panel DNAs  
(2100000201101010001000000101110101221000111001110110101000100010001010010  
21000001111001010000), and the mapping program available at the internet address  
<<http://www-genome.wi.mit.edu/cgi-bin/contig/rhmapper.pl>>, localized the STEAP-1  
gene to chromosome 7p22.3, telomeric to D7S531.

10

The following PCR primers were used for 98P4B6/STEAP-2:

98P4B6.1 5' GACTGAGCTGGAAGTGAATTTGT 3' (SEQ ID NO: 20)

98P4B6.2 5' TTTGAGGAGACTTCATCTCACTGG 3' (SEQ ID NO: 21)

[illegible]

20 The following PCR primers were used for AI139607:

AI139607.1 5' TTAGGACAACCTTGATCACCAGCA 3' (SEQ ID NO: 16)

AI139607.2 5'TGTCCAGTCCAAACTGGGTTATTT3' (SEQ ID NO: 17)

The resulting vector (0000000010000000000000000000100010000020000000100010000000100010001010000010), and the mapping program available at the internet address <<http://www-shgc.stanford.edu/RH/rhserverformnew.html>>, maps AI139607 to chromosome 7q21.

The following PCR primers were used for R80991:

30 R80991.3 5' ACAAGAGCCACCTCTGGGTGAA 3' (SEQ ID NO: 35)

R80991.4 5' AGTTGAGCGAGTTTGCAATGGAC 3' (SEQ ID NO: 36)

The resulting vector (00000000000200001020000000010000000000000000000000010000000001000011100000001001000001), and the mapping program available at the internet address <<http://www-shgc.stanford.edu/RH/rhserverformnew.html>> maps R80991 to chromosome 2q14-q21, near D2S2591.

5 In summary, the above results show that three of the putative human STEAP family members localize to chromosome 7, as is schematically depicted in FIG. 17. In particular, the STEAP-1 gene localizes to the far telomeric region of the short arm of chromosome 7, at 7p22.3, while STEAP-2 and AI139607 localize to the long arm of chromosome 7, at 7q21 (FIG. 17). R80991 maps to chromosome 2q14-q21.

10

#### **Example 10: Identification of Intron-Exon Boundaries of STEAP-1**

Genomic clones for STEAP-1 were identified by searching GenBank for BAC clones containing STEAP-1 sequences, resulting in the identification of accession numbers AC004969 (PAC DJ1121E10) and AC005053 (BAC RG041D11). Using the sequences derived from the PAC and BAC clones for STEAP the intron-exon boundaries were defined (FIG. 18). A total of 4 exons and 3 introns were identified within the coding region of the STEAP gene. Knowledge of the exact exon-intron structure of the STEAP-1 gene may be used for designing primers within intronic sequences which in turn may be used for genomic amplification of exons. Such amplification permits single-stranded conformational polymorphism (SSCP) analysis to search for polymorphisms associated with cancer. Mutant or polymorphic exons may be sequenced and compared to wild type STEAP. Such analysis may be useful to identify patients who are more susceptible to aggressive prostate cancer, as well as other types of cancer, particularly colon, bladder, pancreatic, ovarian, cervical and testicular cancers.

25 Southern blot analysis shows that the STEAP-1 gene exists in several species including mouse (FIG. 19). Therefore, a mouse BAC library (Mouse ES 129-V release I, Genome Systems, FRAC-4431) was screened with the human cDNA for STEAP-1 (clone 10, Example 2). One positive clone, 12P11, was identified and confirmed by southern blotting (FIG. 20). The intron-exon boundary information for human STEAP may be used to identify the mouse STEAP-1 coding sequences.

30 The mouse STEAP-1 genomic clone may be used to study the biological role of STEAP-1 during development and tumorigenesis. Specifically, the mouse genomic

STEAP-1 clone may be inserted into a gene knock-out (K/O) vector for targeted disruption of the gene in mice, using methods generally known in the art. In addition, the role of STEAP in metabolic processes and epithelial cell function may be elucidated. Such K/O mice may be crossed with other prostate cancer mouse models, such as the

5 TRAMP model (Greenberg et al., 1995, PNAS 92:3439), to determine whether STEAP influences the development and progression of more or less aggressive and metastatic prostate cancers.

**Example 11: Predicted HLA-A2 Binding Peptides of STEAP-1 and STEAP-2**

The complete amino acid sequences of the STEAP-1 and STEAP-2 proteins

10 were entered into the HLA Peptide Motif Search algorithm found in the Bioinformatics and Molecular Analysis Section (BIMAS) Web site (<http://bimas.dcrt.nih.gov/>). The HLA Peptide Motif Search algorithm was developed by Dr. Ken Parker based on binding of specific peptide sequences in the groove of HLA Class I molecules and specifically HLA-A2 (Falk et al., 1991, Nature 351: 290-6; Hunt et al., 1992, Science

15 255:1261-3; Parker et al., 1992, J. Immunol. 149:3580-7; Parker et al., 1994, J. Immunol. 152:163-75). This algorithm allows location and ranking of 8-mer, 9-mer, and 10-mer peptides from a complete protein sequence for predicted binding to HLA-A2 as well as other HLA Class I molecules. Most HLA-A2 binding peptides are 9-mers favorably containing a leucine (L) at position 2 and a valine (V) or leucine (L) at position 9.

20 The results of STEAP-1 and STEAP-2 predicted binding peptides are shown in Table 2 below. For both proteins the top 5 ranking candidates are shown along with their location, the amino acid sequence of each specific peptide, and an estimated binding score. The binding score corresponds to the estimated half-time of dissociation of complexes containing the peptide at 37°C at pH 6.5. Peptides with the highest

25 binding score (i.e. 10776.470 for STEAP-1 peptide 165; 1789.612 for STEAP-2 peptide 227) are predicted to be the most tightly bound to HLA Class I on the cell surface and thus represent the best immunogenic targets for T-cell recognition. Actual binding of peptides to HLA-A2 can be evaluated by stabilization of HLA-A2 expression on the antigen-processing defective cell line T2 (Refs. 5,6). Immunogenicity of specific peptides

30 can be evaluated in vitro by stimulation of CD8+ cytotoxic T lymphocytes (CTL) in the presence of dendritic cells (Xue et al., 1997, Prostate 30:73-8; Peshwa et al., 1998, Prostate 36:129-38).



**Table 2: Predicted Binding of STEAP-1 and STEAP-2 Peptide Sequences With Highest Affinity for HLA-A2**

STEAP-1			
Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation)
1	165	GLLSFFFAV	10776.470
2	86	FLYTLLREV	470.951
3	262	LLGTHAL	309.050
4	302	LIFKSILFL	233.719
5	158	MLTRKQFGL	210.633
STEAP-2			
Rank	Start Position	Subsequence Residue Listing	Score (Estimate of Half Time of Disassociation)
1	227	FLYSFVRDV	1789.612
2	402	ALLISTFHV	1492.586
3	307	LLSFFFAMV	853.681
4	306	GLLSFFFAM	769.748
5	100	SLWDLRHLL	726.962

**Example 12: STEAP-2 Induction of Tyrosine Phosphorylation of Cellular Proteins**

Multi-transmembrane proteins have the ability to transmit signal from the membrane and initiate signaling cascades that regulate a variety of downstream events, including gene expression, cellular differentiation, migration and proliferation (Biochim. Biophys. Acta 1997; 1348:56-62, J. Virol. 1995; 69:675-83). In order to determine the involvement of STEAP-1 and STEAP-2 in downstream signaling events, the effect of STEAP-1 and STEAP-2 on tyrosine phosphorylation of PC3 cells was investigated (Figure 26). PC3 cells, stably expressing neo, STEAP-1, or STEAP-2 were grown overnight in 1% fetal bovine serum (FBS) to reduce receptor occupancy and background activity. The cells were then incubated for 5 minutes in the presence of either 1% or 10% FBS, lysed and analyzed by western blotting with anti-phosphotyrosine (4G10 mAb). An overlay using anti-Grb2 Ab was used to show that the gel was equally loaded.

The results (Figure 26) show that expression of STEAP-2 in PC3 cells induces the phosphorylation of several proteins on tyrosine residues, including p150, p120 and p75. In contrast, expression of STEAP-1 induced the de-phosphorylation of p150. Several of the phosphorylated proteins correspond to known signaling proteins, indicating that STEAP-1 and STEAP-2 may be controlling the activation of specific signaling cascades.

### **Example 13: STEAP-1- and STEAP-2- Mediated Activation of MAPK Cascades**

Several multi-transmembrane proteins induce specific biological responses by activating protein kinase cascades, including the MAPK pathways (Curr. Med. Chem. 2000; 7:911-43, Life Sci. 2000; 67:335-6, Biol. Chem. 2000; 275:4660-9). In order to determine whether expression of STEAP-1 and STEAP-2 is sufficient to regulate specific signaling pathways not otherwise active in resting PC3 cells, the effect of these genes on the activation of the p38 MAPK cascade was investigated in the prostate cancer cell line PC3 (Figure 27A-B). Activation of the p38 kinase is dependent on its phosphorylation on tyrosine and serine residues. Phosphorylated p38 can be distinguished from the non-phosphorylated state by a Phospho-p38 mAb. This phospho-specific Ab was used to study the phosphorylation state of p38 in engineered PC3 cell lines.

PC3 cells stably expressing STEAP-1, STEAP-2 or neo were grown overnight in either 1% or 10% FBS. Whole cell lysates were analyzed by western blotting. PC3 cells treated with the known p38 activators, NaSaI or TNF, were used as a positive control. The results show that while expression of the control neo gene has no effect on p38 phosphorylation, expression of STEAP-1 and STEAP-2 in PC3 cells is sufficient to induce the activation of the p38 pathway (Figure 27A). The results were verified using western blotting with an anti-p38 Ab, which shows equal protein loading on the gels (Figure 27B).

In another set of experiments, the sufficiency of expression of STEAP-1 or STEAP-2 in the prostate cancer cell line PC3 to activate the mitogenic MAPK pathway, namely the ERK cascade, was examined (Figure 28A-B). Activation of ERK is dependent on its phosphorylation on tyrosine and serine residues. Phosphorylated ERK can be distinguished from the non-phosphorylated state by a Phospho-ERK mAb. This phospho-specific Ab was used to study the phosphorylation state of ERK in engineered

PC3 cell lines. PC3 cells, expressing an activated form of Ras, were used as a positive control.

The results show that while expression of the control neo gene has no effect on ERK phosphorylation, expression of STEAP-2 in PC3 cells is sufficient to induce at least a 3-fold increase in ERK phosphorylation (Figure 28A). Expression of STEAP-1 also induced some ERK phosphorylation in PC3 cells grown in 1% FBS. These results were verified using anti-ERK western blotting (Figure 28A) and confirm the activation of the ERK pathway by STEAP-1 and STEAP-2.

Since FBS contains several components that may contribute to receptor-mediated ERK activation, we examined the effect of STEAP-1 in low and optimal levels of FBS. PC3 cells expressing neo or STEAP-1 were grown in either 0.1% or 10% FBS overnight. The cells were analyzed by anti-Phospho-ERK western blotting. This experiment shows that STEAP-1 induces the phosphorylation of ERK in 0.1% FBS, and confirms that expression of STEAP-1 is sufficient to induce activation of the ERK signaling cascade in the absence of additional stimuli.

#### **Example 14: Ligand-mediated activation of STEAP-1**

Several factors have been shown to activate multi-transmembrane proteins and mediate downstream signaling events, including ions, leukotrienes, lipophosphatidic acid, etc. (Cell Biochem. Biophys. 1999; 30:213-42). One group of compounds known to activate multi-transmembrane proteins are odorants (Neuron 2000; 25:503-4). In this example, 3 classes of odorants were screened for their ability to induce the activation of tyrosine kinase-mediated signaling in PC3 cells (Figure 29).

PC3 cells, stably expressing neo or STEAP-1 were grown overnight in 0.1% FBS to allow for receptor occupancy. The cells were then treated for 5 min with the indicated concentrations (0.1-10  $\mu$ M) of citralva, ethylvanillin or IBMP. Treatment with 10% FBS was used as a control. Whole cell lysates (20  $\mu$ g) generated from the different treatment conditions were separated by SDS-PAGE and analyzed by anti-phosphotyrosine western blotting.

This experiment shows that, while all 3 classes of odorant induce some tyrosine phosphorylation in PC3-STEAP-1 cells, only IBMP had a measurable effect on the phosphorylation of PC3-neo cells that do not express the STEAP-1 gene. Moreover

both citralva and ethylvanillin induced an increase in the phosphorylation of p136-140 and p200-210 in PC3-STEAP-1 cells in a STEAP-1 specific manner. In addition, citralva induced the *de novo* phosphorylation of a protein at 160-200kDa. The results demonstrate that STEAP-1 is mediating the activation of tyrosine kinase pathways in odorant treated cells.

The finding that citralva induced the tyrosine phosphorylation of proteins in a STEAP-1 mediated manner suggests that STEAP-1 may initiate the activation of one or more signaling cascades. In order to identify potential signaling cascade associated with odorant activation of STEAP-1, the effect of odorants on the activation of the MAPK cascade was studied in PC3 cells (Figure 30). PC3 cells, stably expressing either neo or STEAP-1, were grown overnight in 0.1% FBS. Cells were then treated with citralva for 5 min. Treatment with 10% FBS was used as a control. As observed using phospho-specific mAb (anti-phospho-ERK), citralva activated the ERK pathway in a STEAP-2 specific manner. These results also confirm the findings described in Figures 27-28, that expression of STEAP-1 alone is sufficient to induce the activation of the ERK pathway.

#### **Example 15: Identification of Potential Signal Transduction Pathways**

To confirm that STEAP directly or indirectly activates known signal transduction pathways in cells and to delineate STEAP-mediated downstream events, luciferase (luc) based transcriptional reporter assays are carried out in cells expressing STEAP. These transcriptional reporters contain consensus binding sites for known transcription factors which lie downstream of well characterized signal transduction pathways. The reporters and examples of their associated transcription factors, signal transduction pathways, and activation stimuli are listed below.

25

1. NFkB-luc, NFkB/Rel; Ik-kinase/SAPK; growth/apoptosis/stress
2. SRE-luc, SRF/TCF/ELK1; MAPK/SAPK; growth/differentiation
3. AP-1-luc, FOS/JUN; MAPK/SAPK/PKC; growth/apoptosis/stress
4. ARE-luc, androgen receptor; steroids/MAPK; growth/differentiation/apoptosis
- 30 5. p53-luc, p53; SAPK; growth/differentiation/apoptosis
6. CRE-luc, CREB/ATF2; PKA/p38; growth/apoptosis/stress

STEAP-mediated effects may be assayed in cells showing mRNA expression. Luciferase reporter plasmids may be introduced by lipid mediated transfection (TFX-50, Promega). Luciferase activity, an indicator of relative transcriptional activity, is measured by incubation of cells extracts with luciferin substrate and luminescence of the reaction is  
5 monitored in a luminometer.

**Example 16: In Vitro Assays of STEAP Function**

The expression profile of STEAP in prostate cancer suggests a functional role in tumor initiation, progression and/or maintenance. STEAP function can be assessed in  
10 mammalian cells using in vitro approaches. For mammalian expression, STEAP can be cloned into a number of appropriate vectors, including pcDNA 3.1 myc-His-tag and the retroviral vector pSR $\alpha$ tkneo (Muller et al., 1991, MCB 11:1785). Using such expression vectors, STEAP can be expressed in several cancer cell lines, including for example PC-3, NIH 3T3, LNCaP and 293T. Expression of STEAP can be monitored using anti-  
15 STEAP antibodies.

Mammalian cell lines expressing STEAP can be tested in several in vitro and in vivo assays, including cell proliferation in tissue culture, activation of apoptotic signals, primary and metastatic tumor formation in SCID mice, and in vitro invasion using a membrane invasion culture system (MICS) (Welch et al., Int. J. Cancer 43: 449-457).  
20 STEAP cell phenotype is compared to the phenotype of cells that lack expression of STEAP. In addition, cells treated with and without exogenously added STEAP protein may be analyzed for altered growth parameters.

Cell lines expressing STEAP can also be assayed for alteration of invasive and migratory properties by measuring passage of cells through a matrigel coated porous  
25 membrane chamber (Becton Dickinson). Passage of cells through the membrane to the opposite side is monitored using a fluorescent assay (Becton Dickinson Technical Bulletin #428) using calcein-Am (Molecular Probes) loaded indicator cells. Cell lines analyzed include parental and STEAP overexpressing PC3, 3T3 and LNCaP cells. To assay whether STEAP has chemoattractant properties, parental indicator cells are  
30 monitored for passage through the porous membrane toward a gradient of STEAP conditioned media compared to control media. This assay may also be used to qualify

and quantify specific neutralization of the STEAP induced effect by candidate cancer therapeutic compositions.

In order to establish whether STEAP binds to cellular proteins expressed in prostate cancer cells and other cancer cells or normal cells, two approaches may be taken. In the first approach, in vitro assay for recombinant HIS-tagged STEAP binding to various cell lines are used. In another approach, a recombinant alkaline phosphatase-STEAP fusion protein is generated using the AP-TAG system from GenHunter Corporation (Nashville, TN, cat# Q202), and the AP-TAG fusion used to test STEAP binding to a variety of prostate cancer cell lines as described (Cheng and Flanagan, 1994, Cell 79:157-168). After washing the cells and adding the AP substrate BCIP, which forms an insoluble blue precipitate upon dephosphorylation, STEAP binding is determined by identifying cells staining blue under the light microscope.

Various cancer cell lines can be examined, including without limitation, various prostate cancer cell lines (e.g., LNCaP, PC-3, DU145, TSUPR, LAPC4). Other cell lines such as PREC prostate cell line, 293T, PIN cells, and NIH 3T3, etc. may also be examined. Additionally, the LAPC and other prostate cancer xenografts may be tested. Equilibrium dissociation rate constants may be calculated to evaluate the strength of the binding interaction. In addition, the number of cell surface receptors per cell can be determined. Cell lines or tissues with the highest binding capacity for STEAP would be preferred for cloning a STEAP binding partner.

In another functional assay, NIH-3T3 cells stably expressing STEAP can be analyzed for their ability to form colonies in soft agar. In these experiments, cells used in such procedures (e.g. NIH-3T3 cells), can be transfected to stably express STEAP or neo or activated-Ras (as the test gene, the negative and the positive controls, respectively) in order to assess the transforming capabilities of STEAP. Typically experiments are performed in duplicate and the assays are evaluated approximately 4 weeks after cell plating. Where experimental observations demonstrate that STEAP induces an increase in colony formation relative to a negative control (e.g. neo) such results indicate that STEAP has significant transforming capabilities.

30

**Example 17: In Vivo Assay for STEAP Tumor Growth Promotion**

The effect of the STEAP protein on tumor cell growth may be evaluated in vivo by gene overexpression in tumor-bearing mice. For example, SCID mice can be injected subcutaneously on each flank with  $1 \times 10^6$  of a prostate cell line containing tkNeo empty vector or STEAP. At least two strategies may be used: (1) Constitutive STEAP expression under regulation of an LTR promoter, and (2) Regulated expression under control of an inducible vector system, such as ecdysone, tet, etc. Tumor volume is then monitored at the appearance of palpable tumors and followed over time to determine if STEAP expressing cells grow at a faster rate. Additionally, mice may be implanted with  $1 \times 10^5$  of the same cells orthotopically to determine if STEAP has an effect on local growth in the target tissue (i.e., prostate) or on the ability of the cells to metastasize, specifically to lungs, lymph nodes, liver, bone marrow, etc. The effect of STEAP on bone tumor formation and growth may be assessed by injecting prostate tumor cells intratibially, as described in Example 1.

These assays are also useful to determine the STEAP inhibitory effect of candidate therapeutic compositions, such as for example, STEAP antibodies, STEAP antisense molecules and ribozymes.

**Example 18: Detection of Prostate Cancer Metastases in Mice and Humans Using Anti-STEAP-1 Polyclonal Antibodies***Mice*

STEAP-1 immunohistochemical analysis was performed on 4  $\mu$ m formalin-fixed tissues derived from mice bearing orthotopic LAPC-9 prostate cancer tumors and their derived lung and lymph node metastases. Serial lung sections were tested using anti-STEAP-1 sheep polyclonal antibody and anti-PSA rabbit polyclonal antibody (pAb). Microscopic examination of stained tissue was used to detect LAPC-9 prostate cancer cells, and the results are shown in Figures 31A-F.

The anti-STEAP-1 sheep pAb readily detected human prostate cancer cells in the mouse prostate (Figure 31A), as well as in metastases to lymph nodes (Figure 31B) and lung (Figure 31C-D). Both small (Figure 31C) and large (Figure 31D) micrometastases to the lung were readily detected using the anti-STEAP-1 pAb. To confirm that the lung metastases were of prostate cancer origin, immunohistochemistry on serial lung sections

was performed using the anti-PSA pAb (Figure 31E-F). The strong cell surface staining observed with the anti-STEAP-1 pAb allows for easier detection of micrometastases as compared to the use of the anti-PSA polyclonal antibody.

## 5 *Humans*

The STEAP-1 polyclonal Ab was used in a similar immunohistochemical assay of metastatic prostate cancer specimens from human patients. High expression of STEAP-1 was detected in both the lymph node and bone metastases studied. As shown in Figures 32A (lymph node metastasis) and 32B (bone metastasis), intense pericellular staining is observed in these human patient samples, confirming that STEAP-1 is an excellent marker for the detection of metastatic prostate cancer.

### **Example 19: STEAP-1, STEAP-2 and STEAP-3 Extracellular Loop-Fc Fusion Constructs**

15 To express and secrete the extracellular loops of STEAP-1, STEAP-2, and STEAP-3 into conditioned media of mammalian cell lines, fragments of these genes can be cloned into the pFc vector to generate C-terminal fusions of the human IgG1 Fc region. The pFc vector was generated by cloning human immunoglobulin G1, Fc (bases 74-768 of GenBank accession X70421) preceded by the IgA protease cleavage site (Roche, Cat 1461265) coded by 5' cctcgacctccaacaccgggg 3' (SEQ ID NO: 47) into  
20 pTag-5 (GenHunter Corp. Nashville, TN) using XhoI and ApaI. This construct generates an IgG1 Fc fusion at the C-terminus of STEAP-1, STEAP-2 and STEAP-3 extracellular regions while fusing the IgGK signal sequence to N-terminus.

The resulting recombinant proteins are optimized for secretion into the media of  
25 transfected mammalian cells and can be used to identify proteins such as ligands or receptors that interact with the STEAP-1, STEAP-2, and STEAP-3 protein. These protein fusions can also be used as immunogen to generate antibodies. Protein expression is driven from the CMV promoter. The Zeosin resistance gene allows for selection of mammalian cells expressing the protein and the ampicillin resistance gene  
30 permits selection of the plasmid in *E. coli*.



The following extracellular loops of the STEAP-1, STEAP-2, and STEAP-3 proteins are suitable for cloning into pFc:

- STEAP-1      revihplatshqqyfykipilv (SEQ ID NO: 37)  
 5                rrsyrykllnwayqqvqqnkedawiehdvwrmei (SEQ ID NO: 38)  
                  widikqfvwytpptf (SEQ ID NO: 39)
- STEAP-2      ysfvrdvihpyarnqqsdifykipiei (SEQ ID NO: 40)  
                  rrserylflnmayqqvhanienwneeevwrie (SEQ ID NO: 41)  
 10                krafeeeyryfy (SEQ ID NO: 42)
- STEAP-3      ypyvyekkdntfrmaisipnrifp (SEQ ID NO: 43)  
                  yyvrwrlgnltvtqailkknepfstssawlsdsy (SEQ ID NO: 44)

#### 15      **Example 20: Generation of Polyclonal Antibodies to STEAP-2**

Three immunogens were used to generate antibodies specific to STEAP-2. Two immunogens were peptides encoding amino acids 153-165 (ALQLGPKDASRQV; SEQ ID NO: 45) and amino acids 345-358 (IENSWNEEEVWRIE; SEQ ID NO: 46) of the STEAP-2 protein sequence. The first peptide resides in the N-terminus of STEAP-2, which is intracellular using the membrane topology prediction program SOSUI. The latter peptide resides in the region between transmembrane domains 3 and 4, and this region is predicted to encode the second of 3 extracellular loops. A third immunogen was a glutathione-S-transferase (GST) fusion protein encompassing amino acids 2-204 of the STEAP-2 protein sequence. The recombinant GST-STEAP-2 fusion protein was purified from induced bacteria by glutathione-sepharose affinity chromatography.

In addition to the above antigens, peptides encoding other regions of the STEAP-2 protein, or bacterial and baculovirus virus produced proteins encoding either full length or partial sequences of the STEAP-2 protein, are use to generate a variety of STEAP-2 specific antibodies. These antibodies are directed to regions that may modulate STEAP-2 function.

*Generation of polyclonal antibodies (pAbs)*

To generate pAbs to STEAP-2, the purified GST-fusion protein and the peptides coupled to Keyhole limpet hemacyanin (KLH) were used to immunize individual rabbits as follows. The rabbits were immunized with 200 µg of fusion protein or KLH-peptide antigen mixed in complete Freund's adjuvant. The rabbits were then injected every two weeks with 200 µg of immunogen in incomplete Freund's adjuvant. Test bleeds were taken approximately 7-10 days following each immunization. The titer of peptide 153-165 antiserum was at least 25,000 and of peptide 345-358 antiserum at least 10,000 as determined by ELISA to the respective immunogens. The titer of the GST-fusion serum was at least 300,000. Peptide antiserum is affinity purified by passage of the serum over an affinity column composed of the respective peptide covalently coupled to Affigel matrix (BioRad). Serum raised to the GST-fusion is semi-purified first by removal of GST-reactive antibodies by passage over a GST affinity column. STEAP-2 specific antibody is then isolated by passage over a GST-STEAP-2 affinity column. Alternatively, STEAP-2 specific antisera is isolated by affinity chromatography using a maltose binding protein (MBP)-STEAP-2 fusion protein encoding the same amino acids

**Example 21: Generation of Monoclonal Antibodies to STEAP Proteins**

To generate mAbs to STEAP-1, STEAP-2, STEAP-3, or STEAP-4 proteins, Balb C mice are immunized intraperitoneally with 20-50 µg of KLH-coupled peptide or with bacterial recombinant polypeptides, such as GST-fusion proteins, encoding regions of the respective STEAP member protein sequence mixed in complete Freund's adjuvant. Mice are then subsequently immunized every 2-4 weeks with 20-50 µg of immunogen mixed in Freund's incomplete adjuvant. Alternatively, Ribi adjuvant is used for initial immunizations. Test bleeds are taken 7-10 days following immunization to monitor titer and specificity of the immune response.

Serum from mice immunized with a GST-fusion protein encompassing amino acids 148-251 of STEAP-1 attained a titer of at least  $8 \times 10^6$  as determined by ELISA and specifically recognized STEAP-1 protein by western blot (Figure 33). Once appropriate reactivity and specificity are obtained as determined by ELISA, western blotting, and

flow cytometry analyses, fusion and hybridoma generation is then carried out using established procedures well known in the art (Harlow and Lane, 1988).

Figure 33 is a western blot showing that anti-STEAP-1 murine pAb recognizes STEAP-1 protein in engineered cell lines and endogenous STEAP-1 protein in LNCaP cells. Lysates of LNCaP cells and 293T cells transfected with either pcDNA 3.1 MYC/HIS tagged STEAP-1 or neo empty vector, and RAT1 cells engineered to express STEAP-1 or a neo control gene, were separated by SDS-PAGE and transferred to nitrocellulose. The blot was then subjected to anti-STEAP western analysis using a 1:1000 dilution of serum from mice immunized with a GST-STEAP-1 fusion protein.

Alternative antigens and immunization strategies are also used to generate mAbs with specific reactivity and specificity to various regions of the STEAP proteins. Such antigens may include baculovirus produced recombinant proteins or mammalian expressed and secreted human IgG FC-fusion proteins encoding various regions of each STEAP family member protein sequence, such as predicted extracellular domains. A cell based immunization strategy is also used in which the cDNA of a STEAP family member is overexpressed in cells such as NIH3T3 mouse fibroblasts or 300.19 murine pre-B cells and whole cells or membrane preparations from these cells are used as immunogen. In addition, a DNA-based immunization protocol in which a mammalian expression vector encoding the STEAP-1 or STEAP-2 cDNA such as pcDNA 3.1 is used to immunize mice by direct injection of the plasmid DNA. This protocol is used either alone or in combination with protein and/or cell-based immunization strategies.

---

Throughout this application, various publications are referenced. The disclosures of these publications are hereby incorporated by reference herein in their entireties.

The present invention is not to be limited in scope by the embodiments disclosed herein, which are intended as single illustrations of individual aspects of the invention, and any that are functionally equivalent are within the scope of the invention. Various modifications to the models and methods of the invention, in addition to those described herein, will become apparent to those skilled in the art from the foregoing description and teachings, and are similarly intended to fall within the scope of the invention. Such modifications or other embodiments can be practiced without departing from the true scope and spirit of the invention.

Applicant's or agent's file reference number	29.17WO11	International application No.
--	-----------	-------------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>75</u> , line <u>4-6</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 26 August 1998 (26.08.98)	Accession Number 98849
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer: <b>MISTY WALKER</b> <b>INTERNATIONAL DIVISION</b> <b>703-305-3682</b>	Authorized officer

Applicant's or agent's file reference number	29.17WO11	International application No.
--	-----------	-------------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the microorganism referred to in the description on page <u>84</u> , line <u>1-3</u> .	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 02 July 1999 (02.07.99)	Accession Number PTA-311
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application  Authorized officer: <b>MISTY WALKER</b> <b>INTERNATIONAL DIVISION</b> 703-305-3682	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on:  Authorized officer:
---	--

Applicant's or agent's file reference number	129.17WO11	International application No.
--	------------	-------------------------------

## INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>84</u> , line <u>27-29</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 08 December 1999 (08.12.99)	Accession Number PTA-1033
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer <b>MISTY WALKER</b> <b>INTERNATIONAL DIVISION</b> <b>703-305-2682</b>	Authorized officer

CLAIMS:

1. A STEAP-2 protein having an amino acid sequence shown in FIG. 9A-D (SEQ ID NO: 8).
- 5 2. A STEAP-2 polypeptide that includes at least 10 contiguous amino acids of an amino acid sequence selected from the group consisting of amino acids 1-245, 2-204, 100-108, 121-454, 153-165, 182-454, 183-387, 227-235, 276-453, 306-314, 307-315, 345-358, 402-410, and 419-454 of the amino acid sequence shown in FIG. 9A-D (SEQ ID NO: 8).
- 10 3. A polypeptide comprising an amino acid sequence which is at least 90% identical to the amino acid sequence shown in FIG. 9A-D (SEQ ID NO: 8) over its entire length.
4. A polynucleotide selected from the group consisting of (a) a polynucleotide having the sequence as shown in FIG. 9A-D (SEQ ID NO: 7), wherein T can  
15 also be U; (b) a polynucleotide encoding a STEAP-2 polypeptide whose sequence is encoded by the cDNA contained in plasmid 98P4B6-GTD3 as deposited with American Type Culture Collection as Accession No. PTA-311; (c) a polynucleotide encoding the STEAP-2 protein of claim 1; and (d) a polynucleotide which is fully complementary to a polynucleotide of (a)-(c).
- 20 5. A polynucleotide that encodes the fragment of claim 2.
6. A recombinant expression vector which contains a polynucleotide according to claim 4 or 5.
7. A host cell which contains an expression vector according to claim 6.
8. A polynucleotide according to claim 4 or 5 which is labeled with a detectable  
25 marker.

9. A process for producing a STEAP-2 protein comprising culturing a host cell of claim 7 under conditions sufficient for the production of the polypeptide and recovering the STEAP-2 protein from the culture.
10. An antibody or fragment thereof which specifically binds to an epitope within amino acids 1-245, 2-204, 100-108, 121-454, 153-165, 182-454, 183-387, 227-235, 276-453, 306-314, 307-315, 345-358, 402-410, or 419-454 of the STEAP-2 protein of claim 1.
11. The antibody or fragment thereof of claim 10 which is monoclonal.
12. A recombinant protein comprising the antigen binding region of a monoclonal antibody of claim 11.
13. The antibody or fragment thereof of claim 10 or 11, or the recombinant protein of claim 12, which is labeled with a detectable marker.
14. The antibody or fragment thereof or recombinant protein of claim 13, wherein the detectable marker comprises a radioisotope, metal chelator, enzyme or fluorescent, bioluminescent or chemiluminescent compound.
15. The antibody fragment of claim 13, which is an Fab, F(ab')<sub>2</sub>, Fv or Sfv fragment.
16. The antibody or fragment thereof of any one of claims 10, 11, 13 or 14, which is a human antibody.
17. The antibody of any one of claims 10, 11, 13 or 14, which comprises murine antigen binding region residues and human antibody residues.
18. A transgenic animal producing a monoclonal antibody of claim 11.
19. A hybridoma producing a monoclonal antibody of claim 11.
20. A single chain monoclonal antibody that comprises the variable domains of the heavy and light chains of a monoclonal antibody of claim 11.



21. A vector comprising a polynucleotide encoding a single chain monoclonal antibody of claim 20.
22. The monoclonal antibody or fragment thereof of claim 11 which is conjugated to a toxin, a radioisotope or a therapeutic agent.
- 5 23. The monoclonal antibody or fragment thereof of claim 22, wherein the toxin comprises ricin, ricin A-chain, doxorubicin, daunorubicin, a maytansinoid, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, Pseudomonas exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, 10 alpha-sarcin, gelonin, mitogellin, retstrictocin, phenomycin, enomycin, curicin, crotin, calicheamicin, sapaonaria officinalis inhibitor, or a glucocorticoid.
24. The monoclonal antibody or fragment thereof of claim 22, wherein the radioisotope comprises  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , or  $^{186}\text{Re}$ .
- 15 25. An assay for detecting the presence of a STEAP-2 protein in a biological sample comprising contacting the sample with an antibody of claim 13, and detecting the binding of STEAP-2 protein in the sample thereto.
26. An assay for detecting the presence of a STEAP-2 polynucleotide in a biological sample, comprising:
- 20 (a) contacting the sample with a polynucleotide probe which specifically hybridizes to a polynucleotide of claim 4; and
- (b) detecting the presence of a hybridization complex formed by the hybridization of the probe with STEAP-2 polynucleotide in the sample, wherein the presence of the hybridization complex indicates the presence of STEAP-2 polynucleotide within the sample.
- 25 27. An assay for detecting the presence of STEAP-2 mRNA in a biological sample comprising:

- (a) producing cDNA from the sample by reverse transcription using at least one primer;
- (b) amplifying the cDNA so produced using STEAP-2 polynucleotides as sense and antisense primers to amplify STEAP-2 cDNAs therein;
- 5 (c) detecting the presence of the amplified STEAP-2 cDNA,

wherein the STEAP-2 polynucleotides used as the sense and antisense primers are capable of amplifying the polynucleotide shown in FIG. 9A-D (SEQ ID NO: 7).

28. A pharmaceutical composition comprising an antibody or fragment thereof  
10 according to claim 10, 11, 15, 16 or 17, a STEAP-2 protein according to claim 1 or 2, or a STEAP-2 polynucleotide according to claim 4 or 5, and a physiologically acceptable carrier.
29. A vaccine composition comprising a STEAP-2 protein according to claim 1 or 2,  
15 or a STEAP-2 polynucleotide according to claim 4 or 5, and a physiologically acceptable carrier.
30. A method of inhibiting the development or progression of a cancer expressing STEAP-2 in a patient, comprising administering to the patient an effective amount of the vaccine composition of claim 29.
31. A method of inhibiting the growth of cells expressing STEAP-2 in a patient,  
20 comprising administering to the patient an effective amount of the vaccine composition of claim 29.
32. A method of killing cells that express STEAP-2 in a patient, comprising administering to the patient an effective amount of the vaccine composition of claim 29.
- 25 33. A method of treating a patient with a cancer that expresses STEAP-2, which comprises administering to said patient a vector encoding a single chain monoclonal antibody that comprises the variable domains of the heavy and light

chains of a monoclonal antibody that specifically binds to a STEAP-2 protein of claim 2, such that the vector delivers the single chain monoclonal antibody coding sequence to the cancer cells and the encoded single chain antibody is expressed intracellularly therein.

- 5     34.     A method of inhibiting growth of cells that express STEAP-2, which comprises administering to said patient a vector encoding a single chain monoclonal antibody that comprises the variable domains of the heavy and light chains of a monoclonal antibody that specifically binds to a STEAP-2 protein of claim 2, such that the vector delivers the single chain monoclonal antibody coding  
10     sequence to the cancer cells and the encoded single chain antibody is expressed intracellularly therein.
- 15     35.     A method of killing cells that express STEAP-2, which comprises administering to said patient a vector encoding a single chain monoclonal antibody that comprises the variable domains of the heavy and light chains of a monoclonal  
antibody that specifically binds to a STEAP-2 protein of claim 2, such that the vector delivers the single chain monoclonal antibody coding sequence to the cancer cells and the encoded single chain antibody is expressed intracellularly  
therein.
- 20     36.     A STEAP-3 protein having an amino acid sequence shown in FIG. 10A-E (SEQ ID NO: 10).
37.     A polypeptide of at least 10 contiguous amino acids of the protein of claim 36.
38.     A polypeptide comprising an amino acid sequence which is at least 90% identical to the amino acid sequence shown in FIG. 10A-E (SEQ ID NO: 10) over its entire length.
- 25     39.     A polynucleotide selected from the group consisting of (a) a polynucleotide having the sequence as shown in FIG. 10A-E (SEQ ID NO: 9), wherein T can also be U; (b) a polynucleotide encoding the STEAP-3 protein of claim 1; and (c) a polynucleotide which is fully complementary to a polynucleotide of (a) or (b).

40. An antibody or fragment thereof which specifically binds to the STEAP-3 protein of claim 36.
41. The antibody or fragment thereof of claim 40 which is monoclonal.
42. A recombinant protein comprising the antigen binding region of a monoclonal antibody of claim 41.
43. The antibody or fragment thereof of claim 40 or 41 which is labeled with a detectable marker.
44. The antibody or fragment thereof or recombinant protein of claim 43, wherein the detectable marker comprises a radioisotope, metal chelator, enzyme or fluorescent, bioluminescent or chemiluminescent compound.
45. The monoclonal antibody or fragment thereof of claim 41 which is conjugated to a toxin, a radioisotope or a therapeutic agent.
46. The monoclonal antibody or fragment thereof of claim 41, wherein the toxin comprises ricin, ricin A-chain, doxorubicin, daunorubicin, a maytansinoid, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin, diphtheria toxin, Pseudomonas exotoxin (PE) A, PE40, abrin, abrin A chain, modeccin A chain, alpha-sarcin, gelonin, mitogellin, retstrictocin, phenomycin, enomycin, curicin, crotin, calicheamicin, sapaonaria officinalis inhibitor, or a glucocorticoid.
47. The monoclonal antibody or fragment thereof of claim 41, wherein the radioisotope comprises  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , or  $^{186}\text{Re}$ .
48. The antibody fragment of claim 41, which is an Fab, F(ab')<sub>2</sub>, Fv or Sfv fragment.
49. The antibody or fragment thereof of any one of claims 40, 41, and 43-48, which is a human antibody.
50. The antibody or fragment thereof of any one of claims 40, 41, and 43-48, which comprises murine antigen binding region residues and human antibody residues.

51. A transgenic animal producing a monoclonal antibody of claim 41.
52. A hybridoma producing a monoclonal antibody of claim 41.
53. A single chain monoclonal antibody that comprises the variable domains of the heavy and light chains of a monoclonal antibody of claim 41.
- 5 54. A vector comprising a polynucleotide encoding a single chain monoclonal antibody of claim 53.
55. An assay for detecting the presence of a STEAP-3 protein in a biological sample comprising contacting the sample with an antibody of claim 43 or 44, and detecting the binding of STEAP-3 protein in the sample thereto.
- 10 56. An assay for detecting the presence of a STEAP-3 polynucleotide in a biological sample, comprising
- (a) contacting the sample with a polynucleotide probe which specifically hybridizes to a polynucleotide of claim 39 or its complement; and
- (b) detecting the presence of a hybridization complex formed by the
- 15 hybridization of the probe with STEAP-3 polynucleotide in the sample,
- wherein the presence of the hybridization complex indicates the presence of STEAP-3 polynucleotide within the sample.
57. An assay for detecting the presence of STEAP-3 mRNA in a biological sample comprising:
- 20 (a) producing cDNA from the sample by reverse transcription using at least one primer;
- (b) amplifying the cDNA so produced using STEAP-3 polynucleotides as sense and antisense primers to amplify STEAP-3 cDNAs therein;
- (c) detecting the presence of the amplified STEAP-3 cDNA,

wherein the STEAP-3 polynucleotides used as the sense and antisense primers are capable of amplifying the polynucleotide of FIG. 10A-E (SEQ ID NO: 9).

58. A pharmaceutical composition comprising an antibody or fragment thereof or recombinant protein of any one of claims 40-50, a STEAP-3 protein according to claim 36, or a STEAP-3 polynucleotide according to claim 39, and a physiologically acceptable carrier.

1/51

## FIG. 1A

```

5'  GAG  ACT  CAC  GGT  CAA  GCT  AAG  GCG  AAG  AGT  GGG  TGG  CTG  AAG  CCA  TAC  TAT  TTT
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---

      11      20      29      38      47      56
    ATA  GAA  TTA  ATG  GAA  AGC  AGA  AAA  GAC  ATC  ACA  AAC  CAA  GAA  GAA  CTT  TGG  AAA
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
              M   E   S   R   K   D   I   T   N   Q   E   E   L   W   K

      119      128      137      146      155      164
    ATG  AAG  CCT  AGG  AGA  AAT  TTA  GAA  GAA  GAC  GAT  TAT  TTG  CAT  AAG  GAC  ACG  GGA
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    M   K   P   R   R   N   L   E   E   D   D   Y   L   H   K   D   T   G

      173      182      191      200      209      218
    GAG  ACC  AGC  ATG  CTA  AAA  AGA  CCT  GTG  CTT  TTG  CAT  TTG  CAC  CAA  ACA  GCC  CAT
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    E   T   S   M   L   K   R   P   V   L   L   H   L   H   Q   T   A   H

      227      236      245      254      263      272
    GCT  GAT  GAA  TTT  GAC  TGC  CCT  TCA  GAA  CTT  CAG  CAC  ACA  CAG  GAA  CTC  TTT  CCA
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    A   D   E   F   D   C   P   S   E   L   Q   H   T   Q   E   L   F   P

      281      290      299      308      317      326
    CAG  TGG  CAC  TTG  CCA  ATT  AAA  ATA  GCT  GCT  ATT  ATA  GCA  TCT  CTG  ACT  TTT  CTT
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Q   W   H   L   P   I   K   I   A   A   I   I   A   S   L   T   F   L

      335      344      353      362      371      380
    TAC  ACT  CTT  CTG  AGG  GAA  GTA  ATT  CAC  CCT  TTA  GCA  ACT  TCC  CAT  CAA  CAA  TAT
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    Y   T   L   L   R   E   V   I   H   P   L   A   T   S   H   Q   Q   Y

      389      398      407      416      425      434
    TTT  TAT  AAA  ATT  CCA  ATC  CTG  GTC  ATC  AAC  AAA  GTC  TTG  CCA  ATG  GTT  TCC  ATC
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    F   Y   K   I   P   I   L   V   I   N   K   V   L   P   M   V   S   I

      443      452      461      470      479      488
    ACT  CTC  TTG  GCA  TTG  GTT  TAC  CTG  CCA  GGT  GTG  ATA  GCA  GCA  ATT  GTC  CAA  CTT
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    T   L   L   A   L   V   Y   L   P   G   V   I   A   A   I   V   Q   L

      497      506      515      524      533      542
    CAT  AAT  GGA  ACC  AAG  TAT  AAG  AAG  TTT  CCA  CAT  TGG  TTG  GAT  AAG  TGG  ATG  TTA
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    H   N   G   T   K   Y   K   K   F   P   H   W   L   D   K   W   M   L

      551      560      569      578      587      596
    ACA  AGA  AAG  CAG  TTT  GGG  CTT  CTC  AGT  TTC  TTT  TTT  GCT  GTA  CTG  CAT  GCA  ATT
    ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---  ---
    T   R   K   Q   F   G   L   L   S   F   F   F   A   V   L   H   A   I

```

2/51

## FIG. 1B

TAT	AGT	605	CTG	TCT	TAC	614	CCA	ATG	AGG	623	CGA	TCC	TAC	632	AGA	TAC	AAG	641	TTG	CTA	AAC	650	TGG
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<u>Y</u>	<u>S</u>	<u>L</u>	<u>S</u>	<u>Y</u>	<u>P</u>	<u>M</u>		R		R	S	Y	R	Y	K	L	L	N	W				
GCA	TAT	659	CAA	CAG	GTC	668	CAA	CAA	AAT	677	AAA	GAA	GAT	686	GCC	TGG	ATT	695	GAG	CAT	GAT	704	GTT
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
A	Y	Q	Q	V	Q	Q	N	K	E	D	A	W	I	E	H	D	V						
TGG	AGA	713	ATG	GAG	ATT	722	TAT	GTG	TCT	731	CTG	GGA	ATT	740	GTG	GGA	TTG	749	GCA	ATA	CTG	758	GCT
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
W	R	M	E	I	<u>Y</u>	<u>V</u>	<u>S</u>	<u>L</u>	<u>G</u>	<u>I</u>	<u>V</u>	<u>G</u>	<u>L</u>	<u>A</u>	<u>I</u>	<u>L</u>	<u>A</u>						
CTG	TTG	767	GCT	GTG	ACA	776	TCT	ATT	CCA	785	TCT	GTG	AGT	794	GAC	TCT	TTG	803	ACA	TGG	AGA	812	GAA
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<u>L</u>	<u>L</u>	<u>A</u>	<u>V</u>	<u>T</u>	<u>S</u>	<u>I</u>	<u>P</u>	<u>S</u>	<u>V</u>	S	D	S	L	T	W	R	E						
TTT	CAC	821	TAT	ATT	CAG	830	AGC	AAG	CTA	839	GGA	ATT	GTT	848	TCC	CTT	CTA	857	CTG	GGC	ACA	866	ATA
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
F	H	Y	I	<u>Q</u>	<u>S</u>	<u>K</u>	<u>L</u>	<u>G</u>	<u>I</u>	<u>V</u>	<u>S</u>	<u>L</u>	<u>L</u>	<u>L</u>	<u>G</u>	<u>T</u>	<u>I</u>						
CAC	GCA	875	TTG	ATT	TTT	884	GCC	TGG	AAT	893	AAG	TGG	ATA	902	GAT	ATA	AAA	911	CAA	TTT	GTA	920	TGG
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<u>H</u>	<u>A</u>	<u>L</u>	<u>I</u>	<u>F</u>	<u>A</u>	<u>W</u>	<u>N</u>	<u>K</u>	W	I	D	I	K	Q	F	V	W						
TAT	ACA	929	CCT	CCA	ACT	938	TTT	ATG	ATA	947	GCT	GTT	TTC	956	CTT	CCA	ATT	965	GTT	GTC	CTG	974	ATA
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Y	T	P	P	T	F	<u>M</u>	<u>I</u>	<u>A</u>	<u>V</u>	<u>F</u>	<u>L</u>	<u>P</u>	<u>I</u>	<u>V</u>	<u>V</u>	<u>L</u>	<u>I</u>						
TTT	AAA	983	AGC	ATA	CTA	992	TTC	CTG	CCA	1001	TGC	TTG	AGG	1010	AAG	AAG	ATA	1019	CTG	AAG	ATT	1028	AGA
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<u>F</u>	<u>K</u>	<u>S</u>	<u>I</u>	<u>L</u>	<u>F</u>	<u>L</u>	<u>P</u>	<u>C</u>	<u>L</u>	R	K	K	I	L	K	I	R						
CAT	GGT	1037	TGG	GAA	GAC	1046	GTC	ACC	AAA	1055	ATT	AAC	AAA	1064	ACT	GAG	ATA	1073	TGT	TCC	CAG	1082	TTG
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
H	G	W	E	D	V	T	K	I	N	K	T	E	I	C	S	Q	L						
TAG	AAT	1091	TAC	TGT	TTA	1100	CAC	ACA	TTT	1109	TTG	TTC	AAT	1118	ATT	GAT	ATA	1127	TTT	TAT	CAC	1136	CAA
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
*	N	Y	C	L	H	T	F	L	F	N	I	D	I	F	Y	H	Q						
CAT	TTC	1145	AAG	TTT	GTA	1154	TTT	GTT	AAT	1163	AAA	ATG	ATT	1172	ATT	CAA	GGA	1181	AAA	AAA	AAA	1190	AAA
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
H	F	K	F	V	F	V	N	K	M	I	I	Q	G	K	K	K	K						


AAA AA 3'

---

K



FIG. 1C



4/51

## FIG. 1D

```
5'   GGC GGA GGC GGA GGC GGA GGG CGA GGG GCG GGG AGC GCC GCC TGG AGC GCG  
    GCA GGT CAT ATT GAA CAT TCC AGA TAC CTA TCA TTA CTC GAT GCT GTT GAT  
AAC AGC AAG   3'
```

FIG. 2A





FIG. 2B

FIG. 2C

FIG. 3A





FIG. 3B



## FIG. 4A

[illegible]

9/51

## FIG. 4B

CTCCATCTGAATTAGTCCAGTGTGGCTTAGAGTTGGTACAACATTCTCACAGAATTCCTAATTTGTAGGTTTCAG  
CCTGATAACCACTGGAGTTCCTTTGGTCCTCATTAAATAGCTTTCTTCACACATTGCTCTGCCTGTTACACATATGA  
TGAACACTGCTTTTTAGACTTCATTAGGAATTTAGGACTGCATCTTGACAACCTGAGCCTATTCTACTATATGTACA  
ATACCTAGCCCATAATAGGTATACAATACACATTTGGTAAAACATAATTTCAACCAATGACATGTATTTTCAACT  
AGTAACCTAGAAATGTTTCACTTAAAATCTGAGAACTGGTTACACTACAAGTTACCTTGGAGATTCATATATGAAA  
ACGCAAACTTAGCTATTTGATTGTATTCACTGGGACTTAAGAATGCGCCTGAATAATTGTGAGTTCGATTGTCTCT  
GGCAGGCTAATGACCATTTCCAGTAAAGTGAATAGAGGTCAGAAGTCGTATAAAAAGAGGTGTTGTCAGAACACCGT  
TGAGATTACATAGGTGAACAACTATTTTTAAGCAACTTTATTTGTGTAGTGACAAAGCATCCCAATGCAGGCTGAA  
ATGTTTCATCACATCTCTGGATCTCTCTATTTTGTGCAGACATTGAAAAAATGTTTCATATTATTTCCATGTTATC  
AGAATATTTGATTTTTTAAAAACATAGGCCAAGTTCATTCACCTCATTATTCAATTTATCAAAATCAGAGTGAATCA  
CATTAAGTCGCCTTCACAACTGATAAAGATCACTGAAGTCAAATGATTTTTGCTATAATCTTCAATCTACCTATAT  
TTAATTGAGAATCTAAAATGTACAAATCATTGTGTTGATTCTGCAGTGATCCTGCTATAAGTAAGACTCAGTCCCT  
GATTTTAGGTATCCTGTGAAAAGCAGAATTAAGACAAATACACAAGAGACAAAGCACAAAAAATAAATATCATAAG  
GGGATGAACAAAATGGTGGAGAAAGAGTAGACAAAGTTTTTGATCACCTGCCTTCAAAGAAAGGCTGTGAATTTG  
TTCACCTAGACAGCTTGGAGACAAGAAATTACCCAAAAGTAAGGTGAGGAGGATAGGCAAAAAGAGCAGAAAGATG  
TGAATGGACATTGTTGAGAAATGTGATAGGAAAACAATCATAGATAAAGGATTTCCAAGCAACAGAGCATATCCAG  
ATGAGGTAGGATGGGATAAACTCTTATTGAACCAATCTTCACCAATTTTGTCTTTCTTTTGCAGAGCAAGCTAGGA  
ATTGTTTCCCTTCTACTGGGCACAATACACGCATTGATTTTTGCCTGGAATAAGTGGATAGATATAAAACAATTTG  
TATGGTATACACCTCCAACTTTTATGATAGCTGTTTTCTTCCAATTGTTGTCCTGATATTTAAAGCATACTATT  
CCTGCCATGCTTGAGGAAGAAGATACTGAAGATTAGACATGGTTGGGAAGACGTCACCAAAATTAACAAAACCTGAG  
ATATGTTCCCAGTTGTAGAAATTACTGTTTACACACATTTTGTTCATATTGATATATTTTATCACCACATTTCA  
AGTTTGTATTTGTTAATAAAATGATTATTCAAGGAAAAAAAAAAAAAAAAAAAAA

FIG. 5A




FIG. 5B






FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D




FIG. 8E




FIG. 8F




FIG. 8G





FIG. 8H



14/51

## FIG. 9A

5'		10		19		28		37		46		55						
	GGA	CGC	GTG	GGC	GGA	CGC	GTG	GGT	TCC	TCG	GGC	CCT	CGG	CGC	CAC	AAG	CTG	TCC
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
		64		73		82		91		100		109						
	GGG	CAC	GCA	GCC	CCT	AGC	GGC	GCG	TCG	CTG	CCA	AGC	CGG	CCT	CCG	CGC	GCC	TCC
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
		118		127		136		145		154		163						
	CTC	CTT	CCT	TCT	CCC	CTG	GCT	GTT	CGC	GAT	CCA	GCT	TGG	GTA	GGC	GGG	GAA	GCA
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
		172		181		190		199		208		217						
	GCT	GGA	GTG	CGA	CCG	CCA	CGG	CAG	CCA	CCC	TGC	AAC	CGC	CAG	TCG	GAG	GTG	CAG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
		226		235		244		253		262		271						
	TCC	GTA	GGC	CCT	GGC	CCC	CGG	GTG	GGC	CCT	TGG	GGA	GTC	GGC	GCC	GCT	CCC	GAG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
		280		289		298		307		316		325						
	GAG	CTG	CAA	GGC	TCG	CCC	CTG	CCC	GGC	GTG	GAG	GGC	GCG	GGG	GGC	GCG	GAG	GAT
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
		334		343		352		361		370		379						
	ATT	CTT	GGT	GAT	CTT	GGA	AGT	GTC	CGT	<b>ATC</b>	<b>ATG</b>	<b>GAA</b>	TCA	ATC	TCT	ATG	ATG	GGA
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
										<b>M</b>	<b>E</b>	<b>S</b>	<b>I</b>	<b>S</b>	<b>M</b>	<b>M</b>	<b>G</b>	
		388		397		406		415		424		433						
	AGC	CCT	AAG	AGC	CTT	AGT	GAA	ACT	TGT	TTA	CCT	AAT	GGC	ATA	AAT	GGT	ATC	AAA
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	<b>S</b>	<b>P</b>	<b>K</b>	<b>S</b>	<b>L</b>	<b>S</b>	<b>E</b>	<b>T</b>	<b>C</b>	<b>L</b>	<b>P</b>	<b>N</b>	<b>G</b>	<b>I</b>	<b>N</b>	<b>G</b>	<b>I</b>	<b>K</b>
		442		451		460		469		478		487						
	GAT	GCA	AGG	AAG	GTC	ACT	GTA	GGT	GTG	ATT	GGA	AGT	GGA	GAT	TTT	GCC	AAA	TCC
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	<b>D</b>	<b>A</b>	<b>R</b>	<b>K</b>	<b>V</b>	<b>T</b>	<b>V</b>	<b>G</b>	<b>V</b>	<b>I</b>	<b>G</b>	<b>S</b>	<b>G</b>	<b>D</b>	<b>F</b>	<b>A</b>	<b>K</b>	<b>S</b>
		496		505		514		523		532		541						
	TTG	ACC	ATT	CGA	CTT	ATT	AGA	TGC	GGC	TAT	CAT	GTG	GTC	ATA	GGA	AGT	AGA	AAT
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	<b>L</b>	<b>T</b>	<b>I</b>	<b>R</b>	<b>L</b>	<b>I</b>	<b>R</b>	<b>C</b>	<b>G</b>	<b>Y</b>	<b>H</b>	<b>V</b>	<b>V</b>	<b>I</b>	<b>G</b>	<b>S</b>	<b>R</b>	<b>N</b>
		550		559		568		577		586		595						
	CCT	AAG	TTT	GCT	TCT	GAA	TTT	TTT	CCT	CAT	GTG	GTA	GAT	GTC	ACT	CAT	CAT	GAA
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	<b>P</b>	<b>K</b>	<b>F</b>	<b>A</b>	<b>S</b>	<b>E</b>	<b>F</b>	<b>F</b>	<b>P</b>	<b>H</b>	<b>V</b>	<b>V</b>	<b>D</b>	<b>V</b>	<b>T</b>	<b>H</b>	<b>H</b>	<b>E</b>
		604		613		622		631		640		649						
	GAT	GCT	CTC	ACA	AAA	ACA	AAT	ATA	ATA	TTT	GTT	GCT	ATA	CAC	AGA	GAA	CAT	TAT
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	<b>D</b>	<b>A</b>	<b>L</b>	<b>T</b>	<b>K</b>	<b>T</b>	<b>N</b>	<b>I</b>	<b>I</b>	<b>F</b>	<b>V</b>	<b>A</b>	<b>I</b>	<b>H</b>	<b>R</b>	<b>E</b>	<b>H</b>	<b>Y</b>
		658		667		676		685		694		703						
	ACC	TCC	CTG	TGG	GAC	CTG	AGA	CAT	CTG	CTT	GTG	GGT	AAA	ATC	CTG	ATT	GAT	GTG
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	<b>T</b>	<b>S</b>	<b>L</b>	<b>W</b>	<b>D</b>	<b>L</b>	<b>R</b>	<b>H</b>	<b>L</b>	<b>L</b>	<b>V</b>	<b>G</b>	<b>K</b>	<b>I</b>	<b>L</b>	<b>I</b>	<b>D</b>	<b>V</b>
		712		721		730		739		748		757						
	AGC	AAT	AAC	ATG	AGG	ATA	AAC	CAG	TAC	CCA	GAA	TCC	AAT	GCT	GAA	TAT	TTG	GCT
	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
	<b>S</b>	<b>N</b>	<b>N</b>	<b>M</b>	<b>R</b>	<b>I</b>	<b>N</b>	<b>Q</b>	<b>Y</b>	<b>P</b>	<b>E</b>	<b>S</b>	<b>N</b>	<b>A</b>	<b>E</b>	<b>Y</b>	<b>L</b>	<b>A</b>

15/51

## FIG. 9B

766	775	784	793	802	811
TCA TTA TTC CCA GAT TCT TTG ATT GTC AAA GGA TTT AAT GTT GTC TCA GCT TGG					
S L F P D S L I V K G F N V V S A W					
820	829	838	847	856	865
GCA CTT CAG TTA GGA CCT AAG GAT GCC AGC CGG CAG GTT TAT ATA TGC AGC AAC					
A L Q L G P K D A S R Q V Y I C S N					
874	883	892	901	910	919
AAT ATT CAA GCG CGA CAA CAG GTT ATT GAA CTT GCC CGC CAG TTG AAT TTC ATT					
N I Q A R Q Q V I E L A R Q L N F I					
928	937	946	955	964	973
CCC ATT GAC TTG GGA TCC TTA TCA TCA GCC AGA GAG ATT GAA AAT TTA CCC CTA					
P I D L G S L S S A R E I E N L P <u>L</u>					
982	991	1000	1009	1018	1027
CGA CTC TTT ACT CTC TGG AGA GGG CCA GTG GTG GTA GCT ATA AGC TTG GCC ACA					
<u>R L F T L W R G P V V V A I S L A T</u>					
1036	1045	1054	1063	1072	1081
TTT TTT TTC CTT TAT TCC TTT GTC AGA GAT GTG ATT CAT CCA TAT GCT AGA AAC					
<u>F F F L</u> Y S F V R D V I H P Y A R N					
1090	1099	1108	1117	1126	1135
CAA CAG AGT GAC TTT TAC AAA ATT CCT ATA GAG ATT GTG AAT AAA ACC TTA CCT					
Q Q S D F Y K I P I E I <u>V N K T L P</u>					
1144	1153	1162	1171	1180	1189
ATA GTT GCC ATT ACT TTG CTC TCC CTA GTA TAC CTT GCA GGT CTT CTG GCA GCT					
<u>I V A I T L L S L V Y L A G L L A A</u>					
1198	1207	1216	1225	1234	1243
GCT TAT CAA CTT TAT TAC GGC ACC AAG TAT AGG AGA TTT CCA CCT TGG TTG GAA					
A Y Q L Y Y G T K Y R R F P P W L E					
1252	1261	1270	1279	1288	1297
ACC TGG TTA CAG TGT AGA AAA CAG CTT GGA TTA CTA AGT TTT TTC TTC GCT ATG					
T W L Q C R K <u>Q L G L L S F F F A M</u>					
1306	1315	1324	1333	1342	1351
GTC CAT GTT GCC TAC AGC CTC TGC TTA CCG ATG AGA AGG TCA GAG AGA TAT TTG					
<u>V H V A Y S L C L P M</u> R R S E R Y L					
1360	1369	1378	1387	1396	1405
TTT CTC AAC ATG GCT TAT CAG CAG GTT CAT GCA AAT ATT GAA AAC TCT TGG AAT					
F L N M A Y Q Q V H A N I E N S W N					

16/51

## FIG. 9C

1414	1423	1432	1441	1450	1459
GAG GAA GAA GTT TGG AGA ATT GAA ATG TAT ATC TCC TTT GGC ATA ATG AGC CTT					
-----					
E E E V W R I E <u>M Y I S F G I M S L</u>					
1468	1477	1486	1495	1504	1513
GGC TTA CTT TCC CTC CTG GCA GTC ACT TCT ATC CCT TCA GTG AGC AAT GCT TTA					
-----					
<u>G L L S L L A V T S I P S</u> V S N A L					
1522	1531	1540	1549	1558	1567
AAC TGG AGA GAA TTC AGT TTT ATT CAG TCT ACA CTT GGA TAT GTC GCT CTG CTC					
-----					
N W R E F S <u>F I Q S T L G Y V A L L</u>					
1576	1585	1594	1603	1612	1621
ATA AGT ACT TTC CAT GTT TTA ATT TAT GGA TGG AAA CGA GCT TTT GAG GAA GAG					
-----					
<u>I S T F H V L I Y G W</u> K R A F E E E					
1630	1639	1648	1657	1666	1675
TAC TAC AGA TTT TAT ACA CCA CCA AAC TTT GTT CTT GCT CTT GTT TTG CCC TCA					
-----					
Y Y R F Y <u>T P P N F V L A L V L P S</u>					
1684	1693	1702	1711	1720	1729
ATT GTA ATT CTG GAT CTT TTG CAG CTT TGC AGA TAC CCA GAC TGA GCT GGA ACT					
-----					
<u>I V I L D L L Q L C</u> R Y P D *					
1738	1747	1756	1765	1774	1783
GGA ATT TGT CTT CCT ATT GAC TCT ACT TCT TTA AAA GCG GCT GCC CAT TAC ATT					
-----					
1792	1801	1810	1819	1828	1837
CCT CAG CTG TCC TTG CAG TTA GGT GTA CAT GTG ACT GAG TGT TGG CCA GTG AGA					
-----					
1846	1855	1864	1873	1882	1891
TGA AGT CTC CTC AAA GGA AGG CAG CAT GTG TCC TTT TTC ATC CCT TCA TCT TGC					
-----					
1900	1909	1918	1927	1936	1945
TGC TGG GAT TGT GGA TAT AAC AGG AGC CCT GGC AGC TGT CTC CAG AGG ATC AAA					
-----					
1954	1963	1972	1981	1990	1999
GCC ACA CCC AAA GAG TAA GGC AGA TTA GAG ACC AGA AAG ACC TTG ACT ACT TCC					
-----					
2008	2017	2026	2035	2044	2053
CTA CTT CCA CTG CTT TTC CTG CAT TTA AGC CAT TGT AAA TCT GGG TGT GTT ACA					
-----					
2062	2071	2080	2089	2098	2107
TGA AGT GAA AAT TAA TTC TTT CTG CCC TTC AGT TCT TTA TCC TGA TAC CAT TTA					
-----					
2116	2125	2134	2143	2152	2161
ACA CTG TCT GAA TTA ACT AGA CTG CAA TAA TTC TTT CTT TTG AAA GCT TTT AAA					
-----					

17/51

FIG. 9D

2170	2179	2188	2197	2206	2215
GGA TAA TGT GCA ATT CAC ATT AAA ATT GAT TTT CCA TTG TCA ATT AGT TAT ACT					
2224	2233	2242	2251	2260	2269
CAT TTT CCT GCC TTG ATC TTT CAT TAG ATA TTT TGT ATC TGC TTG GAA TAT ATT					
2278	2287	2296	2305	2314	2323
ATC TTC TTT TTA ACT GTG TAA TTG GTA ATT ACT AAA ACT CTG TAA TCT CCA AAA					
2332	2341	2350	2359	2368	2377
TAT TGC TAT CAA ATT ACA CAC CAT GTT TTC TAT CAT TCT CAT AGA TCT GCC TTA					
2386	2395	2404	2413	2422	2431
TAA ACA TTT AAA TAA AAA GTA CTA TTT AAT GAT TTA AAA AAA AAA AAA AAA AAA					
2440	2449				
AAA AAA AAA AAA AAA AAA AAA AA 3'					

18/51

## FIG. 10A

```

1  CGAAACTTCC CTCTACCCGC CCGGCCCGCG GCGCGCACCG TTGGCGCTGG ACGCTTCCTC
   GCTTTGAAGG GAGATGGGCG GGCCGGGCGC CGCGCGTGGC AACCGCGACC TGCGAAGGAG

                                     M E K T C I D A L P L T
61  CTTGGAAGCG CCTCTCCCTC AGTTATGGAG AAAACTTGTA TAGATGCACT TCCTCTTACT
   GAACCTTCGC GGAGAGGGAG TCAATACCTC TTTTGAACAT ATCTACGTGA AGGAGAATGA

                                     M N S S E K Q E T V C I F G T G D F G R
121 ATGAATTCTT CAGAAAAGCA AGAGACTGTA TGTATTTTGG GAACTGGTGA TTTTGGAAAGA
   TACTTAAGAA GTCTTTTCGT TCTCTGACAT ACATAAAAAC CTTGACCACT AAAACCTTCT

                                     S L G L K M L Q C G Y S V V F G S R N P
181 TCACTGGGAT TGAAAATGCT CCAGTGTGGT TATTCTGTTG TTTTGGAAAG TCGAAACCCC
   AGTGACCCTA ACTTTTACGA GGTACACCA ATAAGACAAC AAAACCTTC AGCTTTGGGG

                                     Q K T T L L P S G A E V L S Y S E A A K
241 CAGAAGACCA CCCTACTGCC CAGTGGTGCA GAAGTCTTGA GCTATTCAGA AGCAGCCAAG
   GTCTTCTGGT GGGATGACGG GTCACCACGT CTTCAGAACT CGATAAGTCT TCGTCGGTTC

                                     K S G I I I I A I H R E H Y D F L T E L
301 AAGTCTGGCA TCATAATCAT AGCAATCCAC AGAGAGCATT ATGATTTTCT CACAGAATTA
   TTCAGACCGT AGTATTAGTA TCGTTAGGTG TCTCTCGTAA TACTAAAAGA GTGTCTTAAT

                                     T E V L N G K I L V D I S N N L K I N Q
361 ACTGAGGTTT TCAATGGAAA AATATTGGTA GACATCAGCA ACAACCTCAA AATCAATCAA
   TGACTCCAAG AGTTACCTTT TTATAACCAT CTGTAGTCGT TGTGGAGTT TTAGTTAGTT

                                     Y P E S N A E Y L A H L V P G A H V V K
421 TATCCAGAAT CTAATGCAGA GTACCTTGCT CATTTGGTGC CAGGAGCCCA CGTGGTAAAA
   ATAGGTCTTA GATTACGTCT CATGGAACGA GTAAACCACG GTCCTCGGTG GCACCATTTT

                                     A F N T I S A W A L Q S G A L D A S R Q
481 GCATTTAACA CCATCTCAGC CTGGGCTCTC CAGTCAGGAG CACTGGATGC AAGTCGGCAG
   CGTAAATTGT GGTAGAGTCG GACCCGAGAG GTCAGTCCTC GTGACCTACG TTCAGCCGTC

                                     V F V C G N D S K A K Q R V M D I V R N
541 GTGTTTGTGT GTGGAAATGA CAGCAAAGCC AAGCAAAGAG TGATGGATAT TGTTTCGTAAT
   CACAAACACA CACCTTTACT GTCGTTTCGG TTCGTTTCTC ACTACCTATA ACAAGCATTA

                                     L G L T P M D Q G S L M A A K E I E K Y
601 CTTGGACTTA CTCCAATGGA TCAAGGATCA CTCATGGCAG CCAAAGAAAT TGAAAAGTAC
   GAACCTGAAT GAGGTTACCT AGTTCCTAGT GAGTACCGTC GGTTTCTTTA ACTTTTTCATG

                                     P L Q L F P M W R F P F Y L S A V L C V
661 CCCCTGCAGC TATTTCCAAT GTGGAGGTTC CCCTTCTATT TGTCTGCTGT GCTGTGTGTC
   GGGGACGTCG ATAAAGGTTA CACCTCCAAG GGAAGATAA ACAGACGACA CGACACACAG

                                     F L F F Y C V I R D V I Y P Y V Y E K K
721 TTCTTGTTTT TCTATTGTGT TATAAGAGAC GTAATCTACC CTTATGTTTA TGAAAAGAAA
   AAGAACAAA AGATAACACA ATATTCTCTG CATTAGATGG GAATACAAAT ACTTTTCTTT

```



19/51

## FIG. 10B

D N T F R M A I S I P N R I F P I T A L  
 781 GATAATACAT TTCGTATGGC TATTTCCATT CCAAATCGTA TCTTTCCAAT AACAGCACTT  
 CTATTATGTA AAGCATACCG ATAAAGGTAA GGTTTAGCAT AGAAAGGTTA TTGTCGTGAA

T L L A L V Y L P G V I A A I L Q L Y R  
 841 AACTGCTTG CTTTGGTTTA CCTCCCTGGT GTTATTGCTG CCATTCTACA ACTGTACCGA  
 TGTGACGAAC GAAACCAAAT GGAGGGACCA CAATAACGAC GGTAAGATGT TGACATGGCT

G T K Y R R F P D W L D H W M L C R K Q  
 901 GGCACAAAAT ACCGTCGATT CCCAGACTGG CTTGACCACT GGATGCTTTG CCGAAAGCAG  
 CCGTGTTTTA TGGCAGCTAA GGGTCTGACC GAACTGGTGA CCTACGAAAC GGCTTTTCGTC

L G L V A L G F A F L H V L Y T L V I P  
 961 CTTGGCTTGG TAGCTCTGGG ATTTGCCTTC CTTTCATGTCC TCTACACACT TGTGATTCTC  
 GAACCGAACC ATCGAGACCC TAAACGGAAG GAAGTACAGG AGATGTGTGA AACTAAGGA

I R Y Y V R W R L G N L T V T Q A I L K  
 1021 ATTCGATATT ATGTACGATG GAGATTGGGA AACTTAACCG TTACCCAGGC AATACTCAAG  
 TAAGCTATAA TACATGCTAC CTCTAACCCCT TTGAATTGGC AATGGGTCCG TTATGAGTTC

K E N P F S T S S A W L S D S Y V A L G  
 1081 AAGGAGAATC CATTTAGCAC CTCCTCAGCC TGGCTCAGTG ATTCATATGT GGCTTTGGGA  
 TTCCTCTTAG GTAAATCGTG GAGGAGTCGG ACCGAGTCAC TAAGTATACA CCGAAACCCCT

I L G F F L F V L L G I T S L P S V S N  
 1141 ATACTTGGGT TTTTCTGTGTT TGTACTCTTG GGAATCACTT CTTTGCCATC TGTTAGCAAT  
 TATGAACCCA AAAAAGACAA ACATGAGAAC CCTTAGTGAA GAAACGGTAG ACAATCGTTA

A V N W R E F R F V Q S K L G Y L T L I  
 1201 GCAGTCAACT GGAGAGAGTT CCGATTGTG CAGTCCAAAC TGGGTTATTT GACCCTGATC  
 CGTCAGTTGA CCTCTCTCAA GGCTAAACAG GTCAGGTTTG ACCCAATAAA CTGGGACTAG

L C T A H T L V Y G G K R F L S P S N L  
 1261 TTGTGTACAG CCCACACCCCT GGTGTACGGT GGGAAAGAGAT TCCTCAGCCC TTCAAATCTC  
 AACACATGTC GGGTGTGGGA CCACATGCCA CCCTTCTCTA AGGAGTCGGG AAGTTTAGAG

R W Y L P A A Y V L G L I I P C T V L V  
 1321 AGATGGTATC TTCCTGCAGC CTACGTGTTA GGGCTTATCA TTCCTTGCAC TGTGCTGGTG  
 TCTACCATAG AAGGACGTCG GATGCACAAT CCCGAATAGT AAGGAACGTG ACACGACCAC

I K F V L I M P C V D N T L T R I R Q G  
 1381 ATCAAGTTTG TCCTAATCAT GCCATGTGTA GACAACACCC TTACAAGGAT CCGCCAGGGC  
 TAGTTCAAAC AGGATTAGTA CGGTACACAT CTGTTGTGGG AATGTTCTTA GGCGGTCCCG

W E R N S K H  
 1441 TGGGAAAGGA ACTCAAAACA CTAGAAAAAG CATTGAATGG AAAATCAATA TTTAAAACAA  
 ACCCTTTCCT TGAGTTTTGT GATCTTTTTC GTAACCTACC TTTTAGTTAT AAATTTTGTT

20/51

## FIG. 10C

1501	AGTTCAATTT	AGCTGGATTT	CTGAACTATG	GTTTTGAATG	TTTAAAGAAG	AATGATGGGT
	TCAAGTTAAA	TCGACCTAAA	GACTTGATAC	CAAAACTTAC	AAATTTCTTC	TTACTACCCA
1561	ACAGTTAGGA	AAGTTTTTTT	CTTACACCGT	GACTGAGGGA	AACATTGCTT	GTCTTTGAGA
	TGTCATCCT	TTCAAAAAAA	GAATGTGGCA	CTGACTCCCT	TTGTAACGAA	CAGAAACTCT
1621	AATTGACTGA	CATACTGGAA	GAGAACACCA	TTTTATCTCA	GGTTAGTGAA	GAATCAGTGC
	TTAACTGACT	GTATGACCTT	CTCTTGTTGGT	AAAATAGAGT	CCAATCACTT	CTTAGTCACG
1681	AGGTCCCTGA	CTCTTATTTT	CCCAGAGGCC	ATGGAGCTGA	GATTGAGACT	AGCCTTGTGG
	TCCAGGGACT	GAGAATAAAA	GGGTCTCCCG	TACCTCGACT	CTAACTCTGA	TCGGAACACC
1741	TTTCACACTA	AAGAGTTTCC	TTGTTATGGG	CAACATGCAT	GACCTAATGT	CTTGCAAAAT
	AAAGTGTGAT	TTCTCAAAGG	AACAATACCC	GTTGTACGTA	CTGGATTACA	GAACGTTTTA
1801	CCAATAGAAG	TATTGCAGCT	TCCTTCTCTG	GCTCAAGGGC	TGAGTTAAGT	GAAAGGAAAA
	GGTTATCTTC	ATAACGTCGA	AGGAAGAGAC	CGAGTTCCCG	ACTCAATTCA	CTTTCCTTTT
1861	ACAGCACAAT	GGTGACCACT	GATAAAGGCT	TTATTAGGTA	TATCTGAGGA	AGTGGGTCAC
	TGTCGTGTTA	CCACTGGTGA	CTATTTCCGA	AATAATCCAT	ATAGACTCCT	TCACCCAGTG
1921	ATGAAATGTA	AAAAGGGAAT	GAGGTTTTTG	TTGTTTTTTG	GAAGTAAAGG	CAAACATAAA
	TACTTTACAT	TTTTCCCTTA	CTCCAAAAAC	AACAAAAAAC	CTTCATTTCC	GTTTGTATTT
1981	TATTACCATG	ATGAATTCTA	GTGAAATGAC	CCCTTGACTT	TGCTTTTCTT	AATACAGATA
	ATAATGGTAC	TACTTAAGAT	CACTTTACTG	GGGAACTGAA	ACGAAAAGAA	TTATGTCTAT
2041	TTTACTGAGA	GGAACATTTT	TTATAACACA	AGAAAAATTT	ACAATTGATT	AAAAGTATCC
	AAATGACTCT	CCTTGATAAA	AATATTGTGT	TCTTTTAA	TGTTAACTAA	TTTTCATAGG
2101	ATGTCTTGGA	TACATACGTA	TCTATAGAGC	TGGCATGTAA	TTCTTCCTCT	ATAAAGAATA
	TACAGAACCT	ATGTATGCAT	AGATATCTCG	ACCGTACATT	AAGAAGGAGA	TATTTCTTAT
2161	GGTATAGGAA	AGACTGAATA	AAAATGGAGG	GATATCCCCT	TGGATTTTAC	TTGCATTGTG
	CCATATCCTT	TCTGACTTAT	TTTTACCTCC	CTATAGGGGA	ACCTAAAGTG	AACGTAACAC
2221	CAATAAGCAA	AGAAGGGTTG	ATAAAAAGTT	TTGATCAAAA	AGTTCAAAGA	AACCAGAATT
	GTTATTCGTT	TCTTCCCAAC	TATTTTCAAG	AACTAGTTTT	TCAAGTTTCT	TTGGTCTTAA
2281	TTAGACAGCA	AGCTAAATAA	ATATTGTAAA	ATTGCACTAT	ATTAGGTAA	GTATTATTTA
	AATCTGTCGT	TCGATTTATT	TATAACATTT	TAACGTGATA	TAATCCAATT	CATAATAAAT
2341	GGTATTATAA	TATGCTTTGT	AAATTTTATA	TTCCAAATAT	TGCTCAATAT	TTTTCATCTA
	CCATAATATT	ATACGAAACA	TTTAAAAATAT	AAGGTTTATA	ACGAGTTATA	AAAAGTAGAT
2401	TTAAATTAAT	TTCTAGTGTA	AATAAGTAGC	TTCTATATCT	GTCTTAGTCT	ATTATAATTG
	AATTTAATTA	AAGATCACAT	TTATTCATCG	AAGATATAGA	CAGAATCAGA	TAATATTAAC

21/51

## FIG. 10D

2461	TAAGGAGTAA	AATTAAATGA	ATAGTCTGCA	GGTATAAATT	TGAACAATGC	ATAGATGATC
	ATTCCTCATT	TTAATTTACT	TATCAGACGT	CCATATTTAA	ACTTGTTACG	TATCTACTAG
2521	GAAAATTACG	GAAAATCATA	GGGCAGAGAG	GTGTGAAGAT	TCATCATTAT	GTGAAATTTG
	CTTTTAATGC	CTTTTAGTAT	CCCGTCTCTC	CACACTTCTA	AGTAGTAATA	CACTTTAAAC
2581	GATCTTTCTC	AAATCCTTGC	TGAAATTTAG	GATGGTTCTC	ACTGTTTTTC	TGTGCTGATA
	CTAGAAAGAG	TTTAGGAACG	ACTTTAAATC	CTACCAAGAG	TGACAAAAAG	ACACGACTAT
2641	GTACCCTTTC	CAAGGTGACC	TTCAGGGGGA	TTAACCTTCC	TAGCTCAAGC	AATGAGCTAA
	CATGGGAAAG	GTTCCACTGG	AAGTCCCCCT	AATTGGAAGG	ATCGAGTTCG	TTACTCGATT
2701	AAGGAGCCTT	ATGCATGATC	TTCCACATA	TCAAAATAAC	TAAAAGGCAC	TGAGTTTGGC
	TTCTCGGAA	TACGTACTAG	AAGGGTGTAT	AGTTTTATTG	ATTTTCCGTG	ACTCAAACCG
2761	ATTTTTCTGC	CTGCTCTGCT	AAGACCTTTT	TTTTTTTTTT	ACTTTCATTA	TAACATATTA
	TAAAAAGACG	GACGAGACGA	TTCTGGAAAA	AAAAAAAAAA	TGAAAGTAAT	ATTGTATAAT
2821	TACATGACAT	TATACAAAAA	TGATTAAAAAT	ATATTAAAAC	AACATCAACA	ATCCAGGATA
	ATGTACTGTA	ATATGTTTTT	ACTAATTTTA	TATAATTTTG	TTGTAGTTGT	TAGGTCCTAT
2881	TTTTTCTATA	AAACTTTTTA	AAAATAATTG	TATCTATATA	TTCAATTTTA	CATCCTTTTT
	AAAAAGATAT	TTTGAAAAAT	TTTTATTAAC	ATAGATATAT	AAGTTAAAAT	GTAGGAAAAA
2941	CAAAGGCTTT	GTTTTTCTAA	AGGCTTTGTT	TTCTTTTTTA	TTATTTTTTT	CTTTTTTATT
	GTTTCCGAAA	CAAAAAGATT	TCCGAAACAA	AAGGAAAAAT	AATAAAAAAA	GAAAAAATAA
3001	TTTTTGAGAC	AGTCTTGCTC	TGTCGCTCAG	GCTGGAGTGC	AGTGGCACGA	TCTCAGCTCA
	AAAAACTCTG	TCAGAACGAG	ACAGCGAGTC	CGACCTCACG	TCACCGTGCT	AGAGTCGAGT
3061	CTGCAACCTC	CTCCTCCCAG	GTTCAAGTGA	TTCTTGTTCA	TCAGCCTCCC	GAGTAGCTGG
	GACGTTGGAG	GAGGAGGGTC	CAAGTTCACT	AAGAACAAGT	AGTCGGAGGG	CTCATCGACC
3121	GACTACAGGC	ATGTGCCACT	ATGCCCAGCT	AATTTTTGTA	CTTTTAGTAG	AGACAGGGTT
	CTGATGTCCG	TACACGGTGA	TACGGGTCGA	TTAAAAACAT	GAAAATCATC	TCTGTCCCAA
3181	TCACCACATT	GGTCAGGCTG	GTCTTGAAAT	GCTGGCGTCA	AGTGATCTGC	CTGCCTCCGC
	AGTGGTGTA	CCAGTCCGAC	CAGAACTTTA	CGACCGCAGT	TACTAGACG	GACGGAGGCG
3241	CTTACGTAAT	ATATTTTCTT	AATGGCTGCA	TAATATCACA	TCAAATAGGC	ATTTTTCAAA
	GAATGCATTA	TATAAAAGAA	TTACCGACGT	ATTATAGTGT	AGTTTATCCG	TAAAAAGTTT
3301	CCTCTTTCCT	TATTAAACAT	GTAGACTATA	TCCATTTTTT	ACTAAAATAA	ATAACATTTT
	GGAGAAAGGA	ATAATTTGTA	CATCTGATAT	AGGTAAAAAA	TGATTTTATT	TATTGTAAAG
3361	AGATAATATC	TTTGCACTGA	TAATGTTGCC	AAGCCATTTT	TAAAGTGACC	TTATCAATTT
	TCTATTATAG	AAACGTGACT	ATTACAACGG	TTCGGTAAAG	ATTTCACTGG	AATAGTTAAA

22/51

## FIG. 10E

3421	AATTACCATT	GGATGAGGGT	GTTGCTTTCA	TCGCACCATT	GTAGATTGTC	TTTTTTATTT
	TTAATGGTAA	CCTACTCCCA	CAACGAAAGT	AGCGTGGTAA	CATCTAACAG	AAAAAATAAA
3481	CAATTTGCGT	TTATTTATAA	CTGGTTGCAA	AGGTACACAG	AACACACGCT	CCTTCAACTT
	GTAAACGCA	AATAAATATT	GACCAACGTT	TCCATGTGTC	TTGTGTGCGA	GGAAGTTGAA
3541	ATCTTTGATA	AACCCAAGCA	AGGATACAAA	AAGTTGGACG	ACATTGAGTA	GAGTCATGGT
	TAGAAACTAT	TTGGGTTCGT	TCCTATGTTT	TTCAACCTGC	TGTAACTCAT	CTCAGTACCA
3601	ATACGGTGCT	GACCCTACAG	TATCAGTGGA	AAAGATAAGG	AAAATGTCAC	TACTCACCTA
	TATGCCACGA	CTGGGATGTC	ATAGTCACCT	TTTCTATTCC	TTTTACAGTG	ATGAGTGGAT
3661	TGTTATGCAA	AACAGTTAGG	TGTGCTGGGG	CTGGATACTG	CTCTTTTACT	TGAGCATTGG
	ACAATACGTT	TTGTCAATCC	ACACGACCCC	GACCTATGAC	GAGAAAATGA	ACTCGTAACC
3721	TTGATTAAAG	TTTAGGTACC	ATCCAGGCTG	GTCTAGAGAA	GTCTTTGGAG	TTAACCATGC
	AACTAATTTT	AAATCCATGG	TAGGTCCGAC	CAGATCTCTT	CAGAAACCTC	AATTGGTACG
3781	TCTTTTTGTT	AAAGAAGAGA	GTAATGTGTT	TATCCTGGCT	CATAGTCCGT	CACCGAAAAT
	AGAAAAACAA	TTTCTTCTCT	CATTACACAA	ATAGGACCGA	GTATCAGGCA	GTGGCTTTTA
3841	AGAAAATGCC	ATCCATAGGT	AAAATGCTGA	CCTATAGAAA	AAAATGAACT	CTACTTTTAT
	TCTTTTACGG	TAGGTATCCA	TTTTACGACT	GGATATCTTT	TTTACTTGA	GATGAAAATA
3901	AGCCTAGTAA	AAATGCTCTA	CCTGAGTAGT	TAAAAGCAAT	TCATGAAGCC	TGAAGCTAAA
	TCGGATCATT	TTTACGAGAT	GGACTCATCA	ATTTTCGTTA	AGTACTTCGG	ACTTCGATTT
3961	GAGCACTCTG	ATGGTTTTTG	CATAATAGCT	GCATTTCCAG	ACCTGACCTT	TGGCCCCAAC
	CTCGTGAGAC	TACCAAAACC	GTATTATCGA	CGTAAAGGTC	TGGACTGGAA	ACCGGGGTTG
4021	CACAAGTGCT	CCAAGCCCCA	CCAGCTGACC	AAAGAAAGCC	CAAGTTCTCC	TTCTGTCTTT
	GTGTTACACG	GGTTCGGGGT	GGTCGACTGG	TTTCTTTTCG	GTTCAAGAGG	AAGACAGGAA
4081	CCCACAACCT	CCCTGCTCCC	AAAACATATG	AATTAATTTG	ACCATATTAA	CACAGCTGAC
	GGGTGTTGGA	GGGACGAGGG	TTTTGATACT	TTAATTAAAC	TGGTATAATT	GTGTCGACTG
4141	TCCTCCAGTT	TACTTAAGGT	AGAAAGAATG	AGTTTACAAC	AGATGAAAAT	AAGTGCTTTG
	AGGAGGTCAA	ATGAATTCCA	TCTTTCTTAC	TCAAATGTTG	TCTACTTTTA	TTCACGAAAC
4201	GGCGAACTGT	ATTCTTTTTA	ACAGATCCAA	ACTATTTTAC	ATTTAAAAAA	AAAGTTAAAC
	CCGCTTGACA	TAAGGAAAAT	TGTCTAGGTT	TGATAAAATG	TAAATTTTTT	TTTCAATTTG
4261	TAAACTTCTT	TACTGCTGAT	ATGTTTCCTG	TATTCTAGAA	AAATTTTTTAC	ACTTTCACAT
	ATTTGAAGAA	ATGACGACTA	TACAAAGGAC	ATAAGATCTT	TTTAAAAATG	TGAAAGTGTA
4321	TATTTTTGTA	CACTTTCCCC	ATGTTAAGGG	ATGATGGCTT	TTATAAATGT	GTATTCATTA
	ATAAAAACAT	GTGAAAGGGG	TACAATTCCC	TACTACCGAA	AATATTTACA	CATAAGTAAT
4381	AATGTTACTT	TAAAAATAAA	AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	
	TTACAATGAA	ATTTTTATTT	TTTTTTTTTT	TTTTTTTTTT	TTTTTTTTTT	

23/51

## FIG. 10F

STEAP-2, AA508880 (NCI\_CGAP Pr6)

ggtcgacttttccctttattcctttgtcagagatctgattcatccatattgctagaaaccaacagagtgacttttaca  
aaattccctatagagattgtgaataaaaaccttacctatagttgccattactttgctctccctagtataccttgcagg  
tcttctggcagctgcttatcaactttattacggcaccaagtataggagatttccaccttggttggaacctgggtta  
cagtgtagaaaacagcttggtactaagttgtttcttcgctatggtccatgttgctacagcctctgcttaccca  
tgagaaggtcagagagat

STEAP-2, 98P4B6 SSH fragment

TTTGAGCTTTGCAGATACCCAGACTGAGCTGGAAGTGAATTTGTCTTCCTATTGACTCTACTTCTTTAAAGCG  
GCTGCCCATTAACATTCTCAGCTGTCCTTGACAGTTAGGTGTACATGTGACTGAGTGTGGCCAGTGAGATGAAGTC  
TCCTCAAAGGAAGGCAGCATGTGTCCTTTTT

STEAP-3, AI139607 (testis EST)

aagaaggagaatccatttagcacctcctcagcctggctcagtgattcatatgtggctttgggaataacttgggtttt  
ttctgtttgtactcttgggaatcacttctttgccatctgttagcaatgcagtcactggagagagtccgatttgt  
ccagtcctaaactgggttatttgaccctgatcttgtgtacagcccacaccctggtgtacggtgggaagagattcctc  
agcccttcaaactctcagatgggtatcttctcgcagcctacgtgttagggcttatcattccttgcactgtgctgggtga  
tcaagtttgtcctaatacatgccatgtgtagacaacacccttacaaggatccgccagggtgggaaggaactcaaa  
acactagaaaaagcattgaatggaaaatcaatatttaaaacaaagtccaatttagctggaaaaaaaaa

STEAP 4, R80991 (placental EST)

ggccgcggcanccgctacgacctgggtcaacctggcagtcagcaggtcttgccanacaagagccacctctgggtg  
aaggaggaggtctggcgatggagatctacctctccctgggagtgctggccctcggcacggtgtccctgctggccg  
tgacctcactgccgtccattgcaaactcgctcaactggagggagttcagcttcgttcagtcctcactgggctttgt  
ggcctnctgtgctgagcacactncacacgctcacctacggctggacccgcgccttcgaggagagccgctacaagttc  
tacctnctcccacctcacgntcacgctgctgggtgcctcgcttcggttcacctgggcccaggccctgtttntac  
tgccctgcattcagccgnaga

FIG. 11A

1	15	16	30	31	45	46	60	61	75	76	90
2	STEAP2	MESISMMGSPKSLSE	TC- - -LPNGINGIKD	ARKVTVGWIGSGDFA	KSLTIRLIRCGYHVV	IGSRNPKFASEFFPH	VVDVTHHEDALTKTN	87			
3	STEAP3	-----MEK	TCIDALPLTNSSSE-	-KQETVCIFGTGDFG	RSLGLKMLQCGYSVV	FGSRNPQKTT-LLPS	GAEVLSYSEAAKKS	75			
4	STEAP1	-----	-----	-----	-----	-----	-----	0			
5	STEAP4	-----	-----	-----	-----	-----	-----	0			
91	105	106	120	121	135	136	150	151	165	166	180
2	STEAP2	TEFVAIHREHYTSIM	DLRHLLVGKILIDVS	NNMRINQYPESNAEY	LASLFPDLSLVKGFN	VVSAAWALQIGPKDAS	QOVYICSNNTQARQQ	177			
3	STEAP3	ITIIIAIHREHYDELT	ELTEVLNGKILIDIS	NNLKINQYPESNAEY	LAHLVPGAHVVKAFN	TISAAWALQSGALDAS	QOVFVCGNDSKAKQR	165			
4	STEAP1	-----	-----	-----	-----MESRKDITN	QEELWKNK-PRRNLE	EDDYLL--HKDTGETS	36			
5	STEAP4	-----	-----	-----	-----	-----	-----	0			
181	195	196	210	211	225	226	240	241	255	256	270
2	STEAP2	WIELARQLNF-IPID	LGSLSASAREIENLPL	RLFTLWRGPVVVAIS	LATFEEFLYSFVRDVI	HPYARNQSSDFYKIP	TEETVKNKTLPIVAITL	266			
3	STEAP3	YMDIVRNIGL-TPMD	QGSIMAAKEIEKYPL	QLFPPWRFPFYL SAV	LCVFELEFYCVIRDVI	YPYVYVEKDKNTFRMA	HSPNRIFFPTALTLL	254			
4	STEAP1	MLKRPVLLHHQTAH	ADEFDCPSELQH-TQ	ELFPQWHLPIKIAAI	IASUTFLYTLLEREVI	HPLATSHQQYFYKIP	ILLVINKVLPMVVSITL	125			
5	STEAP4	-----	-----	-----	-----	-----	-----	0			
271	285	286	300	301	315	316	330	331	345	346	360
2	STEAP2	LSLVYLAAGLLAAAYQ	LYYGTKYRRFPFPWLE	TWLCRKQQLGLLSEF	FAMVHVAYSICLPMR	RSERYLLFLNMAYQQV	HANIENSWNEEEVWR	356			
3	STEAP3	LALVYLPGVIAAILQ	LYRGTKYRRFPDMLD	HWMLCRKQLGLVALG	FATLHVLYTIVIPIR	YYVRWRIGNLITVQA	ILKKENPFSTSSAWL	344			
4	STEAP1	LALVYLPGVIAAIVQ	LHNGTKYKKFPFPHLD	KWMLTRKQFGLLSEF	FAVLHAIFYSHSYPMR	RSYRYKLLNWAYQQV	QONKEDAWIEHDVWR	215			
5	STEAP4	-----	-----	-----	-----A	AAAXATTWSTWQSSRS	WPXKSHLWVKEEVR	31			



```

STEAP-1      67  LFPQWHLPIKIAAIIASLTFLYTLLEVIHPLATSHQQYFYKIPILVINKVLPMSITLL
STEAP-2     208  LFTLWRGPVVVAISLATFFFLYSFVRDVIHPYARNQQSDFYKIPIEIVNKTLP IVAITLL
               **  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *
STEAP-1     127  ALVYLPGVIAAIVQLHNGTKYKKFPHWLDKMWLTRKQFGLLSFFF AVLHAIYSLSPMRR
STEAP-2     268  SLVYLAGLLAAAYQLYYGTYKRRFPWPLETWLQCRKQLGLLSFFF AMVHVAYSLCLPMRR
               **** *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *
STEAP-1     187  SYRYKLLNWAYQQVQONKEDAWIEHDVWRMEIYVSLGIVGLAILALLAVTSIPSVSDSLT
STEAP-2     328  SERYLFLNMAYQQVHANIENSWNEEVWRIEMYISFGTMSLGLLSLLAVTSIPSVSNALN
               *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *  *
STEAP-1     247  WREFHYIQSKLGIVSLLLGTHALIFAWNKWIDIKQFVWYTPPTFMIAVFLPIVVLI
STEAP-2     388  WREFSFIQSTLGYVALLISTFHVLIYGWKRAFEEEYRYFTPPNFVLALVLPISVIL
               ***** *  *  *  *  *  *  *  *  *  *  *  *  *  *  *

```



27/51

## FIG. 11D

```
STEAP1  66  ELFPQWHLPIKIAAIIASLTFLYTLLREVIHPLATSHQQYFYKIPILVINKVLPMSITL
STEAP3 195  QLFPMWRFPPFYLSAVLCVFLFFYCVIRDVIYPYVEKKDNTFRMAISIPNRIFPITALTL
          *** * * * * * * * * * * * * * * * * * * * * * * *
STEAP1 126  LALVYLPGVIAAIVQLHNGTKYKKFPHWLDKWMLTRKQFGLLSFFFFAVLHAIYSLSPMR
STEAP3 255  LALVYLPGVIAAILQLYRGTKYRRFPDWLDHWMLCRKQLGLVALGFAFLHVLYTLVIPIR
          ***** ** * * * * * * * * * * * * * * * * * * * *
STEAP1 186  RSYRYKLLNWAYQQVQQNKEDAWIEHVDVRMEIYVSLGIVGLAILALLAVTSIPSVSDSL
STEAP3 315  YYVRWRLGNLTVTQAILKKENPFSTSSAWLSDSYVALGILGFFLFVLLGITSLSVSNV
          * * * * * * * * * * * * * * * * * * * * * * * *
STEAP1 246  TWREFHYIQSKLGIVSLLLGTIHALIFAWNKWIDIKQFVWYTPPTFMIAVFLPIVVLIK
STEAP3 375  NWREFRFVQSKLGYLTLILCTAHTLVYGGKRFLSPSNLRWYLPAAAYVLGLIIPCTVLVIK
          **** * * * * * * * * * * * * * * * * * * * * * *
STEAP1 306  SILFLPCLRKKILKIRHWEDVTK
STEAP3 435  FVLIMPCVDNTLTRIRQGWERNK
          * ** * * * *
```

28/51

## FIG. 11E

```
STEAP2  29  RKVTVGVIGSGDFAKSLTIRLIRCGYHVVIGSRNPKFASEFFPHVVDVTHHEDALTKTNI
STEAP3  18  KQETVCIFGTGDFGRSLGLKMLQCGYSVVFGSRNPQ-KTTLLPSGAEVLSYSEAAKSGI
          **      *  ***   **          ***  **  *****          *      *      *  *  *

STEAP2  89  IFVAIHREHYTSLWDLRHLLVGKILIDVSNMRINQYPESNAEYLASLFPDSLIVKGFNV
STEAP3  77  IIIIAIHREHYDFLTEVLNGKILVDISNNLKINQYPESNAEYLAHLVPGAHVVKAFNT
          *  *****   *  *      *  *****  *  ***  *****  *  *      **  **

STEAP2 149  VSAWALQLGPKDASRQVYICSNNIQARQQVIELARQLNFIPIIDLGLSSAREIENLPLRL
STEAP3 137  ISAWALQSGALDASRQVFVCGNDSKAKQVRMDIVRNGLTPMDQGSMAAKEIEKYPLQL
          *****  *  *****   *  *      *  *  *      *  *      *  *  ***  *  *  *


STEAP2 209  FTLWRGPVVVAISLATFFFLYSFVRDVIHPYARNQQSDFYKIPIEIVNKTLPIVAITLLS
STEAP3 197  FPMWRFPFYLSAVLCVFLFFYCVIRDVIYPYVEKKDNTFRMAISIPNRIFPITALTLA
          *  **  *      *  *  *  *      *****  *      *  *  *      **  *  ***


STEAP2 269  LVYLAGLLAAAYQLYYGTKYRRFPWLETWLQCRKQLGLLSFFFAMVHVAYSCLCLPMRRS
STEAP3 257  LVYLPGVIAAILQLYRGTKYRRFPDWLDHWMLCRKQLGLVALGFAFLHVLYTLVIPIRY
          *****  *  **   ***  *****  **  *  *****          **  **  *  *  *  *

STEAP2 329  ERYLFLNMAYQQVHANIENSWNEEEVWRIEMYISFGIMSLGLLSLLAVTSIPSVSNALNW
STEAP3 317  VRWRLGNLTVTQAILKKENPFSTSSAWLSDSYVALGILGFFLFVLLGITSLSVSNVNW
          *      *      *      **          *      *      **      *  **  **  *****  **

STEAP2 389  REFSFIQSTLGYVALLISTFHVLIYGWKRAFEEEYRFYTPPNFVLALVLPISIVIL
STEAP3 377  REFRFVQSKLGYLTLILCTAHTLVYGGKRFLSPSNLRWYLPAAAYVLGLIIPCTVLV
          ***  *  **  ***   *      *  *  *  *  *      *  *  *      **  *  *  *
```

FIG. 12A





31/51




FIG.14A




FIG.14B




FIG. 15A




FIG. 15B




FIG. 15C




FIG. 16






FIG. 17





FIG. 18



37/51

**FIG. 19**

38/51


**FIG. 20**

Fig. 21B




Fig. 21D



Fig. 21A



Fig. 21C




40/51

**Fig. 22A**




**Fig. 22B**



41/51

**Fig. 23A**



**Fig. 23B**





Fig. 24





43/51



**Fig. 25**

FIG. 26






FIG. 27A






FIG. 27B







FIG. 30




Fig. 31A




Fig. 31B




Fig. 31C




Fig. 31D




Fig. 31E





Fig. 31F



50/51

Fig. 32A




Fig. 32B







FIG. 33



## SEQUENCE LISTING

&lt;110&gt; UroGenesys, Inc.

<120> NOVEL SERPENTINE TRANSMEMBRANE ANTIGENS  
EXPRESSED IN HUMAN CANCERS AND USES THEREOF

&lt;130&gt; 129.17WOI1

&lt;150&gt; 09/455,486

&lt;151&gt; 1999-12-06

&lt;160&gt; 47

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 1130

&lt;212&gt; DNA

&lt;213&gt; Homo Sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (64)...(1083)

&lt;400&gt; 1

atggaaagca gaaaagacat cacaaaccaa gaagaacttt ggaaaatgaa gcctaggaga	60
aat tta gaa gaa gac gat tat ttg cat aag gac acg gga gag acc agc	108
Leu Glu Glu Asp Asp Tyr Leu His Lys Asp Thr Gly Glu Thr Ser	
1 5 10 15	
atg cta aaa aga cct gtg ctt ttg cat ttg cac caa aca gcc cat gct	156
Met Leu Lys Arg Pro Val Leu Leu His Leu His Gln Thr Ala His Ala	
20 25 30	
gat gaa ttt gac tgc cct tca gaa ctt cag cac aca cag gaa ctc ttt	204
Asp Glu Phe Asp Cys Pro Ser Glu Leu Gln His Thr Gln Glu Leu Phe	
35 40 45	
cca cag tgg cac ttg cca att aaa ata gct gct att ata gca tct ctg	252
Pro Gln Trp His Leu Pro Ile Lys Ile Ala Ala Ile Ile Ala Ser Leu	
50 55 60	
act ttt ctt tac act ctt ctg agg gaa gta att cac cct tta gca act	300
Thr Phe Leu Tyr Thr Leu Leu Arg Glu Val Ile His Pro Leu Ala Thr	
65 70 75	
tcc cat caa caa tat ttt tat aaa att cca atc ctg gtc atc aac aaa	348
Ser His Gln Gln Tyr Phe Tyr Lys Ile Pro Ile Leu Val Ile Asn Lys	
80 85 90 95	
gtc ttg cca atg gtt tcc atc act ctc ttg gca ttg gtt tac ctg cca	396
Val Leu Pro Met Val Ser Ile Thr Leu Leu Ala Leu Val Tyr Leu Pro	
100 105 110	
ggc gtg ata gca gca att gtc caa ctt cat aat gga acc aag tat aag	444
Gly Val Ile Ala Ala Ile Val Gln Leu His Asn Gly Thr Lys Tyr Lys	
115 120 125	
aag ttt cca cat tgg ttg gat aag tgg atg tta aca aga aag cag ttt	492
Lys Phe Pro His Trp Leu Asp Lys Trp Met Leu Thr Arg Lys Gln Phe	
130 135 140	

```

ggg ctt ctc agt ttc ttt ttt gct gta ctg cat gca att tat agt ctg      540
Gly Leu Leu Ser Phe Phe Phe Ala Val Leu His Ala Ile Tyr Ser Leu
   145                               150                               155

tct tac cca atg agg cga tcc tac aga tac aag ttg cta aac tgg gca      588
Ser Tyr Pro Met Arg Arg Ser Tyr Arg Tyr Lys Leu Leu Asn Trp Ala
   160                               165                               170                               175

tat caa cag gtc caa caa aat aaa gaa gat gcc tgg att gag cat gat      636
Tyr Gln Gln Val Gln Gln Asn Lys Glu Asp Ala Trp Ile Glu His Asp
                               180                               185                               190

gtt tgg aga atg gag att tat gtg tct ctg gga att gtg gga ttg gca      684
Val Trp Arg Met Glu Ile Tyr Val Ser Leu Gly Ile Val Gly Leu Ala
                               195                               200                               205

ata ctg gct ctg ttg gct gtg aca tct att cca tct gtg agt gac tct      732
Ile Leu Ala Leu Leu Ala Val Thr Ser Ile Pro Ser Val Ser Asp Ser
                               210                               215                               220

ttg aca tgg aga gaa ttt cac tat att cag agc aag cta gga att gtt      780
Leu Thr Trp Arg Glu Phe His Tyr Ile Gln Ser Lys Leu Gly Ile Val
   225                               230                               235

tcc ctt cta ctg ggc aca ata cac gca ttg att ttt gcc tgg aat aag      828
Ser Leu Leu Leu Gly Thr Ile His Ala Leu Ile Phe Ala Trp Asn Lys
   240                               245                               250                               255

tgg ata gat ata aaa caa ttt gta tgg tat aca cct cca act ttt atg      876
Trp Ile Asp Ile Lys Gln Phe Val Trp Tyr Thr Pro Pro Thr Phe Met
                               260                               265                               270

ata gct gtt ttc ctt cca att gtt gtc ctg ata ttt aaa agc ata cta      924
Ile Ala Val Phe Leu Pro Ile Val Val Leu Ile Phe Lys Ser Ile Leu
                               275                               280                               285

ttc ctg cca tgc ttg agg aag aag ata ctg aag att aga cat ggt tgg      972
Phe Leu Pro Cys Leu Arg Lys Lys Ile Leu Lys Ile Arg His Gly Trp
                               290                               295                               300

gaa gac gtc acc aaa att aac aaa act gag ata tgt tcc cag ttg tag     1020
Glu Asp Val Thr Lys Ile Asn Lys Thr Glu Ile Cys Ser Gln Leu *
   305                               310                               315

aat tac tgt tta cac aca ttt ttg ttc aat att gat ata ttt tat cac     1068
Asn Tyr Cys Leu His Thr Phe Leu Phe Asn Ile Asp Ile Phe Tyr His
   320                               325                               330

caa cat ttc aag ttt gtatttggtta ataaaatgat tattcaagga aaaaaaaaaa     1123
Gln His Phe Lys Phe
   335

aaaaaaaaa                                                                1130

<210> 2
<211> 339
<212> PRT
<213> Homo Sapiens

<400> 2
Leu Glu Glu Asp Asp Tyr Leu His Lys Asp Thr Gly Glu Thr Ser Met
   1           5           10           15
Leu Lys Arg Pro Val Leu Leu His Leu His Gln Thr Ala His Ala Asp

```

[illegible]

```
<210> 3
<211> 111
<212> DNA
<213> Homo Sapiens
```

```
<400> 3
ggcggaggcg gaggcggagg gcgaggggcg gggagcgccg cctggagcgc ggcaggtcat      60
attgaacatt ccagatacct atcattactc gatgctgttg ataacagcaa g                    111
```

```
<210> 4
<211> 24
<212> DNA
<213> Artificial Sequence
```

```
<220>
<223> primer
```

```
<400> 4
actttgttga tgaccaggat tggg
```

```
<210> 5
<211> 24
```

<212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer

<400> 5  
 cagaacttca gcacacacag gaac

24

<210> 6  
 <211> 3627  
 <212> DNA  
 <213> Homo Sapiens

<400> 6  
 ggggcccgca cctctgggca gcagcggcag ccgagactca cggcgaagct aaggcgaaga 60  
 gtgggtgggt gaagccatac ttttttatag aattaatgga aagcagaaaa gacatcacaa 120  
 accaagaaga actttggaaa atgaagccta ggagaaattt agaagaagac gattatttgc 180  
 ataaggacac gggagagacc agcatgctaa aaagacctgt gcttttgcac ttgcacccaa 240  
 cagcccatgc tgatgaattt gactgccctt cagaacttca gcacacacag gaactctttc 300  
 cacagtggca cttgcccaatt aaaatagctg ctattatagc atctctgact tttctttaca 360  
 ctctctcgag ggaagtaatt cacccttag caacttccca tcaacaatat ttttataaaa 420  
 ttccaatcct ggtcatcaac aaagtcttgc caatgggttc catcactctc ttggcattgg 480  
 tttacctgcc aggtgtgata gcagcaattg tccaacttca taatggaacc aagtataaga 540  
 agtttccaca ttggttggtt aagtggatgt taacaagaaa gcagtttggg cttctcagtt 600  
 tcttttttgc tgtactgcat gcaatttata gtctgtctta cccaatgagg cgatcctaca 660  
 gatacaagtt gctaaactgg gcatatcaac aggtccaaca aaataaagaa gatgcctgga 720  
 ttgagcatga tgtttggaga atggagattt atgtgtctct ggggaattgtg ggattggcaa 780  
 tactggctct gttggctgtg acatctattc catctgtgag tgactctttg acatggagag 840  
 aatttcacta tattcaggta aataatatat aaaataaccc taagaggtaa atcttctttt 900  
 tgtgtttatg atatagaata tgttgacttt accccataaa aaataacaaa tgtttttcaa 960  
 cagcaaagat cttataactg ttccaattaa taatgtgtct tcctgttgtt ttccctattg 1020  
 cttctaatta ggacaagtgt ttcttagaca taaataaaaag gcattaaaaat attcttttgt 1080  
 tttttttttt tgtttgtttt tttttgtttt gtttgtttgt ttttttgaga tgaagtctcg 1140  
 ctctgttgcc catgctggag tacagtggca cgatctcgcc tcaactgcaac ctgcgcctcc 1200  
 tgggttcagg cgattctctt gcctcagcct cctgagtagc tgggattaca ggcacccatc 1260  
 accatgtcca gctaattttt gtatttttag tagagacagg gttttcccat gttggccagg 1320  
 ctggctctga tctcctgacc tcaaagtatc cgccacctc ggcctcccaa agtgcctgga 1380  
 tgacagttgt gagccaccac actcagcctg ctctttctaa tatttgaaac ttgttagaca 1440  
 atttgctacc catctaattg gatatttttag gaatccaata tgcattggtt attatttctt 1500  
 aaaaaaata ttcttttacc tgtcacctga atttagtaat gccttttatg ttacacaact 1560  
 tagcactttc cagaacacaa aactctctcc ttgaataat agagttttta tctaccaaag 1620  
 atatgtagt gtctcatttc aaaggctgct ttttccagct tacattttat atacttactc 1680  
 acttgaagtt tcttaaatatt ctgttaattt taaaactatc tcagatttac tgaggtttat 1740  
 cttctggtgg tagattatcc ataagaagag tgatgtgcca gaatcactct gggatccttg 1800  
 tctgacaaga ttcaaaggac taaatttaat tcagtcata acactgcca ttaccgttta 1860  
 tgggtagaca tctttggaaa tttccacaag gtcagacatt cgcaactatc ccttctacat 1920  
 gtccacacgt atactccaac actttattag gcatctgatt agtttggaaa gtatgcctcc 1980  
 atctgaatta gtccagtgtg gcttagagtt ggtacaacat tctcacagaa tttcctaatt 2040  
 ttgtagggtc agcctgataa ccactggagt tctttggtcc tcattaaata gctttcttca 2100  
 cacattgctc tgctgtttac acatagatg aacactgctt ttttagactt attaggaatt 2160  
 taggactgca tcttgacaac tgagcctatt ctactatatg tacaatacct agcccataat 2220  
 aggtatacaa tacacatttg gtaaaactaa ttttcaacca atgacatgta tttttcaact 2280  
 agtaacctag aaatgtttca cttaaaatct gagaactggg tacactacaa gttaccttgg 2340  
 agattcatat atgaaaacgc aaacttagct atttgattgt attcactggg acttaagaat 2400  
 gcgcctgaat aattgtgagt tcgatttgtt ctggcaggct aatgaccatt tccagtaaag 2460  
 tgaatagagg tcagaagtgc tataaaagag gtgtgtgcag aacaccgttg agattacata 2520  
 ggtgaacaac tatttttaag caactttatt tgtgtagtga caaagcatcc caatgcaggc 2580  
 tgaaatgttt catcacatct ctggatctct ctattttgtg cagacattga aaaaattgtt 2640  
 catatttttt ccatgttatc agaattttg atttttttaa aacataggcc aagttcattc 2700  
 acttcattat tcatattatc aaatcagagt gaatcacatt agtcgccttc acaactgata 2760  
 aagatcactg aagtcaaatt gattttttgt ataattctca atctacctat atttaattga 2820  
 gaatctaaaa tgtacaaatc attgtgttga ttctgcagt atcctgctat aagtaagact 2880  
 cagtccttga ttttaggtat cctgtgaaaa gcagaattaa gacaaatata caagagacaa 2940

```

agcacaaaaa ataaatatca taaggggatg aacaaaatgg tggagaaaaga gtagacaaag 3000
tttttgatca cctgccttca aagaaaggct gtgaattttg ttcacttaga cagcttggag 3060
acaagaaatt acccaaaagt aaggtgagga ggataggcaa aaagagcaga aagatgtgaa 3120
tggacattgt tgagaaatgt gataggaaaa caatcataga taaaggattt ccaagcaaca 3180
gagcatatcc agatgaggta ggatgggata aactcttatt gaaccaatct tcaccaatct 3240
tgtttttctt ttgcagagca agctaggaat tgtttccctt ctactgggca caatacacgc 3300
attgattttt gcctggaata agtggataga tataaaacaa tttgtatggg atacacctcc 3360
aacttttatg atagctgttt tccttccaat tgttgcctg atatttaaaa gcatactatt 3420
cctgccatgc ttgaggaaaga agatactgaa gattagacat gggtgggaag acgtcaccaa 3480
aattaacaaa actgagatat gttcccagtt gtagaattac tgtttacaca catttttggt 3540
caatattgat atattttatc accaacattt caagtttgta tttgttaata aaatgattat 3600
tcaaggaaaa aaaaaaaaaa aaaaaaa 3627

```

&lt;210&gt; 7

&lt;211&gt; 2099

&lt;212&gt; DNA

&lt;213&gt; Homo Sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (355)...(1719)

&lt;400&gt; 7

```

atggaatcaa tctctatgat gggaagccct aagagcctta gtgaaacttg tttacctaat 60
ggcataaatg gtatcaaaga tgcaaggaag gtcactgtag gtgtgattgg aagtggagat 120
tttgccaaat ccttgaccat tcgacttatt agatgaggct atcatgtggg cataggaagt 180
agaaatccta agtttgcttc tgaatttttt cctcatgtgg tagatgtcac tcatcatgaa 240
gatgctctca caaaaacaaa tataatatgt gttgctatac acagagaaca ttatacctcc 300
ctgtggggacc tgagacatct gcttgtgggt aaaatcctga ttgatgtgag caat aac 357
                                     Asn
                                     1

```

```

atg agg ata aac cag tac cca gaa tcc aat gct gaa tat ttg gct tca 405
Met Arg Ile Asn Gln Tyr Pro Glu Ser Asn Ala Glu Tyr Leu Ala Ser
                5                10                15

```

```

tta ttc cca gat tct ttg att gtc aaa gga ttt aat gtt gtc tca gct 453
Leu Phe Pro Asp Ser Leu Ile Val Lys Gly Phe Asn Val Val Ser Ala
                20                25                30

```

```

tgg gca ctt cag tta gga cct aag gat gcc agc cgg cag gtt tat ata 501
Trp Ala Leu Gln Leu Gly Pro Lys Asp Ala Ser Arg Gln Val Tyr Ile
                35                40                45

```

```

tgc agc aac aat att caa gcg cga caa cag gtt att gaa ctt gcc cgc 549
Cys Ser Asn Asn Ile Gln Ala Arg Gln Gln Val Ile Glu Leu Ala Arg
                50                55                60                65

```

```

cag ttg aat ttc att ccc att gac ttg gga tcc tta tca tca gcc aga 597
Gln Leu Asn Phe Ile Pro Ile Asp Leu Gly Ser Leu Ser Ser Ala Arg
                70                75                80

```

```

gag att gaa aat tta ccc cta cga ctc ttt act ctc tgg aga ggg cca 645
Glu Ile Glu Asn Leu Pro Leu Arg Leu Phe Thr Leu Trp Arg Gly Pro
                85                90                95

```

```

gtg gtg gta gct ata agc ttg gcc aca ttt ttt ttc ctt tat tcc ttt 693
Val Val Val Ala Ile Ser Leu Ala Thr Phe Phe Phe Leu Tyr Ser Phe
                100                105                110

```

```

gtc aga gat gtg att cat cca tat gct aga aac caa cag agt gac ttt 741
Val Arg Asp Val Ile His Pro Tyr Ala Arg Asn Gln Gln Ser Asp Phe
                115                120                125

```

tac aaa att cct ata gag att gtg aat aaa acc tta cct ata gtt gcc Tyr Lys Ile Pro Ile Glu Ile Val Asn Lys Thr Leu Pro Ile Val Ala 130 135 140 145	789
att act ttg ctc tcc cta gta tac ctt gca ggt ctt ctg gca gct gct Ile Thr Leu Leu Ser Leu Val Tyr Leu Ala Gly Leu Leu Ala Ala Ala 150 155 160	837
tat caa ctt tat tac ggc acc aag tat agg aga ttt cca cct tgg ttg Tyr Gln Leu Tyr Tyr Gly Thr Lys Tyr Arg Arg Phe Pro Pro Trp Leu 165 170 175	885
gaa acc tgg tta cag tgt aga aaa cag ctt gga tta cta agt ttt ttc Glu Thr Trp Leu Gln Cys Arg Lys Gln Leu Gly Leu Leu Ser Phe Phe 180 185 190	933
ttc gct atg gtc cat gtt gcc tac agc ctc tgc tta ccg atg aga agg Phe Ala Met Val His Val Ala Tyr Ser Leu Cys Leu Pro Met Arg Arg 195 200 205	981
tca gag aga tat ttg ttt ctc aac atg gct tat cag cag gtt cat gca Ser Glu Arg Tyr Leu Phe Leu Asn Met Ala Tyr Gln Gln Val His Ala 210 215 220 225	1029
aat att gaa aac tct tgg aat gag gaa gaa gtt tgg aga att gaa atg Asn Ile Glu Asn Ser Trp Asn Glu Glu Glu Val Trp Arg Ile Glu Met 230 235 240	1077
tat atc tcc ttt ggc ata atg agc ctt ggc tta ctt tcc ctc ctg gca Tyr Ile Ser Phe Gly Ile Met Ser Leu Gly Leu Leu Ser Leu Leu Ala 245 250 255	1125
gtc act tct atc cct tca gtg agc aat gct tta aac tgg aga gaa ttc Val Thr Ser Ile Pro Ser Val Ser Asn Ala Leu Asn Trp Arg Glu Phe 260 265 270	1173
agt ttt att cag tct aca ctt gga tat gtc gct ctg ctc ata agt act Ser Phe Ile Gln Ser Thr Leu Gly Tyr Val Ala Leu Leu Ile Ser Thr 275 280 285	1221
ttc cat gtt tta att tat gga tgg aaa cga gct ttt gag gaa gag tac Phe His Val Leu Ile Tyr Gly Trp Lys Arg Ala Phe Glu Glu Glu Tyr 290 295 300 305	1269
tac aga ttt tat aca cca cca aac ttt gtt ctt gct ctt gtt ttg ccc Tyr Arg Phe Tyr Thr Pro Pro Asn Phe Val Leu Ala Leu Val Leu Pro 310 315 320	1317
tca att gta att ctg gat ctt ttg cag ctt tgc aga tac cca gac tga Ser Ile Val Ile Leu Asp Leu Leu Gln Leu Cys Arg Tyr Pro Asp * 325 330 335	1365
gct gga act gga att tgt ctt cct att gac tct act tct tta aaa gcg Ala Gly Thr Gly Ile Cys Leu Pro Ile Asp Ser Thr Ser Leu Lys Ala 340 345 350	1413
gct gcc cat tac att cct cag ctg tcc ttg cag tta ggt gta cat gtg Ala Ala His Tyr Ile Pro Gln Leu Ser Leu Gln Leu Gly Val His Val 355 360 365	1461
act gag tgt tgg cca gtg aga tga agt ctc ctc aaa gga agg cag cat Thr Glu Cys Trp Pro Val Arg * Ser Leu Leu Lys Gly Arg Gln His 370 375 380	1509

```

gtg tcc ttt ttc atc cct tca tct tgc tgc tgg gat tgt gga tat aac      1557
Val Ser Phe Phe Ile Pro Ser Ser Cys Cys Trp Asp Cys Gly Tyr Asn
    385                      390                      395

agg agc cct ggc agc tgt ctc cag agg atc aaa gcc aca ccc aaa gag      1605
Arg Ser Pro Gly Ser Cys Leu Gln Arg Ile Lys Ala Thr Pro Lys Glu
    400                      405                      410                      415

taa ggc aga tta gag acc aga aag acc ttg act act tcc cta ctt cca      1653
* Gly Arg Leu Glu Thr Arg Lys Thr Leu Thr Thr Ser Leu Leu Pro
    420                      425                      430

ctg ctt ttc ctg cat tta agc cat tgt aaa tct ggg tgt gtt aca tga      1701
Leu Leu Phe Leu His Leu Ser His Cys Lys Ser Gly Cys Val Thr *
    435                      440                      445

agt gaa aat taa ttc ttt ctgcccttca gttctttatc ctgataccat      1749
Ser Glu Asn * Phe Phe
    450

ttaacactgt ctgaattaac tagactgcaa taattctttc ttttgaaagc ttttaaagga      1809
taatgtgcaa ttcacattaa aattgatttt ccattgtcaa ttagttatac tcattttcct      1869
gccttgatct ttcattagat attttgtatc tgcttggaat atattatctt ctttttaact      1929
gtgtaattgg taattactaa aactctgtaa tctccaaaat attgctatca aattacacac      1989
catgttttct atcattctca tagatctgcc ttataaacat ttaaataaaaa agtactattt      2049
aatgatttaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      2099

```

<210> 8  
 <211> 450  
 <212> PRT  
 <213> Homo Sapiens

```

<400> 8
Asn Met Arg Ile Asn Gln Tyr Pro Glu Ser Asn Ala Glu Tyr Leu Ala
 1          5          10          15
Ser Leu Phe Pro Asp Ser Leu Ile Val Lys Gly Phe Asn Val Val Ser
    20          25          30
Ala Trp Ala Leu Gln Leu Gly Pro Lys Asp Ala Ser Arg Gln Val Tyr
    35          40          45
Ile Cys Ser Asn Asn Ile Gln Ala Arg Gln Gln Val Ile Glu Leu Ala
    50          55          60
Arg Gln Leu Asn Phe Ile Pro Ile Asp Leu Gly Ser Leu Ser Ser Ala
    65          70          75          80
Arg Glu Ile Glu Asn Leu Pro Leu Arg Leu Phe Thr Leu Trp Arg Gly
    85          90          95
Pro Val Val Val Ala Ile Ser Leu Ala Thr Phe Phe Phe Leu Tyr Ser
    100          105          110
Phe Val Arg Asp Val Ile His Pro Tyr Ala Arg Asn Gln Gln Ser Asp
    115          120          125
Phe Tyr Lys Ile Pro Ile Glu Ile Val Asn Lys Thr Leu Pro Ile Val
    130          135          140
Ala Ile Thr Leu Leu Ser Leu Val Tyr Leu Ala Gly Leu Leu Ala Ala
    145          150          155          160
Ala Tyr Gln Leu Tyr Tyr Gly Thr Lys Tyr Arg Arg Phe Pro Pro Trp
    165          170          175
Leu Glu Thr Trp Leu Gln Cys Arg Lys Gln Leu Gly Leu Leu Ser Phe
    180          185          190
Phe Phe Ala Met Val His Val Ala Tyr Ser Leu Cys Leu Pro Met Arg
    195          200          205
Arg Ser Glu Arg Tyr Leu Phe Leu Asn Met Ala Tyr Gln Gln Val His
    210          215          220
Ala Asn Ile Glu Asn Ser Trp Asn Glu Glu Glu Val Trp Arg Ile Glu
    225          230          235          240
Met Tyr Ile Ser Phe Gly Ile Met Ser Leu Gly Leu Leu Ser Leu Leu

```



245 250 255  
 Ala Val Thr Ser Ile Pro Ser Val Ser Asn Ala Leu Asn Trp Arg Glu  
 260 265 270  
 Phe Ser Phe Ile Gln Ser Thr Leu Gly Tyr Val Ala Leu Leu Ile Ser  
 275 280 285  
 Thr Phe His Val Leu Ile Tyr Gly Trp Lys Arg Ala Phe Glu Glu Glu  
 290 295 300  
 Tyr Tyr Arg Phe Tyr Thr Pro Pro Asn Phe Val Leu Ala Leu Val Leu  
 305 310 315 320  
 Pro Ser Ile Val Ile Leu Asp Leu Leu Gln Leu Cys Arg Tyr Pro Asp  
 325 330 335  
 Ala Gly Thr Gly Ile Cys Leu Pro Ile Asp Ser Thr Ser Leu Lys Ala  
 340 345 350  
 Ala Ala His Tyr Ile Pro Gln Leu Ser Leu Gln Leu Gly Val His Val  
 355 360 365  
 Thr Glu Cys Trp Pro Val Arg Ser Leu Leu Lys Gly Arg Gln His Val  
 370 375 380  
 Ser Phe Phe Ile Pro Ser Cys Cys Trp Asp Cys Gly Tyr Asn Arg  
 385 390 395 400  
 Ser Pro Gly Ser Cys Leu Gln Arg Ile Lys Ala Thr Pro Lys Glu Gly  
 405 410 415  
 Arg Leu Glu Thr Arg Lys Thr Leu Thr Thr Ser Leu Leu Pro Leu Leu  
 420 425 430  
 Phe Leu His Leu Ser His Cys Lys Ser Gly Cys Val Thr Ser Glu Asn  
 435 440 445  
 Phe Phe  
 450

<210> 9  
 <211> 4419  
 <212> DNA  
 <213> Homo Sapiens

<220>  
 <221> CDS  
 <222> (85)...(1464)

<400> 9  
 cgaaacttcc ctctaccgcg ccggcccgcg gcgcgcaccg ttggcgctgg acgcttcctc 60  
 cttggaagcg cctctccctc agtt atg gag aaa act tgt ata gat gca ctt 111  
 Met Glu Lys Thr Cys Ile Asp Ala Leu  
 1 5  
  
 cct ctt act atg aat tct tca gaa aag caa gag act gta tgt att ttt 159  
 Pro Leu Thr Met Asn Ser Ser Glu Lys Gln Glu Thr Val Cys Ile Phe  
 10 15 20 25  
  
 gga act ggt gat ttt gga aga tca ctg gga ttg aaa atg ctc cag tgt 207  
 Gly Thr Gly Asp Phe Gly Arg Ser Leu Gly Leu Lys Met Leu Gln Cys  
 30 35 40  
  
 ggt tat tct gtt gtt ttt gga agt cga aac ccc cag aag acc acc cta 255  
 Gly Tyr Ser Val Val Phe Gly Ser Arg Asn Pro Gln Lys Thr Thr Leu  
 45 50 55  
  
 ctg ccc agt ggt gca gaa gtc ttg agc tat tca gaa gca gcc aag aag 303  
 Leu Pro Ser Gly Ala Glu Val Leu Ser Tyr Ser Glu Ala Ala Lys Lys  
 60 65 70  
  
 tct ggc atc ata atc ata gca atc cac aga gag cat tat gat ttt ctc 351  
 Ser Gly Ile Ile Ile Ile Ala Ile His Arg Glu His Tyr Asp Phe Leu  
 75 80 85  
  
 aca gaa tta act gag gtt ctc aat gga aaa ata ttg gta gac atc agc 399

Thr	Glu	Leu	Thr	Glu	Val	Leu	Asn	Gly	Lys	Ile	Leu	Val	Asp	Ile	Ser	
90					95					100					105	
aac	aac	ctc	aaa	atc	aat	caa	tat	cca	gaa	tct	aat	gca	gag	tac	ctt	447
Asn	Asn	Leu	Lys	Ile	Asn	Gln	Tyr	Pro	Glu	Ser	Asn	Ala	Glu	Tyr	Leu	
				110					115					120		
gct	cat	ttg	gtg	cca	gga	gcc	cac	gtg	gta	aaa	gca	ttt	aac	acc	atc	495
Ala	His	Leu	Val	Pro	Gly	Ala	His	Val	Val	Lys	Ala	Phe	Asn	Thr	Ile	
			125					130					135			
tca	gcc	tgg	gct	ctc	cag	tca	gga	gca	ctg	gat	gca	agt	cgg	cag	gtg	543
Ser	Ala	Trp	Ala	Leu	Gln	Ser	Gly	Ala	Leu	Asp	Ala	Ser	Arg	Gln	Val	
		140					145					150				
ttt	gtg	tgt	gga	aat	gac	agc	aaa	gcc	aag	caa	aga	gtg	atg	gat	att	591
Phe	Val	Cys	Gly	Asn	Asp	Ser	Lys	Ala	Lys	Gln	Arg	Val	Met	Asp	Ile	
	155					160					165					
gtt	cgt	aat	ctt	gga	ctt	act	cca	atg	gat	caa	gga	tca	ctc	atg	gca	639
Val	Arg	Asn	Leu	Gly	Leu	Thr	Pro	Met	Asp	Gln	Gly	Ser	Leu	Met	Ala	
170					175					180					185	
gcc	aaa	gaa	att	gaa	aag	tac	ccc	ctg	cag	cta	ttt	cca	atg	tgg	agg	687
Ala	Lys	Glu	Ile	Glu	Lys	Tyr	Pro	Leu	Gln	Leu	Phe	Pro	Met	Trp	Arg	
				190					195					200		
ttc	ccc	ttc	tat	ttg	tct	gct	gtg	ctg	tgt	gtc	ttc	ttg	ttt	ttc	tat	735
Phe	Pro	Phe	Tyr	Leu	Ser	Ala	Val	Leu	Cys	Val	Phe	Leu	Phe	Phe	Tyr	
			205					210					215			
tgt	gtt	ata	aga	gac	gta	atc	tac	cct	tat	gtt	tat	gaa	aag	aaa	gat	783
Cys	Val	Ile	Arg	Asp	Val	Ile	Tyr	Pro	Tyr	Val	Tyr	Glu	Lys	Lys	Asp	
		220					225					230				
aat	aca	ttt	cgt	atg	gct	att	tcc	att	cca	aat	cgt	atc	ttt	cca	ata	831
Asn	Thr	Phe	Arg	Met	Ala	Ile	Ser	Ile	Pro	Asn	Arg	Ile	Phe	Pro	Ile	
		235				240					245					
aca	gca	ctt	aca	ctg	ctt	gct	ttg	gtt	tac	ctc	cct	ggt	gtt	att	gct	879
Thr	Ala	Leu	Thr	Leu	Leu	Ala	Leu	Val	Tyr	Leu	Pro	Gly	Val	Ile	Ala	
250					255					260					265	
gcc	att	cta	caa	ctg	tac	cga	ggc	aca	aaa	tac	cgt	cga	ttc	cca	gac	927
Ala	Ile	Leu	Gln	Leu	Tyr	Arg	Gly	Thr	Lys	Tyr	Arg	Arg	Phe	Pro	Asp	
			270						275					280		
tgg	ctt	gac	cac	tgg	atg	ctt	tgc	cga	aag	cag	ctt	ggc	ttg	gta	gct	975
Trp	Leu	Asp	His	Trp	Met	Leu	Cys	Arg	Lys	Gln	Leu	Gly	Leu	Val	Ala	
			285					290					295			
ctg	gga	ttt	gcc	ttc	ctt	cat	gtc	ctc	tac	aca	ctt	gtg	att	cct	att	1023
Leu	Gly	Phe	Ala	Phe	Leu	His	Val	Leu	Tyr	Thr	Leu	Val	Ile	Pro	Ile	
		300					305					310				
cga	tat	tat	gta	cga	tgg	aga	ttg	gga	aac	tta	acc	gtt	acc	cag	gca	1071
Arg	Tyr	Tyr	Val	Arg	Trp	Arg	Leu	Gly	Asn	Leu	Thr	Val	Thr	Gln	Ala	
		315				320					325					
ata	ctc	aag	aag	gag	aat	cca	ttt	agc	acc	tcc	tca	gcc	tgg	ctc	agt	1119
Ile	Leu	Lys	Lys	Glu	Asn	Pro	Phe	Ser	Thr	Ser	Ser	Ala	Trp	Leu	Ser	
330					335					340					345	
gat	tca	tat	gtg	gct	ttg	gga	ata	ctt	ggg	ttt	ttt	ctg	ttt	gta	ctc	1167

Asp Ser Tyr Val	Ala Leu Gly Ile	Leu Gly Phe Phe Leu Phe Val Leu	
	350	355	360
ttg gga atc act tct ttg cca tct gtt agc aat gca gtc aac tgg aga			1215
Leu Gly Ile Thr Ser Leu Pro Ser Val Ser Asn Ala Val Asn Trp Arg			
	365	370	375
gag ttc cga ttt gtc cag tcc aaa ctg ggt tat ttg acc ctg atc ttg			1263
Glu Phe Arg Phe Val Gln Ser Lys Leu Gly Tyr Leu Thr Leu Ile Leu			
	380	385	390
tgt aca gcc cac acc ctg gtg tac ggt ggg aag aga ttc ctc agc cct			1311
Cys Thr Ala His Thr Leu Val Tyr Gly Gly Lys Arg Phe Leu Ser Pro			
	395	400	405
tca aat ctc aga tgg tat ctt cct gca gcc tac gtg tta ggg ctt atc			1359
Ser Asn Leu Arg Trp Tyr Leu Pro Ala Ala Tyr Val Leu Gly Leu Ile			
	410	415	420
att cct tgc aca tca agt ttg tcc taa tca tgc cat gtg tag aca aca			1407
Ile Pro Cys Thr Ser Ser Leu Ser * Ser Cys His Val * Thr Thr			
	430	435	
ccc tta caa gga tcc gcc agg gct ggg aaa gga act caa aac act aga			1455
Pro Leu Gln Gly Ser Ala Arg Ala Gly Lys Gly Thr Gln Asn Thr Arg			
	440	445	450
aaa agc att gaatggaaaa tcaatatatta aaacaaagtt caatttagct			1504
Lys Ser Ile			

ggattttctga	actatgggtt	tgaatgttta	aagaagaatg	atgggtacag	ttaggaaagt	1564
ttttttctta	caccgtgact	gagggaaaca	ttgcttgtct	ttgagaaatt	gactgacata	1624
ctggaagaga	acaccatttt	atctcagggt	agtgaagaat	cagtgcaggt	ccctgactct	1684
tattttccca	gaggccatgg	agctgagatt	gagactagcc	ttgtgggttc	acactaaaga	1744
gtttccttgt	tatgggcaac	atgcatgacc	taatgtcttg	caaaatccaa	tagaagtatt	1804
gcagcttctt	tctctggctc	aagggctgag	ttaagtgaag	ggaaaaacag	cacaatgggt	1864
accactgata	aaggctttat	taggtatatc	tgaggaaagt	ggtcacatga	aatgtaaaaa	1924
gggaatgagg	tttttgggtg	tttttggagg	taaaggcaaa	cataaatatt	accatgatga	1984
attctagtga	aatgacccct	tgactttgct	tttcttaata	cagatatatta	ctgagaggaa	2044
ctatttttat	aacacaagaa	aaatttacaa	ttgattaaaa	gtatccatgt	cttggatata	2104
tacgtatcta	tagagctggc	atgtaattct	tcctctataa	agaataggta	taggaaagac	2164
tgaataaaaa	tggagggata	tccccttggg	tttccactgc	attgtgcaat	aagcaaaagaa	2224
gggttgataa	aagttcttga	tcaaaaagtt	caaagaaacc	agaatttttag	acagcaagct	2284
aaataaatat	tgtaaaattg	cactatatta	gggttaagat	tatttaggta	ttataaatatg	2344
ctttgtaaat	tttatattcc	aaatattgct	caatattttt	catctattaa	attaatttct	2404
agtgtaaata	agtagcttct	atatctgtct	tagtctatta	taattgtaag	gagtaaaatt	2464
aaatgaatag	tctgcaggta	taaatttgaa	caatgcatag	atgacgaaa	attacggaaa	2524
atcatagggc	agagagggtg	gaagattcat	cattatgtga	aatttggatc	tttctcaaat	2584
ccttgctgaa	atttaggatg	gttctcactg	tttttctgtg	ctgatagtac	cctttccaag	2644
gtgaccttca	gggggattaa	ccttccttagc	tcaagcaatg	agctaaaagg	agccttatgc	2704
atgatcttcc	cacatatcaa	aataactaaa	aggcactgag	tttggcattt	ttctgcctgc	2764
tttgctaaga	cctttttttt	ttttttactt	tcattataac	atattataca	tgacattata	2824
caaaaatgat	taaaatatat	taaaacaaca	tcaacaatcc	aggatatatt	tctataaaaac	2884
tttttaaaaa	taattgtatc	tatatattca	attttacatc	ctttttcaaa	ggctttgttt	2944
ttctaaaggc	tttggtttcc	tttttattat	ttttttcttt	tttatttttt	tgagacagtc	3004
ttgctctgtc	gctcaggctg	gagtgcagtg	gccacgatctc	agctcactgc	aacctcctcc	3064
cccagggttc	aagtgtattct	tgttcatcag	cctcccgagt	agctgggact	acaggcatgt	3124
gccactatgc	ccagctaatt	tttgtacttt	tagtagagag	agggtttcac	cacattgggtc	3184
aggctgggtc	tgaaatgctg	gcgtcaagtg	atctgcctgc	ctccgcctta	cgtaatatat	3244
tttcttaatg	gctgcataat	atcacatcaa	ataggcatatt	ttcaaacctc	tttccttatt	3304
aaacatgtag	actatatcca	ttttttacta	aaataaataa	catttcagat	aatatctttg	3364
cactgataat	gttgccaagc	catttctaaa	gtgaccttat	caatttaatt	accattggat	3424
gaggggtgtg	ctttcatcgc	accattgtag	atgtgtcttt	ttatttcaat	ttgcgtttat	3484

```

ttataactgg ttgcaaaggt acacagaaca cacgctcctt caacttatct ttgataaacc 3544
caagcaagga tacaaaaagt tggacgacat tgagtagagt catgggtatac ggtgctgacc 3604
ctacagtatc agtggaaaaag ataaggaaaa tgctactact cacctatggt atgcaaaaca 3664
gttaggtgtg ctggggctgg atactgctct tttacttgag cattgggtga ttaaagttta 3724
ggtaccatcc aggctggtct agagaagtct ttggagttaa ccatgctctt tttgttaaag 3784
aagagagtaa tgtgtttatc ctggctcata gtcgctcacc gaaaatagaa aatgccatcc 3844
ataggtaaaa tgctgaccta tagaaaaaaa tgaactctac ttttatagcc tagtaaaaat 3904
gctctacctg agtagttaa agcaattcat gaagcctgaa gctaaagagc actctgatgg 3964
ttttggcata atagctgcat ttccagacct gacctttggc cccaaccaca agtgctccaa 4024
gccccaccag ctgaccaaag aaagcccaag ttctccttct gtccttccca caacctccct 4084
gctcccaaaa ctatgaaatt aatttgacca tattaacaca gctgactcct ccagtttact 4144
taaggtagaa agaatgagtt tacaacagat gaaaataagt gctttgggag aactgtattc 4204
cttttaacag atccaaacta ttttacattt aaaaaaaaaa ttaaactaaa cttctttact 4264
gctgatatgt ttctgtatt ctagaaaaat ttttacactt tcacattatt tttgtacact 4324
ttccccatgt taagggatga tggtttttat aaatgtgtat tcattaaatg ttactttaaa 4384
aataaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 4419

```

&lt;210&gt; 10

&lt;211&gt; 458

&lt;212&gt; PRT

&lt;213&gt; Homo Sapiens

&lt;400&gt; 10

```

Met Glu Lys Thr Cys Ile Asp Ala Leu Pro Leu Thr Met Asn Ser Ser
 1          5          10          15
Glu Lys Gln Glu Thr Val Cys Ile Phe Gly Thr Gly Asp Phe Gly Arg
 20          25          30
Ser Leu Gly Leu Lys Met Leu Gln Cys Gly Tyr Ser Val Val Phe Gly
 35          40          45
Ser Arg Asn Pro Gln Lys Thr Thr Leu Leu Pro Ser Gly Ala Glu Val
 50          55          60
Leu Ser Tyr Ser Glu Ala Lys Lys Ser Gly Ile Ile Ile Ile Ala
 65          70          75          80
Ile His Arg Glu His Tyr Asp Phe Leu Thr Glu Leu Thr Glu Val Leu
 85          90          95
Asn Gly Lys Ile Leu Val Asp Ile Ser Asn Asn Leu Lys Ile Asn Gln
100          105          110
Tyr Pro Glu Ser Asn Ala Glu Tyr Leu Ala His Leu Val Pro Gly Ala
115          120          125
His Val Val Lys Ala Phe Asn Thr Ile Ser Ala Trp Ala Leu Gln Ser
130          135          140
Gly Ala Leu Asp Ala Ser Arg Gln Val Phe Val Cys Gly Asn Asp Ser
145          150          155          160
Lys Ala Lys Gln Arg Val Met Asp Ile Val Arg Asn Leu Gly Leu Thr
165          170          175          180
Pro Met Asp Gln Gly Ser Leu Met Ala Ala Lys Glu Ile Glu Lys Tyr
180          185          190
Pro Leu Gln Leu Phe Pro Met Trp Arg Phe Pro Phe Tyr Leu Ser Ala
195          200          205
Val Leu Cys Val Phe Leu Phe Phe Tyr Cys Val Ile Arg Asp Val Ile
210          215          220
Tyr Pro Tyr Val Tyr Glu Lys Lys Asp Asn Thr Phe Arg Met Ala Ile
225          230          235          240
Ser Ile Pro Asn Arg Ile Phe Pro Ile Thr Ala Leu Thr Leu Leu Ala
245          250          255
Leu Val Tyr Leu Pro Gly Val Ile Ala Ala Ile Leu Gln Leu Tyr Arg
260          265          270
Gly Thr Lys Tyr Arg Arg Phe Pro Asp Trp Leu Asp His Trp Met Leu
275          280          285
Cys Arg Lys Gln Leu Gly Leu Val Ala Leu Gly Phe Ala Phe Leu His
290          295          300
Val Leu Tyr Thr Leu Val Ile Pro Ile Arg Tyr Tyr Val Arg Trp Arg
305          310          315          320
Leu Gly Asn Leu Thr Val Thr Gln Ala Ile Leu Lys Lys Glu Asn Pro

```

```

          325          330          335
Phe Ser Thr Ser Ser Ala Trp Leu Ser Asp Ser Tyr Val Ala Leu Gly
          340          345          350
Ile Leu Gly Phe Phe Leu Phe Val Leu Leu Gly Ile Thr Ser Leu Pro
          355          360          365
Ser Val Ser Asn Ala Val Asn Trp Arg Glu Phe Arg Phe Val Gln Ser
          370          375          380
Lys Leu Gly Tyr Leu Thr Leu Ile Leu Cys Thr Ala His Thr Leu Val
          385          390          395          400
Tyr Gly Gly Lys Arg Phe Leu Ser Pro Ser Asn Leu Arg Trp Tyr Leu
          405          410          415
Pro Ala Ala Tyr Val Leu Gly Leu Ile Ile Pro Cys Thr Ser Ser Leu
          420          425          430
Ser Ser Cys His Val Thr Thr Pro Leu Gln Gly Ser Ala Arg Ala Gly
          435          440          445
Lys Gly Thr Gln Asn Thr Arg Lys Ser Ile
          450          455

```

&lt;210&gt; 11

&lt;211&gt; 322

&lt;212&gt; DNA

&lt;213&gt; Homo Sapiens

&lt;400&gt; 11

```

ggctcgacttt tcctttatttc ctttgtcaga gatctgattc atccatatgc tagaaaccaa      60
cagagtgactt tttacaaaat tcctatagag attgtgaata aaaccttacc tatagttgcc      120
attacttttgc tctccctagt ataccttgca ggtcttctgg cagctgctta tcaactttat      180
tacggcacca agtataggag atttccacct tggttggaaa cctggttaca gtgtagaaaa      240
cagcttggat tactaagttg tttcttcgct atgggtccatg ttgcctacag cctctgctta      300
ccgatgagaa ggtcagagag at

```

&lt;210&gt; 12

&lt;211&gt; 183

&lt;212&gt; DNA

&lt;213&gt; Homo Sapiens

&lt;400&gt; 12

```

tttgcagctt tgcagatacc cagactgagc tggaaactgga atttgtcttc ctattgactc      60
tacttcttta aaagcggctg cccattacat tcctcagctg tccttgagct taggtgtaca      120
tgtgactgag tgttggccag tgagatgaag tctcctcaaa ggaaggcagc atgtgtcctt      180
ttt

```

&lt;210&gt; 13

&lt;211&gt; 448

&lt;212&gt; DNA

&lt;213&gt; Homo Sapiens

&lt;400&gt; 13

```

aagaaggaga atccatttag cacctcctca gcctggctca gtgattcata tgtggctttg      60
ggaatacttg ggttttttct gtttgtactc ttgggaatca cttctttgcc atctgttagc      120
aatgcagtca actggagaga gttccgattt gtccagtcga aactgggtta tttgacctg      180
atcttgtgta cagccacac cctggtgtac ggtgggaaga gattcctcag cccttcaaat      240
ctcagatggt atcttcctgc agcctacgtg ttagggctta tcattccttg cactgtgctg      300
gtgatcaagt ttgtccta atgcatgt gtagacaaca cccttacaag gatccgccag      360
ggctgggaaa ggaactcaaa aactagaaa aagcattgaa tggaaaatca atatttaaaa      420
caaagtcaaa tttagctgga aaaaaaaa

```

&lt;210&gt; 14

&lt;211&gt; 401

&lt;212&gt; DNA

&lt;213&gt; Homo Sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

<222> 11, 56, 233, 250, 310, 326, 377, 398

<223> n = a, t, c, or g

<400> 14

```

ggccgcggca nccgctacga cctgggtcaac ctggcagtcag agcaggtctt ggccanacaa      60
gagccacctc tgggtgaagg aggaggtctg gcggatggag atctacctct ccctggggagt      120
gctggccctc ggcacgttgt ccctgctggc cgtgacctca ctgccgtcca ttgcaaactc      180
gctcaactgg agggagtcca gcttcgttca gtcctcactg ggctttgtgg ccntcgtgct      240
gagcacactn cacacgctca cctacggctg gaccgcgcc ttcgaggaga gccgctacaa      300
gttctacctn cctccacact tcacgntcac gctgctgggt ccctgcgttc gttcatcctg      360
ggccaaagcc ctgtttntac tgccttgcat tcagccgnag a                          401

```

<210> 15

<211> 118

<212> PRT

<213> Homo Sapiens

<220>

<223> Xaa = Any amino acid

<400> 15

```

Ala Ala Ala Xaa Ala Thr Thr Trp Ser Thr Trp Gln Ser Ser Arg Ser
 1          5          10          15
Met Glu Ile Tyr Leu Ser Leu Gly Val Leu Ala Leu Gly Thr Leu Ser
      20          25          30
Leu Leu Ala Val Thr Ser Leu Pro Ser Ile Ala Asn Ser Leu Asn Trp
      35          40          45
Arg Glu Phe Ser Phe Val Gln Ser Ser Leu Gly Phe Val Ala Xaa Val
      50          55          60
Leu Ser Thr Leu His Thr Leu Thr Tyr Gly Trp Thr Arg Ala Phe Glu
      65          70          75          80
Glu Ser Arg Tyr Lys Phe Tyr Leu Pro Pro Thr Phe Thr Xaa Thr Leu
      85          90          95
Leu Val Pro Cys Val Arg Ser Ser Trp Ala Lys Ala Leu Phe Xaa Leu
      100          105          110
Pro Cys Ile Gln Pro Xaa
      115

```

<210> 16

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> primer

<400> 16

```

ttaggacaac ttgatcacca gca      23

```

<210> 17

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> primer

<400> 17

```

tgtccagtcc aaactggggt attt      24

```

<210> 18

<211> 23

<212> DNA

<213> Artificial Sequence

<220>  
 <223> primer  
  
 <400> 18  
 agggagttca gcttcgttca gtc 23  
  
 <210> 19  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 19  
 ggtagaactt gtagcggctc tcct 24  
  
 <210> 20  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 20  
 gactgagctg gaactggaat ttgt 24  
  
 <210> 21  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 21  
 tttgaggaga cttcatctca ctgg 24  
  
 <210> 22  
 <211> 15  
 <212> PRT  
 <213> Homo Sapiens  
  
 <400> 22  
 Trp Lys Met Lys Pro Arg Arg Asn Leu Glu Glu Asp Asp Tyr Leu  
 1 5 10 15  
  
 <210> 23  
 <211> 14  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 23  
 ttttgatcaa gctt 14  
  
 <210> 24  
 <211> 42  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer  
  
 <400> 24  
 ctaatacgac tcactatagg gctcgagcgg ccgcccgggc ag 42  
  
 <210> 25  
 <211> 12  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 25  
 ggcccgtcct ag 12  
  
 <210> 26  
 <211> 40  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 26  
 gtaatacgac tcactatagg gcagcgtggt cgcggccgag 40  
  
 <210> 27  
 <211> 10  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 27  
 cggctcctag 10  
  
 <210> 28  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 28  
 ctaatacgac tcactatagg gc 22  
  
 <210> 29  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 29  
 tcgagcggcc gcccgggcag ga 22  
  
 <210> 30  
 <211> 20



<212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 30  
 agcgtggtcg cggccgagga 20  
  
 <210> 31  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 31  
 atatcgccgc gctcgtcgtc gacaa 25  
  
 <210> 32  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 32  
 agccacacgc agctcattgt agaagg 26  
  
 <210> 33  
 <211> 15  
 <212> PRT  
 <213> Homo Sapiens  
  
 <220>  
 <223> primer  
  
 <400> 33  
 Tyr Gln Gln Val Gln Gln Asn Lys Glu Asp Ala Trp Ile Glu His  
 1 5 10 15  
  
 <210> 34  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> FLAG sequence  
  
 <400> 34  
 gattacaagg atgacgacga taag 24  
  
 <210> 35  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> primer  
  
 <400> 35  
 acaagagcca cctctgggtg aa 22

<210> 36  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> primer

<400> 36  
 agttgagcga gtttgcaatg gac

23

<210> 37  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> IgA protease cleavage site

<400> 37  
 cctcgacctc caacaccggg g

21

<210> 38  
 <211> 22  
 <212> PRT  
 <213> Homo Sapiens

<400> 38  
 Arg Glu Val Ile His Pro Leu Ala Thr Ser His Gln Gln Tyr Phe Tyr  
 1 5 10 15  
 Lys Ile Pro Ile Leu Val  
 20

<210> 39  
 <211> 34  
 <212> PRT  
 <213> Homo Sapiens

<400> 39  
 Arg Arg Ser Tyr Arg Tyr Lys Leu Leu Asn Trp Ala Tyr Gln Gln Val  
 1 5 10 15  
 Gln Gln Asn Lys Glu Asp Ala Trp Ile Glu His Asp Val Trp Arg Met  
 20 25 30  
 Glu Ile

<210> 40  
 <211> 15  
 <212> PRT  
 <213> Homo Sapiens

<400> 40  
 Trp Ile Asp Ile Lys Gln Phe Val Trp Tyr Thr Pro Pro Thr Phe  
 1 5 10 15

<210> 41  
 <211> 26  
 <212> PRT  
 <213> Homo Sapiens

<400> 41  
 Tyr Ser Phe Val Arg Asp Val Ile His Pro Tyr Ala Arg Asn Gln Gln  
 1 5 10 15

Ser Asp Phe Tyr Lys Ile Pro Ile Glu Ile  
 20 25

<210> 42  
 <211> 33  
 <212> PRT  
 <213> Homo Sapiens

<400> 42  
 Arg Arg Ser Glu Arg Tyr Leu Phe Leu Asn Met Ala Tyr Gln Gln Val  
 1 5 10 15  
 His Ala Asn Ile Glu Asn Ser Trp Asn Glu Glu Glu Val Trp Arg Ile  
 20 25 30  
 Glu

<210> 43  
 <211> 12  
 <212> PRT  
 <213> Homo Sapiens

<400> 43  
 Lys Arg Ala Phe Glu Glu Glu Tyr Tyr Arg Phe Tyr  
 1 5 10

<210> 44  
 <211> 24  
 <212> PRT  
 <213> Homo Sapiens

<400> 44  
 Tyr Pro Tyr Val Tyr Glu Lys Lys Asp Asn Thr Phe Arg Met Ala Ile  
 1 5 10 15  
 Ser Ile Pro Asn Arg Ile Phe Pro  
 20

<210> 45  
 <211> 34  
 <212> PRT  
 <213> Homo Sapiens

<400> 45  
 Tyr Tyr Val Arg Trp Arg Leu Gly Asn Leu Thr Val Thr Gln Ala Ile  
 1 5 10 15  
 Leu Lys Lys Glu Asn Pro Phe Ser Thr Ser Ser Ala Trp Leu Ser Asp  
 20 25 30  
 Ser Tyr

<210> 46  
 <211> 13  
 <212> PRT  
 <213> Homo Sapiens

<400> 46  
 Ala Leu Gln Leu Gly Pro Lys Asp Ala Ser Arg Gln Val  
 1 5 10

<210> 47  
 <211> 14  
 <212> PRT  
 <213> Homo Sapiens

<400> 47

Ile Glu Asn Ser Trp Asn Glu Glu Glu Val Trp Arg Ile Glu  
1 5 10